Neuroplasticity Mediated by Altered Gene Expression

Colleen A McClung¹ and Eric J Nestler^{*,1}

¹Department of Psychiatry and Center for Basic Neuroscience, UT Southwestern Medical Center, Dallas, TX, USA

Plasticity in the brain is important for learning and memory, and allows us to respond to changes in the environment. Furthermore, long periods of stress can lead to structural and excitatory changes associated with anxiety and depression that can be reversed by pharmacological treatment. Drugs of abuse can also cause long-lasting changes in reward-related circuits, resulting in addiction. Each of these forms of long-term plasticity in the brain requires changes in gene expression. Upon stimulation, second messenger pathways are activated that lead to an enhancement in transcription factor activity at gene promoters. This stimulation results in the expression of new growth factors, ion channels, structural molecules, and other proteins necessary to alter the neuronal circuit. With repeated stimulation, more permanent modifications to transcription factors and chromatin structure are made that result in either sensitization or desensitization of a circuit. Studies are beginning to uncover the molecular mechanisms that lead to these types of long-term changes in the brain. This review summarizes some of the major transcriptional mechanisms that are thought to underlie neuronal and behavioral plasticity. *Neuropsychopharmacology Reviews* (2008) **33**, 3–17; doi:10.1038/sj.npp.1301544; published online 29 August 2007

Keywords: addiction; depression; transcription; learning and memory

INTRODUCTION

Neuroplasticity refers to the brain's ability to adapt and change over time. This plasticity underlies all types of memory formation (Barco *et al*, 2006). These can be memories of places, individuals or events. In addition, plasticity occurs in reward- and stress-related centers of the brain, so that one can remember and seek out reinforcing stimuli that are advantageous to survival and avoid potentially dangerous situations. However, drugs of abuse can hijack these circuits and induce abnormally persistent drug reward-related memories and in the extreme addiction (Hyman *et al*, 2006—review of the memory forming processes associated with addiction). Likewise, exposure to physical or emotional stresses can lead to depression or other mood or anxiety disorders (Duman, 2002; Nestler *et al*, 2002). Antidepressants and psychotherapeutic treatments can reverse these stress-related changes, and this is thought to mediate their efficacy. Therefore, long-term plasticity in the brain is highly relevant to both normal learning and memory, and the development and treatment of psychiatric disorders.

Work from diverse fields has suggested that common molecular and cellular mechanisms underlie aspects of all of these types of plasticity, and that long-term plasticity in the

brain requires changes in gene expression. The most well-characterized potential cellular mechanisms of plasticity are long-term potentiation (LTP) and long-term depression (LTD) (Malenka and Bear, 2004—a review of what is known about the regulation of LTP and LTD and the role of these events in memory formation). LTP and LTD describe long-lasting changes in the efficiency of synaptic transmission that occur in response to repeated stimulation. The diverse signal transduction pathways implicated in mediating LTP and LTD in different brain regions have been reviewed recently (Malenka and Bear, 2004). Both LTP and LTD appear to be important in the development of all types of memory formation including the reward-related memories associated with drug addiction (Hyman *et al*, 2006). The first indication that new gene expression is required for long-term plasticity came from several studies showing that LTP does not persist when animals are injected with protein synthesis inhibitors (Krug *et al*, 1984; Stanton and Sarvey, 1984—one of the first studies to show that LTP in the hippocampus is dependent on new gene expression; Montarolo *et al*, 1988; Otani *et al*, 1992). These drugs also have severe effects on memory formation, reactivation, and reconsolidation, suggesting that new protein expression is necessary for these processes as well (Davis and Squire, 1984; Rudy *et al*, 2006). Although it has long been thought that protein synthesis was only needed for the induction of the late phase of LTP or LTD, a recent study by Fonseca *et al* (2006) shows that early LTP can also be dependent on protein synthesis when there are high levels of synaptic activation, and that the requirement for new protein synthesis can extend well beyond LTP induction, depending

*Correspondence: Dr EJ Nestler, Department of Psychiatry, Center for Basic Neuroscience, UT Southwestern Medical Center, 5323 Harry Hines Blvd, Dallas, TX 75390-9070, USA, Tel: +214 648 1111, Fax: +214 648 4947, E-mail: Eric.Nestler@utsouthwestern.edu
Received 23 April 2007; revised 26 June 2007; accepted 22 July 2007

on the degree of synaptic stimulation. Therefore, the regulation of new gene expression in response to stimulation may be important in most aspects of LTP and LTD.

TRANSCRIPTION FACTORS AND PLASTICITY

Gene expression is controlled by a series of DNA-binding proteins known as transcription factors. With cellular stimulation, modifications are made to transcription factors that allow nuclear entry, change protein stability, enhance DNA binding, or allow binding to essential co-factors. Several studies have identified specific transcription factors that are important in the persistence of LTP, LTD, and other forms of neural and behavioral plasticity.

A Role for CREB

The most well-studied transcription factor in the context of neural plasticity is the cAMP response element-binding protein (CREB). CREB binds to the cAMP response element (CRE) in many gene promoters, including several growth factors, enzymes, structural proteins, and other transcription factors (Lonze and Ginty, 2002—an overview of CREB function and CREB family members and how they regulate various processes). CREB can be modified in multiple ways leading to changes in activity or stability of the protein. For example, it can be phosphorylated at multiple sites, acetylated by CREB-binding protein (CBP), ubiquitinated and targeted for degradation, glycosylated, or stabilized through SUMOylation (Johannessen *et al*, 2004). CREB can also interact with at least 30 other proteins, all of which affect the ability of CREB to influence gene expression (Johannessen *et al*, 2004). Upon stimulation, CREB can be activated via phosphorylation at Serine-133 by several signal transduction cascades involving protein kinase A (PKA), Ca^{2+} /calmodulin (CaM)-kinases, and growth-factor related kinases (eg ribosomal S6 kinase, which is down-

stream of extracellular signal-regulated kinases (ERK)) (Figure 1) (Mayr and Montminy, 2001; Lonze and Ginty, 2002). CREB can also be phosphorylated at Ser-142 by CaMKII, which generally leads to a decrease in activity, however, positive effects of this phosphorylation on transcription have been observed as well (Johannessen *et al*, 2004). In addition, CREB can be phosphorylated at Ser-129 by glycogen synthase kinase β (GSK-3 β); the function of this event is still unclear, but it may reduce the activity of CREB at certain gene promoters and enhance activity at others (Salas *et al*, 2003; Hansen *et al*, 2004; Johannessen *et al*, 2004).

Late-phase LTP at cortical synapses leads to an increase in the Ser-133 phosphorylation of CREB (Hottel *et al*, 2007) and an increase in CREB activity is observed in the hippocampus following training in certain tasks, or memory provoking stimuli (Bito *et al*, 1996; Taubenfeld *et al*, 1999). Furthermore, in hippocampal neuronal cultures, activation of NMDA receptors increases Ser-133 CREB phosphorylation, whereas chronic cellular stimulation by bicuculline further increases this NMDA-dependent phosphorylation of CREB (Ehlers, 2003). There is evidence to suggest that multiple signal transduction cascades are required for full activation of CREB. Consequently, CREB-mediated gene expression can be different depending on the strength of the stimulation (Johannessen *et al*, 2004). In addition, activation of certain signal transduction cascades leads to a prolonged Ser-133 phosphorylation of CREB, producing more long-lasting changes in gene expression, while others produce only a transient phosphorylation (Johannessen *et al*, 2004). This prolonged Ser-133 phosphorylation of CREB is seen with stimulation of the ERK pathway, one that is required for LTP maintenance and normal memory (Atkins *et al*, 1998; Schafe *et al*, 2000; Wu *et al*, 2001; Deisseroth and Tsien, 2002). Expression of a constitutively active form of CREB (VP16-CREB) in the CA1 region of the hippocampus lowers the threshold for eliciting the persistent late-phase LTP (Barco *et al*, 2002; Barco *et al*, 2005),

Figure 1 Regulation of CREB activity. Stimulation of D1 dopamine receptors and glutamate receptors on striatal neurons activates several second messenger cascades. Shown in the cell nucleus is a model of binding sites from the *cFos* promoter including a serum response element (SRE), activator protein-1 element (AP-1), and a cyclic AMP (cAMP) response element (CRE). CBP, CREB-binding protein; CREB, cAMP response element binding protein; MAPK, MAP kinase; NMDAR, NMDA receptor; PKA, protein kinase A; TBP, TATA binding protein. Modified from Hyman *et al* (2006).

and promotes NMDA receptor function (Marie *et al*, 2005). Some of these effects are dependent on the CREB target gene, *brain-derived neurotrophic factor* (BDNF) (Barco *et al*, 2005). Moreover, inducible overexpression of CREB in *Drosophila* enhances long-term memory, and overexpression of CREB in the amygdala of rodents leads to enhanced fear memory (Yin *et al*, 1995—one of the original studies implicating CREB in the formation of long-term memory; Josselyn *et al*, 2001; Jasnow *et al*, 2005). Likewise, genetic disruptions of CREB activity in several animal species leads to severe deficits in learning and memory (Yin and Tully, 1996; Mayford and Kandel, 1999; Josselyn *et al*, 2004). Therefore, CREB appears to be central in the formation of long-term memory.

CREB activity is also increased in reward-related areas of the brain following acute or chronic treatment with opiate drugs (such as morphine and heroin) or psychostimulants (such as amphetamine and cocaine) (Konradi *et al*, 1994; Hyman *et al*, 1995; Shaw-Lutchman *et al*, 2002; Shaw-Lutchman *et al*, 2003; Walters *et al*, 2003; Carlezon *et al*, 2005; Olson *et al*, 2005). This induction of CREB activity appears to be involved in regulating reward-related measures associated with addiction. Increasing CREB levels by the use of inducible transgenic mice or viral-mediated gene transfer in the nucleus accumbens (NAc), a crucial brain reward region, decreases the conditioned preference for these drugs, whereas blocking CREB function through expression of a dominant-negative protein has the opposite effect (Carlezon *et al*, 1998; Barrot *et al*, 2002; McClung and Nestler, 2003). Studies of constitutive CREB knockdown mice generally support these findings (Walters and Blendy, 2001). Furthermore, induction of CREB increases the excitability of GABAergic NAc neurons, whereas the dominant-negative CREB (mCREB) decreases neuronal excitability (Dong *et al*, 2006). It is thought that these acute actions of CREB represent a feed-back mechanism in which

the increased activation of CREB by drugs of abuse in the NAc induce a decrease in dopaminergic activity, leading to drug tolerance and dependence, as manifested in dysphoria during withdrawal (Figure 2) (Carlezon *et al*, 2005). Indeed, one of the target genes of CREB, *prodynorphin*, encodes a peptide (dynorphin) that is released from NAc neuron projections in the ventral tegmental area (VTA) where it binds to κ opioid receptors on dopaminergic neurons and decreases dopamine release (Cole *et al*, 1995). A recent study by found that inhibition of CREB expression by antisense oligonucleotides in the NAc selectively reduced cocaine self-administration and increased the threshold dose of cocaine needed to reinstate self-administration after a period of withdrawal. These findings suggest that activation of CREB in the NAc, and the downregulation of reward mechanisms, can lead to increased drug self-administration and relapse, perhaps as a result of the associated dysphoria.

CREB activity in other brain regions is also involved in plasticity associated with drug addiction. The actions of CREB in the VTA are complex, in that CREB overexpression in the rostral vs caudal subregions of the VTA have opposing effects on opiate and psychostimulant preference (Walters *et al*, 2003; Olson *et al*, 2005). This may be due to differences in the proportion of dopaminergic and GABAergic neurons in these two subregions. CREB in the VTA is also implicated in nicotine reward (Walters *et al*, 2005—an important study that outlines a crucial role for CREB in the ventral tegmental area in nicotine reward). In addition to the VTA, CREB may modulate drug reward through its actions in the lateral hypothalamus (LH). CREB activity is induced in this area following chronic opiate or psychostimulant treatment (Shaw-Lutchman *et al*, 2002; Georgescu *et al*, 2003; Shaw-Lutchman *et al*, 2003; Georgescu *et al*, 2005), and CREB overexpression in the LH increases the conditioned preference for morphine,

Figure 2 Feedback between the NAc and VTA via CREB activation. Cocaine and amphetamine have been shown to activate *prodynorphin* gene expression in the nucleus accumbens (NAc) and dorsal striatum via D1 dopamine receptor stimulation, the cyclic AMP pathway, and the phosphorylation of CREB. The resulting dynorphin peptides are transported to presynaptic terminals including terminals found on recurrent collateral axons that feedback on dopaminergic neurons. Dynorphin peptides are agonists at inhibitory κ opiate receptors, resulting in decreased dopamine release. Modified from Hyman *et al* (2006).

whereas expression of a dominant-negative CREB decreases morphine preference (Olson *et al*, 2007). Together, these results show that CREB is involved in regulating drug-induced plasticity in multiple brain regions.

CREB also appears to be involved in the development and treatment of mood and anxiety disorders. However, studies showing the effects of stress on CREB activity are highly dependent on the method and duration of stress and brain region examined. For example, CREB activity is increased in the NAc following stress associated with short-term foot shock, restraint, and unfamiliar social interactions, whereas long-term social isolation decreases CREB activity in the NAc (Barrot *et al*, 2002; Barrot *et al*, 2005). Chronic, unpredictable stress also decreases Ser-133 CREB phosphorylation in the dentate gyrus of the hippocampus and the frontal cortex (Gronli *et al*, 2006; Xu *et al*, 2006). However, restraint stress increases CREB activity in the frontal cortex with no effects on the hippocampus (Miller *et al*, 2006).

Induction of CREB causes very different functional effects—either antidepressant-like or pro-depression-like effects—depending on the brain region involved (Duman, 2002; Blendy, 2006; Nestler and Carlezon, 2006). Selective overexpression of CREB in the dentate gyrus of the hippocampus has an antidepressant effect in multiple behavioral measures (Chen *et al*, 2001a). In contrast, overexpression of CREB in the amygdala or NAc has a prodepressant effect, whereas dominant-negative CREB (mCREB) expression results in an antidepressant-like response in several behavioral measures (Pliakas *et al*, 2001; Newton *et al*, 2002; Wallace *et al*, 2004). These results highlight the importance of CREB in this type of plasticity, and how the changes evoked from these types of stimuli differ throughout the brain.

Other CREB family members act as modulators of CRE-mediated transcription and contribute to the regulation of neuronal plasticity. The actions of CREB can be inhibited by the inducible cAMP early repressor (ICER) proteins, which are protein products of the *CRE modulator* gene (*Crem*) (Mioduszevska *et al*, 2003). Although overexpression of CREB in the hippocampus of older rats (15 months old) leads to an improvement in long-term memory over wild-type animals, overexpression of an ICER protein decreases long-term memory in spatial-navigation and passive avoidance tasks (Mouravlev *et al*, 2006). Furthermore, reported recently that acute exposure to either stress or psychostimulants induces ICER proteins in the NAc, and overexpression of an ICER protein in this region through viral-mediated gene transfer, like overexpression of the artificial dominant-negative mCREB, enhances responses to drugs of abuse and natural rewards, and induces an antidepressant-like effect. These results suggest that natural inhibitors of CREB are involved in modulating neural plasticity relevant to drug addiction and stress-related disorders.

A Role for Δ FosB

The Fos family of transcription factors, including cFos, FosB, Δ FosB, Fos-related antigen 1 (Fra1), and Fra2, dimerize with Jun proteins to form an AP-1 transcription factor complex (Hess *et al*, 2004). LTP stimulation in the

dentate gyrus leads to an increase in AP-1 DNA binding, 4 h after stimulation, suggesting that this enhanced binding is associated with late-phase LTP and long-term plasticity (Williams *et al*, 2000). All Fos family proteins are induced rapidly in many brain regions following various types of acute stimulation. With more chronic treatments, this induction becomes less and less prominent. However, Δ FosB, a splice variant of the *FosB* gene, is an unusually stable protein that accumulates with chronic treatments (McClung *et al*, 2004—a recent review of the actions and regulation of Δ FosB). As this protein is so persistent, it is thought that it is important in the induction and maintenance of long-term plasticity. Indeed, Δ FosB is induced by a wide range of chronic stimuli, including repeated exposure to many types of drugs of abuse or stress, in specific brain regions (McClung *et al*, 2004; Perrotti *et al*, 2004). Δ FosB also appears to have different functions over time, in that, it acts much like a transcriptional repressor at AP-1 sites with short-term treatments, but then functions as a transcriptional activator, as it accumulates with more chronic treatments (McClung and Nestler, 2003).

When the FosB gene is knocked out, resulting in the absence of both FosB and Δ FosB, mice have behavioral abnormalities in response to drugs of abuse, and in measures of anxiety and depression, that are more pronounced when mice are stressed, suggesting that this gene is important in the regulation of these responses (Zhu *et al*, 2007). Long-term overexpression specifically of Δ FosB in the NAc mimics an overall addiction-like phenotype seen in response to drugs of abuse (Kelz *et al*, 1999; Colby *et al*, 2003; Zachariou *et al*, 2006). Conversely, the expression of a dominant-negative c-Jun, termed Δ c-Jun, which acts as a repressor of AP-1 activity, in NAc opposes this addiction-like phenotype (Peakman *et al*, 2003; Zachariou *et al*, 2006). Long-term Δ FosB expression in the NAc also leads to enhanced food-reinforced instrumental performance and an increase in voluntary wheel running, suggesting that it is important in the response to natural as well as drug rewards (Werme *et al*, 2002; Olausson *et al*, 2006). Recently, found that Δ FosB is phosphorylated by casein kinase 2 (CK2) at Serine 27, and that this phosphorylation opposes protein degradation, contributing to Δ FosB's unusual stability. Furthermore, this phosphorylation event increases the transcriptional activity of Δ FosB (Ulery and Nestler, 2007). As it has been shown that LTP induction rapidly increases the activity of CK2 in the hippocampus (Chariat-Marlangue *et al*, 1991), it is possible that this provides a mechanism by which the activity of Δ FosB is increased and stabilized following chronic stimulation, leading to some of the long-term gene expression changes responsible for plasticity. Δ FosB's unusual stability is also due to the fact that it lacks two degron domains present in the C-terminus of full-length FosB and all other Fos family proteins, which target these proteins for rapid degradation (Carle *et al*, 2007).

A Role for NF- κ B

The transcription factor, NF- κ B, can exist as a latent form or a constitutively active form in neurons (Kaltschmidt *et al*, 2005). Although the constitutive form is located in the nucleus, the latent form is held in the cytoplasm through an

interaction with inhibitory kappaB ($I\kappa B$). Upon stimulation by numerous factors, $I\kappa B$ is degraded, allowing NF- κB to enter the nucleus. In addition, the RelA subunit of NF- κB can be acetylated upon stimulation and this prevents its interaction with $I\kappa B\alpha$ (Chen *et al*, 2001b). RelA can subsequently be deacetylated by histone deacetylase 3 (HDAC3), and this promotes nuclear export and $I\kappa B\alpha$ binding (Chen *et al*, 2001b). This reversible acetylation contributes to the duration of NF- κB activity in the nucleus. NF- κB is activated in response to neurodegeneration, cellular stress, or trauma, and through its regulation of antiapoptotic genes, it promotes neuronal survival (Mattson, 2005). NF- κB activity is also induced following LTP in the hippocampus and after fear conditioning in the amygdala (Yeh *et al*, 2002; Romano *et al*, 2006). Studies have found that the administration of a competing decoy κB can prevent LTD and severely impair LTP in both the hippocampus and amygdala (Albensi and Mattson, 2000; Yeh *et al*, 2002). In addition, members of the NF- κB family, p50, p65, and c-Rel, show an increase in DNA binding in the hippocampus following glutamate receptor stimulation, and mice lacking c-Rel have a deficit specifically in the late-phase of LTD (O'Riordan *et al*, 2006). Furthermore, animals that overexpress a 'super repressor' $I\kappa B$ -AA in neurons selectively using a bitransgenic system have defects in learning and memory (Kaltschmidt *et al*, 2006). The late phase of hippocampal LTP in these mice is also compromised and LTD can not be induced (Kaltschmidt *et al*, 2006). Interestingly, this study identified the PKA catalytic α -subunit as a transcriptional target for NF- κB , and the loss of NF- κB function reduces PKA-mediated CREB phosphorylation (Kaltschmidt *et al*, 2006). Thus, the plasticity and memory defects in the $I\kappa B$ -AA overexpressing mice may come from a lack of activated CREB.

In addition to the effects on memory, NF- κB is also induced by chronic cocaine in the NAC, where it may be a transcriptional target of $\Delta FosB$ (Ang *et al*, 2001). Therefore, NF- κB may also be involved in the plasticity associated with addiction, a possibility, which now requires further investigation.

A Role for Circadian Transcription Factors

Recent studies have implicated the transcription factors that make up the circadian clock in neural plasticity. Circadian rhythms, or 24-h cycles, occur in many bodily functions, including sleep/wake, body temperature, and hormone levels. Circadian rhythms are largely regulated by a transcriptional feed-back loop located in the suprachiasmatic nucleus (SCN) in the hypothalamus. However, the genes that make up the clock are expressed throughout the brain and other organs, and can form SCN-independent oscillators that respond to many types of external stimuli (King and Takahashi, 2000—a review of the circadian clock and rhythms in mammals; Abe *et al*, 2001; Reppert and Weaver, 2001; Stokkan *et al*, 2001; Iijima *et al*, 2002; Granados-Fuentes *et al*, 2006; McDearmon *et al*, 2006; Mieda *et al*, 2006). In fact, it is estimated that >10% of all mammalian messenger RNA (mRNA) transcripts are regulated in a circadian manner (Akhtar *et al*, 2002; Duffield *et al*, 2002).

The major transcriptional activator consists of a dimer between the Circadian Locomotor Output Cycles Kaput Protein (CLOCK) and Brain and Muscle ARNT-like Protein 1 (BMAL1, also known as ARNTL or MOP3). This complex binds to E-box sequences in the promoters of many genes, including the *Period* (*Per*) and *Cryptochrome* (*Cry*) genes. The PER and CRY proteins are translated in the cytoplasm, and are phosphorylated by casein kinase I ϵ and δ (CK1) and GSK3 β , leading to changes in their stability, association, and nuclear entry (Harms *et al*, 2003; Iitaka *et al*, 2005; Knippschild *et al*, 2005; Kurabayashi *et al*, 2006). These proteins can then repress the actions of CLOCK:BMAL1, which creates a precisely timed transcriptional feedback loop, the molecular basis of circadian rhythms. In addition, there is an adjoining loop in which CLOCK:BMAL1 activates the transcription of *Rev-erba* and *Rora* (Sato *et al*, 2004; Guillaumond *et al*, 2005). These proteins can bind to the promoter of the *Bmal1* gene and both positively and negatively regulate its transcription. Selectively in forebrain regions, Neuronal PAS Domain Protein 2 (NPAS2), a protein very similar to CLOCK, can bind BMAL1 and induce *Per* and *Cry* gene expression independent of CLOCK (Reick *et al*, 2001). NPAS2 may also function in the place of CLOCK in the SCN, if the CLOCK protein is abolished (Debruyne *et al*, 2006).

It is thought that sleep contributes strongly to the process of memory formation and consolidation (Walker and Stickgold, 2006). In addition, phase shifting of circadian rhythms or chronic disruption of circadian rhythms can impair long-term memory and working memory in both rodents and humans (Cho *et al*, 2000; Devan *et al*, 2001). Several studies have found differences in learning and memory when experiments are performed during the light cycle vs the dark cycle. For example, snails display a diurnal regulation of conditioned taste aversion (Wagatsuma *et al*, 2004). Rodents also show diurnal differences in contextual fear conditioning and performance in the Morris water maze (Chaudhury and Colwell, 2002; Valentinuzzi *et al*, 2004). In addition, the magnitude of hippocampal LTP is greater and the decay of LTP is slower during the night in nocturnal mice, even when the mice are kept in constant darkness, suggesting that an endogenous circadian oscillator regulates hippocampal plasticity (Chaudhury *et al*, 2005). Studies in *Aplysia* found that long-term memory formation in response to electrical stimulation or 5-HT is suppressed in diurnal animals that were trained during the night (Fernandez *et al*, 2003; Lyons *et al*, 2005, 2006). This reduction in memory formation during the night was correlated with a circadian rhythm in the induction of the memory-related proteins, phosphorylated ERK, and CCAAT/enhancer-binding protein, such that a greater induction of these proteins was seen when training was performed during the day than at night (Lyons *et al*, 2006). Long-term memory formation at night could be restored to day-time levels when animals were treated with agents that activate ERK and another transcriptional activator together, but not individually, indicating that both ERK phosphorylation and general gene transcription are necessary for normal memory formation, and that both are inhibited by the circadian clock during the night.

Additional studies have implicated individual members of the circadian clock or clock-controlled hormones in

learning and memory. *Drosophila* that carry a mutation in the *Per* gene are defective in long-term memory associated with experience-dependent courtship, whereas overexpression of *Per* enhanced memory formation in this paradigm (Sakai *et al*, 2004). Interestingly, mutations in other circadian genes including *Clock* and *Cycle* did not affect memory formation in this task, suggesting an independent role for *Per* outside of the central clock (Sakai *et al*, 2004). Somewhat surprisingly, mice that lack functional *Per1* or *Per2* show normal spatial and contextual learning in hippocampal-dependent tasks (Zueger *et al*, 2006), however, mice lacking both genes have not been tested. The circadian hormone, melatonin, which is released at night in response to signals from the SCN, can inhibit hippocampal LTP (Wang *et al*, 2005). As well, expression of the circadian-controlled transcription factor, albumin D-element-binding protein (DBP), is suppressed by hippocampal activation of a protein associated with increased learning, glucagon-like peptide-1 receptor, and viral-mediated overexpression of DBP in the hippocampus leads to an inhibition of spatial learning (Klugmann *et al*, 2006). Moreover, induction of LTP leads to the phosphorylation and inhibition of the circadian modulator, GSK-3 β in the hippocampus (Hooper *et al*, 2007). LTP is impaired in mice that conditionally overexpress GSK-3 β and the deficits can be rescued by treatment with the GSK-3 β inhibitor, lithium (Hooper *et al*, 2007). These results suggest that certain components of the molecular clock are involved in learning and memory.

Several mood and substance abuse disorders have also been strongly linked to the circadian clock (Bunney and Bunney, 2000; Mansour *et al*, 2005; Yuferov *et al*, 2005; Wirz-Justice, 2006). Indeed, some of the most pronounced symptoms of diseases like major depressive disorder (MDD) and bipolar disorder (BPD) are abnormal sleep/wake, appetite, and social rhythms (Boivin, 2000; Bunney and Bunney, 2000; Lenox *et al*, 2002; Grandin *et al*, 2006). In addition, nearly all of the successful treatments for mood disorders affect circadian rhythms, and it appears that the shifts, resetting, and stabilization of these rhythms, produced by these treatments, are important for their therapeutic efficacy (McClung, 2007). Interestingly, a recent study found that the circadian genes, *Clock*, *Npas2*, and *Bmal1*, are all strongly induced in the mouse hippocampus after chronic, but not acute, treatment with the antidepressant, fluoxetine (Uz *et al*, 2005). This suggests that these genes may be involved in the plasticity that occurs in this region with chronic antidepressant treatment. In addition, chronic treatment with the mood stabilizer, valproate, decreased the expression of *CK1 δ* and *Cry2* in the amygdala (Ogden *et al*, 2004). These changes did not occur when mice were co-treated with methamphetamine, which was given to induce manic-like symptoms, suggesting that these genes may be involved in the treatment of mania (Ogden *et al*, 2004).

Researchers have begun to examine the roles of individual members of the circadian clock in the regulation of mood and drug addiction. Transgenic mice that overexpress GSK-3 β are hyperactive, have reduced immobility in the forced swim test (interpreted as lower depression-like behavior), and an increased startle response (Prickaerts *et al*, 2006). These behaviors are reminiscent of those of bipolar patients in the manic state. This is perhaps not surprising, as

GSK-3 β is inhibited by the mood stabilizer, lithium (Jope and Roh, 2006). Recently, Roybal *et al* (2007)—the authors report a mania-like phenotype, and its reversal by chronic lithium treatment, in mice lacking functional Clock protein—found that mice harboring a mutation in the *Clock* gene also display a behavioral profile that is strikingly similar to human mania. These mice have a point mutation in the *Clock* gene that results in the expression of a dominant-negative protein (King *et al*, 1997). These mice are extremely hyperactive in response to novelty and throughout the light/dark cycle, they have reduced depression-like behavior in the forced swim and learned helplessness tests, reduced anxiety or increased risk-taking behavior in several measures, and an increase in the reward value of cocaine, sucrose, and intracranial self-stimulation (McClung *et al*, 2005; Roybal *et al*, 2007). In addition, other groups have found that these mice sleep less and have increased exploratory activity, adding to their overall manic-like phenotype (Naylor *et al*, 2000; Easton *et al*, 2003). Interestingly, when these mice were treated chronically with lithium, the majority of their behavioral responses return to wild-type levels (Roybal *et al*, 2007). As the midbrain dopaminergic system has been widely implicated in the development of mania, we performed *in vivo* recordings from the dopaminergic neurons of the VTA in the *Clock* mutant mice, and found an overall increase in dopamine cell firing and bursting (McClung *et al*, 2005; Nestler, 2005; Nestler and Carlezon, 2006). CLOCK is expressed in the VTA, where it regulates several genes that are important in dopaminergic transmission (McClung *et al*, 2005). Therefore, we used viral-mediated gene transfer to express a functional CLOCK protein specifically in the VTA of the mutant mice, and found that this treatment returned several of their behavioral phenotypes to wild-type levels (Roybal *et al*, 2007). These results suggest that CLOCK is important in the development of mania, and that at least a portion of the behavioral and mood-related phenotypes seen in the mutant mice are due to the loss of CLOCK function in the VTA.

As mentioned above, one of the phenotypes of the *Clock* mutant mice is an increase in the reward value for cocaine. This was measured both by conditioned place preference and by levels of intracranial self-stimulation following cocaine treatment (McClung *et al*, 2005; Roybal *et al*, 2007). Other members of the circadian clock have also been implicated in drug reward and addiction. The original studies implicating circadian genes in drug-induced plasticity found that *Drosophila* lacking *Per*, *Clock*, *Cycle*, or *Doubletime* all fail to sensitize to cocaine (Andretic *et al*, 1999—this is the first study to demonstrate a role for the genes that control circadian rhythms in the regulation of cocaine responsiveness). Since then, several studies have found changes in the expression of multiple circadian genes in various brain regions of rodents treated with drugs of abuse (Nikaido *et al*, 2001; Ammon *et al*, 2003; Yuferov *et al*, 2003, 2005; McClung *et al*, 2005; Uz *et al*, 2005; Manev and Uz, 2006). In addition, mice that lack a functional *Per1* gene show reduced behavioral responses to cocaine (Abarca *et al*, 2002). Reduction in the expression of *Per1* by DNase targeting likewise leads to a reduction in the conditioned preference for morphine (Liu *et al*, 2005). However, mice lacking *Per1* show normal levels of alcohol

self-administration and reinstatement of alcohol-seeking behavior, suggesting that *Per1* is perhaps not involved in alcohol reinforcement (Zghoul *et al*, 2007). Interestingly, mice lacking *Per2* show an enhanced sensitization to cocaine, but normal levels of conditioned preference for cocaine (Abarca *et al*, 2002). They also show an increase in alcohol self-administration and enhanced incentive motivation for alcohol (Spanagel *et al*, 2005). These results suggest that *Per1* and *Per2* may serve different functions in response to treatment with drugs of abuse.

CHROMATIN MODIFICATIONS THAT UNDERLIE PLASTICITY

Modifications to Histones

Long-term modifications to chromatin may underlie some of the changes in gene expression that lead to neural plasticity. DNA is tightly packed around octamers of the histone proteins H2A, H2B, H3, and H4, linked together by histone H1 (Berger, 2001). Histone proteins can be modified to allow the DNA to unwind and permit transcription factor binding and gene activation (Figure 3). In turn, other modifications to histone proteins can inhibit transcription factor binding. Several studies have found that gene activation is associated with histone acetylation (Cheung *et al*, 2000). This acetylation is catalyzed by a histone acetyltransferase (HAT) enzyme, such as CBP/p300 or many others, which serve as transcriptional co-activators. In addition, some transcription factors appear to have their own HAT domain that may allow histone acetylation. For example, the CLOCK protein has recently been shown to contain an active HAT domain that is enhanced by BMAL1 binding (Doi *et al*, 2006). HDAC proteins reduce gene expression at these promoters through histone deacetylation. Histone phosphorylation is also generally associated with an increase in transcription, and phosphorylation has

been associated with immediate early gene activation in response to growth factor treatment (Mahadevan *et al*, 1991; Sassone-Corsi *et al*, 1999).

Histones H3 and H4 can also be methylated at arginine or lysine residues, which can lead to gene activation or repression depending on the residue undergoing methylation (Beaujean, 2002). The arginine methyltransferase, co-activator-associated arginine methyltransferase-1, can co-activate myogenic transcription factor-mediated gene expression (Chen *et al*, 2002). However, methylation of histone H3 on lysine residues K9 or K27 is associated with stretches of transcriptional repression and inert heterochromatin (Beaujean, 2002). Finally, ubiquitination, SUMOylation, and ADP-ribosylation of histones have been observed, and these processes influence gene transcription as well (Huletsky *et al*, 1985; Davie and Murphy, 1990; Beaujean, 2002; Nathan *et al*, 2003).

HAT and HDAC proteins can associate with transcription factors, and this association often occurs in response to second messenger stimulation. For example, CREB binds CBP after phosphorylation at Ser-133 (Johannessen *et al*, 2004). Mice that lack full CBP function through various genetic methods have defects in late-phase LTP, spatial learning, and fear conditioning (Alarcon *et al*, 2004; Korzus *et al*, 2004; Wood *et al*, 2005). Furthermore, studies have found that the induction of LTP in hippocampal slices using high-frequency stimulation is enhanced by treatment with the nonspecific HDAC inhibitors, trichostatin A and sodium butyrate (Levenson *et al*, 2004). Trichostatin A also enhanced forskolin-induced LTP in the amygdala (Yeh *et al*, 2004). These results suggest that HDAC inhibition and transcriptional activation may improve the plasticity associated with memory formation.

Long-term modifications to histone proteins have been seen in response to chronic stimuli, where they contribute to neuronal plasticity. Early studies in *Aplysia* found that acetylation of histone H4 at the ApC/EBP promoter was increased following 5-HT-induced long-term facilitation,

Figure 3 Differential states of chromatin. Chromatin can either be open (ie active, allowing gene expression) or condensed (ie inactive, repressing gene expression). This change in state is mediated by the modifications to core histone proteins. Histone acetylation (A) is associated with chromatin relaxation and the binding of transcription factors and co-activators, such as HATs (histone acetyl transferases) and SWI-SNF proteins that mediate the movement of nucleosomes along a strand of DNA. Histone methylation (M) results in condensed chromatin and transcriptional repression (REP).

and acetylation was decreased following FMRFamide-induced LTD through the recruitment of HDAC5 (Guan *et al*, 2002). Later, it was found that multiple signaling pathways modify histones in the rodent hippocampus, including an increase in H3 acetylation following the activation of NMDA receptors and the ERK pathways associated with LTP (Crosio *et al*, 2003—an important paper that delineates some of the intracellular messenger pathways that lead to regulation of histone acetylation; Levenson *et al*, 2004). In studies of depression, the HDAC inhibitor, sodium butyrate, has an antidepressant-like effect in mice that correlates with hyperacetylation at the *Bdnf* promoter in the frontal cortex (Schroeder *et al*, 2006; Tsankova *et al*, 2006—this is the first study to implicate histone modifications in depression- and antidepressant-like behavior). In agreement with these findings, our studies found that a chronic electroconvulsive seizure paradigm, which is antidepressant in humans, changes the acetylation of specific histones at the *Bdnf*, *Creb*, and *c-Fos* promoters in the hippocampus, and these changes correlate with sustained increases or decreases in levels of expression of these genes (Tsankova *et al*, 2004). Furthermore, a chronic social defeat stress paradigm, which produces a depression-like state (Berton *et al*, 2006), decreases the expression of *Bdnf* in the hippocampus (Tsankova *et al*, 2006). The prolonged reduction in *Bdnf* expression was correlated with an increase in repressive histone methylation at the gene promoter (Tsankova *et al*, 2006). The antidepressant, imipramine reversed the depression-associated behavior and increased histone acetylation at the *Bdnf* promoter and increased *Bdnf* expression (Tsankova *et al*, 2006). Furthermore, this increase in histone acetylation with imipramine treatment involves the downregulation of HDAC5, and overexpression of HDAC5 in the hippocampus blocks the ability of imipramine to reverse the depression-like behavior (Tsankova *et al*, 2006). Together, these studies highlight the importance of long-term histone modifications in the plasticity associated with depression and antidepressant treatment.

Chromatin remodeling through histone modification also appears to play a role in drug addiction. One of the key features of addiction is that it is more persistent if drug use begins at an early age, indicating long-lasting changes in brain function. Interestingly, a recent study found that the administration of ascending doses of cocaine to adolescent rats resulted in an increased response to cocaine in adulthood, and an overall decrease in histone H3 methylation at lysines 4 and 27 in the medial pre-frontal cortex (Black *et al*, 2006). Furthermore, acute cocaine leads to a transient increase in histone acetylation at the *c-Fos* promoter in striatal regions, consistent with its transient increase in expression following this treatment (Kumar *et al*, 2005). However, chronic cocaine treatment selectively increases *Bdnf* and *Cdk5* gene expression in these regions, and these increases are associated with hyperacetylation at their promoters (Kumar *et al*, 2005). Kumar *et al* provided several lines of evidence indicating that induction of the *Cdk5* gene is mediated via Δ FosB. Cocaine also increases CBP-mediated acetylation of histone H4 at the *FosB* promoter, and mice lacking one allele of CBP have less accumulation of Δ FosB, and are less sensitive to chronic cocaine (Levine *et al*, 2005). Furthermore, administration of

the HDAC inhibitor, TSA, increases the conditioned place preference for cocaine, whereas viral-mediated overexpression of HDAC4 in the NAc decreases cocaine preference (Kumar *et al*, 2005). These results suggest that chronic cocaine treatment leads to a hyperacetylation at several gene promoters, which increases the reward value for cocaine.

DNA Methylation

DNA (cytosine-5) methylation occurs at methyl CpG islands within a gene sequence, and it was once thought to be a permanent way by which certain genes are silenced. This silencing occurs during X-chromosome inactivation, and several cell-fate determination processes throughout development (Morgan *et al*, 2005). Recent studies, however, show that changes in DNA methylation may be more transient, and underlie certain types of long-term plasticity. For example, DNA methylation can occur in response to stimuli experienced during development or early childhood, and this can lead to lasting changes into adulthood. A recent study by Weaver *et al* (2004)—one of the first studies to show that DNA methylation patterns can be influenced during adolescence and that these patterns affect adult behavior—found that patterns of DNA methylation in the *glucocorticoid receptor* gene were determined during early life based on the quality of maternal care experienced. These patterns or ‘memories’ remained until adulthood, and influenced levels of anxiety and nurturing behavior in the rats as adults. These results suggest that the methylation patterns that occur during early childhood result in long-lasting changes in gene expression that affect adult behavior. Interestingly, these changes in adult behavior were reversible. Infusion of adult rats with the amino-acid L-methionine, a precursor to S-adenosyl-methionine that acts as the donor of methyl groups for DNA methylation, reverses the effect of maternal behavior on DNA methylation and the behavioral response to stress (Weaver *et al*, 2005). This shows that the DNA methylation changes that occur during adolescence are not permanent. Another study, by Zhang *et al* (2007) found that maternal cocaine use induces persistent DNA methylation of the *Pkcx* gene in the heart of the fetus. It is possible that similar changes could occur in the brain, leading to a propensity for drug addiction later in life.

Studies aimed at determining whether or not DNA methylation is involved in more short-term learning and memory are just beginning. Interestingly, a protein that binds methylated cytosines and represses transcription, MeCP2, has been associated with memory formation. Loss of function mutations in this protein cause Rett syndrome, a developmental disorder that severely affects cognitive and motor abilities (Williamson and Christodoulou, 2006). These behavioral abnormalities are associated with impairments in presynaptic release of glutamate (Nelson *et al*, 2006—an important study that demonstrates abnormal presynaptic function in mice lacking MeCP2). Surprisingly, overexpression of MeCP2 enhances hippocampal LTP and long-term memory formation transiently before animals develop seizures, severe hypoactivity, and other ailments (Collins *et al*, 2004). This suggests that the proper balance of MeCP2 is needed for memory function. In addition, studies by Chen *et al* (2003)—one of the first studies to

demonstrate regulation of the *Bdnf* gene by DNA methylation mechanisms—and Martinowich *et al* (2003) found that MeCP2 binds to the *Bdnf* promoter III and inhibits *Bdnf* transcription through CpG methylation. Membrane depolarization leads to a calcium-dependent phosphorylation of MeCP2 that dissociates the MeCP2-HDAC-mSin3A complex from the promoter, allowing *Bdnf* transcription (Chen *et al*, 2003; Martinowich *et al*, 2003). Furthermore, recent studies by Sweatt and co-workers found that the enzymes that catalyze the methylation of cytosine residues, DNA methyltransferases (DNMTs), are important regulators of memory in the hippocampus (Levenson *et al*, 2006; Miller and Sweatt, 2007—a recent study implicating DNA methylation in the formation of long-term memory). Inhibition of DNMT through treatment with phorbol-12,13-diacetate altered the methylation at the *reelin* and *Bdnf* promoters, and blocked the induction of LTP (Levenson *et al*, 2006). Furthermore, DNMT expression is upregulated in the hippocampus with contextual fear conditioning, and DNMT inhibition leads to deficits in memory formation (Miller and Sweatt, 2007). DNMT is involved in the silencing of the memory suppressor gene, *Pp1*, and this gene is rapidly inactivated via DNA methylation following fear conditioning (Miller and Sweatt, 2007). These studies suggest that DNA methylation is far more dynamic than previously described and could be an important mechanism that contributes to long-term neuronal plasticity.

POSTTRANSCRIPTIONAL MODIFICATIONS THAT UNDERLIE PLASTICITY

Polyadenylation Factors

In addition to the long-term modifications to transcription factors and DNA that govern neuronal plasticity, post-

transcriptional mechanisms also appear to be involved. Many mRNAs are transported to synapses where they are locally translated, and participate in synaptic remodeling at those locations. Several studies have identified mRNA-binding proteins, such as the cytoplasmic polyadenylation element-binding protein (CPEB) and cleavage and polyadenylation specificity factor that regulate mRNA translation at the synapse (Wu *et al*, 1998). *Drosophila* carrying a mutation in *orb*, a CPEB homologue, lack long-term memory (Dubnau *et al*, 2003). Furthermore, NMDA receptor stimulation induces the phosphorylation of CPEB by Aurora kinase, and disrupts its association with a repressor protein, Maskin, thereby allowing CPEB to interact with the translation initiation factors, eIF4G and eIF4E at target mRNAs (Huang *et al*, 2002; Si *et al*, 2003—a study showing the importance of postranscriptional regulation by CPEB in plasticity). This action can stabilize synaptic changes at these sites associated with plasticity. Interestingly, the *Aplysia* CPEB was reported to have prion-like properties in that, it is modular and transferable, and it is capable of conferring a change in state to other proteins (Si *et al*, 2003). The prion-like, aggregated state is the active state of CPEB, and it is possible that this conformational change comes about through chronic stimulation, leading to long-lasting, local synaptic changes.

MicroRNAs

Recently, a class of non-coding, micro RNAs (miRNA) has been described that influence the translation of mRNA by sequence-specific pairing with 3'-untranslated regions (Presutti *et al*, 2006). It is predicted that 40–50% of mRNAs are regulated by miRNAs (Presutti *et al*, 2006). A number of these miRNAs appear to be brain specific, and they have been implicated in neuronal development and differentia-

Figure 4 Regulation of mRNA translation by miRNAs. The ability of mRNAs to be locally translated and participate in synaptic plasticity can be altered by miRNAs. This interaction between the miRNA, mRNA, and the RISC complex can lead to the inhibition of translation, mRNA degradation, or the sequestering of the mRNA away from the synapse. Stimulation by glutamate, or growth factors such as BDNF, can lead to the degradation of components of the RISC complex and allow mRNA translation.

tion (Presutti *et al*, 2006). These miRNAs also undergo tissue-specific RNA editing, adding another level of potential regulation to their function (Kawahara *et al*, 2007). It is possible that miRNAs are involved in regulating the local translation of mRNAs at synapses that are involved in plasticity (Figure 4). Many of the structural changes that are thought to underlie synaptic plasticity occur at dendritic spines and branches (Schratt *et al*, 2006—an important study demonstrating the regulation of dendritic spine growth by a microRNA). miRNAs might be involved in repressing translation of proteins at these synapses under normal conditions, with this repression being abolished following stimulation. Indeed, in the hippocampus, a brain-specific miRNA, *miR-134* is localized to the synaptodendritic compartment of neurons where it decreases the size of dendritic spines through its repressive actions at the *LIM-domain kinase 1* (*Limk1*) mRNA (Schratt *et al*, 2006). Neuronal stimulation by BDNF relieves the repression of *Limk1* by *miR-134*, allowing spine growth (Schratt *et al*, 2006). Another recent study in *Drosophila* found that local protein synthesis occurs at synapses when animals are exposed to an odor paired with an electric shock (Ashraf *et al*, 2006). A component of the RNA-silencing complex (RISC), Armitage, that couples with miRNAs is located at these synapses, where it normally silences mRNAs (Ashraf and Kunes, 2006; Ashraf *et al*, 2006; Chekanova and Belostotsky, 2006). Upon stimulation, Armitage is degraded, and the mRNA is now translated. Some of the key proteins involved in plasticity are targets of this pathway, including CaMKII and Staufien (Ashraf and Kunes, 2006; Ashraf *et al*, 2006). Therefore, the RISC complex may play a key role in synaptic plasticity.

Total expression levels of the miRNAs may contribute to the maintenance of long-lasting plasticity. Interestingly, genome-wide screens of transcription factor binding found that CREB may regulate the expression of individual brain-specific miRNAs (33 had identifiable CRE regulatory sequences), and *Mir-132*, a miRNA expressed predominantly in the hippocampus, was identified as a direct target of CREB (Vo *et al*, 2005; Wu and Xie, 2006; Lukiw, 2007). Furthermore, the transcriptional repressor RE1-silencing transcription factor (also known as neuron-restrictive silencer factor), appears to convey long-term repression of some of the miRNAs by recruiting MeCP2 and HDACs to their regulatory regions (Wu and Xie, 2006). Therefore, long-term treatments could change the expression level of these miRNAs through manipulation of these transcription factors, which would result in more or less local translation of proteins that are involved in altering synaptic connections.

FUTURE RESEARCH DIRECTIONS

Although it is clear that transcriptional and post-transcriptional changes are necessary for the development of neuroplasticity, the mechanisms that underlie these changes are complex, and we are only beginning to understand how these changes are mediated. In addition, many of the transcriptional target genes that are involved in long-term plasticity have yet to be identified. Therefore, future research into the stimulus-dependent regulation of tran-

scription factors, chromatin structure, and mRNA processing are important in our understanding of how long-term changes are facilitated in the brain. The use of new technologies such as chromatin immunoprecipitation (ChIP) followed by microarray analysis (ie ChIP on chip) or DNA methylation arrays will allow us to study chromatin modifications that occur following various treatments, and identify a range of direct transcriptional targets of many transcription factors involved in plasticity. In addition, the study of miRNAs not only allows us to determine their importance in plasticity but also provides a technique that is now being utilized to knock down the expression of specific genes in individual brain regions of adult animals to determine their function in these long-term adaptations.

CLINICAL IMPLICATIONS

This research will undoubtedly lead to new, more targeted, treatments for memory disorders, as well as drug addiction, depression, and other psychiatric conditions. Through better understanding of the mechanisms that underlie long-term plasticity, it might be possible to either enhance or block specific aspects of a form of plasticity depending upon the condition. For example, it may be possible to reverse the chromatin modifications that occur in response to chronic exposure to drugs of abuse by modulating the transcription factors or enzymes necessary to facilitate this process. As all forms of plasticity appear to be mediated via a finite number of basic molecular processes, the large task ahead is to determine the specific molecules, changes, and circuits that are involved in the formation of normal memories, as opposed to drug addiction or depression. It is likely, therefore, that future treatments for these disorders will have to be highly targeted toward specific adaptations.

DISCLOSURE

The authors state that they have no conflicts of interest.

REFERENCES

- Abarca C, Albrecht U, Spanagel R (2002). Cocaine sensitization and reward are under the influence of circadian genes and rhythm. *Proc Natl Acad Sci USA* 99: 9026–9030.
- Abe H, Honma S, Namiyama M, Masubuchi S, Ikeda M, Ebihara S *et al* (2001). Clock gene expressions in the suprachiasmatic nucleus and other areas of the brain during rhythm splitting in CS mice. *Brain Res Mol Brain Res* 87: 92–99.
- Akhtar RA, Reddy AB, Maywood ES, Clayton JD, King VM, Smith AG *et al* (2002). Circadian cycling of the mouse liver transcriptome, as revealed by cDNA microarray, is driven by the suprachiasmatic nucleus. *Curr Biol* 12: 540–550.
- Alarcon JM, Malleret G, Touzani K, Vronskaya S, Ishii S, Kandel ER *et al* (2004). Chromatin acetylation, memory, and LTP are impaired in CBP+/- mice: a model for the cognitive deficit in Rubinstein-Taybi syndrome and its amelioration. *Neuron* 42: 947–959.
- Albensi BC, Mattson MP (2000). Evidence for the involvement of TNF and NF-kappaB in hippocampal synaptic plasticity. *Synapse* 35: 151–159.
- Ammon S, Mayer P, Riechert U, Tischmeyer H, Holtt V (2003). Microarray analysis of genes expressed in the frontal cortex of

- rats chronically treated with morphine and after naloxone precipitated withdrawal. *Brain Res Mol Brain Res* 112: 113–125.
- Andretic R, Chaney S, Hirsh J (1999). Requirement of circadian genes for cocaine sensitization in *Drosophila*. *Science* 285: 1066–1068.
- Ang E, Chen J, Zagouras P, Magna H, Holland J, Schaeffer E *et al* (2001). Induction of nuclear factor-kappaB in nucleus accumbens by chronic cocaine administration. *J Neurochem* 79: 221–224.
- Ashraf SI, Kunes S (2006). A trace of silence: memory and microRNA at the synapse. *Curr Opin Neurobiol* 16: 535–539.
- Ashraf SI, McLoon AL, Sclarsic SM, Kunes S (2006). Synaptic protein synthesis associated with memory is regulated by the RISC pathway in *Drosophila*. *Cell* 124: 191–205.
- Atkins CM, Selcher JC, Petraitis JJ, Trzaskos JM, Sweatt JD (1998). The MAPK cascade is required for mammalian associative learning. *Nat Neurosci* 1: 602–609.
- Barco A, Alarcon JM, Kandel ER (2002). Expression of constitutively active CREB protein facilitates the late phase of long-term potentiation by enhancing synaptic capture. *Cell* 108: 689–703.
- Barco A, Bailey CH, Kandel ER (2006). Common molecular mechanisms in explicit and implicit memory. *J Neurochem* 97: 1520–1533.
- Barco A, Patterson S, Alarcon JM, Gromova P, Mata-Roig M, Morozov A *et al* (2005). Gene expression profiling of facilitated L-LTP in VP16-CREB mice reveals that BDNF is critical for the maintenance of LTP and its synaptic capture. *Neuron* 48: 123–137.
- Barrot M, Olivier JD, Perrotti LI, DiLeone RJ, Berton O, Eisch AJ *et al* (2002). CREB activity in the nucleus accumbens shell controls gating of behavioral responses to emotional stimuli. *Proc Natl Acad Sci USA* 99: 11435–11440.
- Barrot M, Wallace DL, Bolanos CA, Graham DL, Perrotti LI, Neve RL *et al* (2005). Regulation of anxiety and initiation of sexual behavior by CREB in the nucleus accumbens. *Proc Natl Acad Sci USA* 102: 8357–8362.
- Beaujean N (2002). Fundamental features of chromatin structure. *Cloning Stem Cells* 4: 355–361.
- Berger SL (2001). An embarrassment of niches: the many covalent modifications of histones in transcriptional regulation. *Oncogene* 20: 3007–3013.
- Berton O, McClung CA, Dileone RJ, Krishnan V, Renthal W, Russo SJ *et al* (2006). Essential role of BDNF in the mesolimbic dopamine pathway in social defeat stress. *Science* 311: 864–868.
- Bito H, Deisseroth K, Tsien RW (1996). CREB phosphorylation and dephosphorylation: a Ca(2+)- and stimulus duration-dependent switch for hippocampal gene expression. *Cell* 87: 1203–1214.
- Black YD, Maclaren FR, Naydenov AV, Carlezon Jr WA, Baxter MG, Konradi C (2006). Altered attention and prefrontal cortex gene expression in rats after binge-like exposure to cocaine during adolescence. *J Neurosci* 26: 9656–9665.
- Blendy JA (2006). The role of CREB in depression and antidepressant treatment. *Biol Psychiatry* 59: 1144–1150.
- Boivin DB (2000). Influence of sleep-wake and circadian rhythm disturbances in psychiatric disorders. *J Psychiatry Neurosci* 25: 446–458.
- Bunney WE, Bunney BG (2000). Molecular clock genes in man and lower animals: possible implications for circadian abnormalities in depression. *Neuropsychopharmacology* 22: 335–345.
- Carle TL, Ohnishi YN, Ohnishi YH, Alibhai IN, Wilkinson MB, Kumar A *et al* (2007). Absence of conserved C-terminal degron domain contributes to Δ FosB's unique stability. *Eur J Neurosci* 25: 3009–3019.
- Carlezon Jr WA, Duman RS, Nestler EJ (2005). The many faces of CREB. *Trends Neurosci* 28: 436–445.
- Carlezon Jr WA, Thome J, Olson VG, Lane-Ladd SB, Brodtkin ES, Hiroi N *et al* (1998). Regulation of cocaine reward by CREB. *Science* 282: 2272–2275.
- Charriaut-Marlangue C, Otani S, Creuzet C, Ben-Ari Y, Loeb J (1991). Rapid activation of hippocampal casein kinase II during long-term potentiation. *Proc Natl Acad Sci USA* 88: 10232–10236.
- Chaudhury D, Colwell CS (2002). Circadian modulation of learning and memory in fear-conditioned mice. *Behav Brain Res* 133: 95–108.
- Chaudhury D, Wang LM, Colwell CS (2005). Circadian regulation of hippocampal long-term potentiation. *J Biol Rhythms* 20: 225–236.
- Chekanova JA, Belostotsky DA (2006). MicroRNAs and messenger RNA turnover. *Methods Mol Biol* 342: 73–85.
- Chen AC, Shirayama Y, Shin KH, Neve RL, Duman RS (2001a). Expression of the cAMP response element binding protein (CREB) in hippocampus produces an antidepressant effect. *Biol Psychiatry* 49: 753–762.
- Chen L, Fischle W, Verdin E, Greene WC (2001b). Duration of nuclear NF-kappaB action regulated by reversible acetylation. *Science* 293: 1653–1657.
- Chen SL, Loffler KA, Chen D, Stallcup MR, Muscat GE (2002). The coactivator-associated arginine methyltransferase is necessary for muscle differentiation: CARM1 coactivates myocyte enhancer factor-2. *J Biol Chem* 277: 4324–4333.
- Chen WG, Chang Q, Lin Y, Meissner A, West AE, Griffith EC *et al* (2003). Derepression of BDNF transcription involves calcium-dependent phosphorylation of MeCP2. *Science* 302: 885–889.
- Cheung WL, Briggs SD, Allis CD (2000). Acetylation and chromosomal functions. *Curr Opin Cell Biol* 12: 326–333.
- Cho K, Ennaceur A, Cole JC, Suh CK (2000). Chronic jet lag produces cognitive deficits. *J Neurosci* 20: RC66.
- Choi KH, Whisler K, Graham DL, Self DW (2006). Antisense-induced reduction in nucleus accumbens cyclic AMP response element binding protein attenuates cocaine reinforcement. *Neuroscience* 137: 373–383.
- Colby CR, Whisler K, Steffen C, Nestler EJ, Self DW (2003). Striatal cell type-specific overexpression of DeltaFosB enhances incentive for cocaine. *J Neurosci* 23: 2488–2493.
- Cole RL, Konradi C, Douglass J, Hyman SE (1995). Neuronal adaptation to amphetamine and dopamine: molecular mechanisms of prodynorphin gene regulation in rat striatum. *Neuron* 14: 813–823.
- Collins AL, Levenson JM, Vilaythong AP, Richman R, Armstrong DL, Noebels JL *et al* (2004). Mild overexpression of MeCP2 causes a progressive neurological disorder in mice. *Hum Mol Genet* 13: 2679–2689.
- Crosio C, Heitz E, Allis CD, Borrelli E, Sassone-Corsi P (2003). Chromatin remodeling and neuronal response: multiple signaling pathways induce specific histone H3 modifications and early gene expression in hippocampal neurons. *J Cell Sci* 116: 4905–4914.
- Davie JR, Murphy LC (1990). Level of ubiquitinated histone H2B in chromatin is coupled to ongoing transcription. *Biochemistry* 29: 4752–4757.
- Davis HP, Squire LR (1984). Protein synthesis and memory: a review. *Psychol Bull* 96: 518–559.
- Debruyne JP, Noton E, Lambert CM, Maywood ES, Weaver DR, Reppert SM (2006). A clock shock: mouse CLOCK is not required for circadian oscillator function. *Neuron* 50: 465–477.
- Deisseroth K, Tsien RW (2002). Dynamic multiphosphorylation passwords for activity-dependent gene expression. *Neuron* 34: 179–182.
- Devan BD, Goad EH, Petri HL, Antoniadis EA, Hong NS, Ko CH *et al* (2001). Circadian phase-shifted rats show normal acquisition but impaired long-term retention of place information in the water task. *Neurobiol Learn Mem* 75: 51–62.
- Doi M, Hirayama J, Sassone-Corsi P (2006). Circadian regulator CLOCK is a histone acetyltransferase. *Cell* 125: 497–508.

- Dong Y, Green T, Saal D, Marie H, Neve R, Nestler EJ *et al* (2006). CREB modulates excitability of nucleus accumbens neurons. *Nat Neurosci* 9: 475–477.
- Dubnau J, Chiang AS, Grady L, Barditch J, Gossweiler S, McNeil J *et al* (2003). The *staufen/pumilio* pathway is involved in *Drosophila* long-term memory. *Curr Biol* 13: 286–296.
- Duffield GE, Best JD, Meurers BH, Bittner A, Loros JJ, Dunlap JC (2002). Circadian programs of transcriptional activation, signaling, and protein turnover revealed by microarray analysis of mammalian cells. *Curr Biol* 12: 551–557.
- Duman RS (2002). Pathophysiology of depression: the concept of synaptic plasticity. *Eur Psychiatry* 17(Suppl 3): 306–310.
- Easton A, Arbuzova J, Turek FW (2003). The circadian clock mutation increases exploratory activity and escape-seeking behavior. *Genes Brain Behav* 2: 11–19.
- Ehlers MD (2003). Activity level controls postsynaptic composition and signaling via the ubiquitin-proteasome system. *Nat Neurosci* 6: 231–242.
- Fernandez RI, Lyons LC, Levenson J, Khabour O, Eskin A (2003). Circadian modulation of long-term sensitization in Aplysia. *Proc Natl Acad Sci USA* 100: 14415–14420.
- Fonseca R, Nagerl UV, Bonhoeffer T (2006). Neuronal activity determines the protein synthesis dependence of long-term potentiation. *Nat Neurosci* 9: 478–480.
- Georgescu D, Sears RM, Hommel JD, Barrot M, Bolanos CA, Marsh DJ *et al* (2005). The hypothalamic neuropeptide melanin-concentrating hormone acts in the nucleus accumbens to modulate feeding behavior and forced-swim performance. *J Neurosci* 25: 2933–2940.
- Georgescu D, Zachariou V, Barrot M, Mieda M, Willie JT, Eisch AJ *et al* (2003). Involvement of the lateral hypothalamic peptide orexin in morphine dependence and withdrawal. *J Neurosci* 23: 3106–3111.
- Granados-Fuentes D, Tseng A, Herzog ED (2006). A circadian clock in the olfactory bulb controls olfactory responsivity. *J Neurosci* 26: 12219–12225.
- Grandin LD, Alloy LB, Abramson LY (2006). The social zeitgeber theory, circadian rhythms, and mood disorders: review and evaluation. *Clin Psychol Rev* 26: 679–694.
- Green TA, Alibhai IN, Hommel JD, DiLeone RJ, Kumar A, Theobald DE *et al* (2006). Induction of inducible cAMP early repressor expression in nucleus accumbens by stress or amphetamine increases behavioral responses to emotional stimuli. *J Neurosci* 26: 8235–8242.
- Gronli J, Bramham C, Murison R, Kanhema T, Fiske E, Bjorvatn B *et al* (2006). Chronic mild stress inhibits BDNF protein expression and CREB activation in the dentate gyrus but not in the hippocampus proper. *Pharmacol Biochem Behav* 85: 842–849.
- Guan Z, Giustetto M, Lomvardas S, Kim JH, Miniaci MC, Schwartz JH *et al* (2002). Integration of long-term-memory-related synaptic plasticity involves bidirectional regulation of gene expression and chromatin structure. *Cell* 111: 483–493.
- Guillaumond F, Dardente H, Giguere V, Cermakian N (2005). Differential control of Bmal1 circadian transcription by REV-ERB and ROR nuclear receptors. *J Biol Rhythms* 20: 391–403.
- Hansen TO, Rehfeld JF, Nielsen FC (2004). GSK-3 β reduces cAMP-induced cholecystokinin gene expression by inhibiting CREB binding. *Neuroreport* 15: 841–845.
- Harms E, Young MW, Saez L (2003). CK1 and GSK3 in the *Drosophila* and mammalian circadian clock. *Novartis Found Symp* 253: 267–277. discussion 102–269, 277–284.
- Hess J, Angel P, Schorpp-Kistner M (2004). AP-1 subunits: quarrel and harmony among siblings. *J Cell Sci* 117: 5965–5973.
- Hooper C, Markevich V, Plattner F, Killick R, Schofield E, Engel T *et al* (2007). Glycogen synthase kinase-3 inhibition is integral to long-term potentiation. *Eur J Neurosci* 25: 81–86.
- Hotte M, Thuaud S, Dineley KT, Hemmings Jr HC, Nairn AC, Jay TM (2007). Phosphorylation of CREB and DARPP-32 during late LTP at hippocampal to prefrontal cortex synapses *in vivo*. *Synapse* 61: 24–28.
- Huang YS, Jung MY, Sarkissian M, Richter JD (2002). N-methyl-D-aspartate receptor signaling results in Aurora kinase-catalyzed CPEB phosphorylation and alpha CaMKII mRNA polyadenylation at synapses. *EMBO J* 21: 2139–2148.
- Huletsky A, Niedergang C, Frechette A, Aubin R, Gaudreau A, Poirier GG (1985). Sequential ADP-ribosylation pattern of nucleosomal histones. ADP-ribosylation of nucleosomal histones. *Eur J Biochem* 146: 277–285.
- Hyman SE, Cole RL, Konradi C, Kosofsky BE (1995). Dopamine regulation of transcription factor-target interactions in rat striatum. *Chem Senses* 20: 257–260.
- Hyman SE, Malenka RC, Nestler EJ (2006). Neural mechanisms of addiction: the role of reward-related learning and memory. *Annu Rev Neurosci* 29: 565–598.
- Iijima M, Nikaido T, Akiyama M, Moriya T, Shibata S (2002). Methamphetamine-induced, suprachiasmatic nucleus-independent circadian rhythms of activity and mPer gene expression in the striatum of the mouse. *Eur J Neurosci* 16: 921–929.
- Iitaka C, Miyazaki K, Akaike T, Ishida N (2005). A role for glycogen synthase kinase-3 β in the mammalian circadian clock. *J Biol Chem* 280: 29397–29402.
- Jasnow AM, Shi C, Israel JE, Davis M, Huhman KL (2005). Memory of social defeat is facilitated by cAMP response element-binding protein overexpression in the amygdala. *Behav Neurosci* 119: 1125–1130.
- Johannessen M, Delghandi MP, Moens U (2004). What turns CREB on? *Cell Signal* 16: 1211–1227.
- Jope RS, Roh MS (2006). Glycogen synthase kinase-3 (GSK3) in psychiatric diseases and therapeutic interventions. *Curr Drug Targets* 7: 1421–1434.
- Josselyn SA, Kida S, Silva AJ (2004). Inducible repression of CREB function disrupts amygdala-dependent memory. *Neurobiol Learn Mem* 82: 159–163.
- Josselyn SA, Shi C, Carlezon Jr WA, Neve RL, Nestler EJ, Davis M (2001). Long-term memory is facilitated by cAMP response element-binding protein overexpression in the amygdala. *J Neurosci* 21: 2404–2412.
- Kaltschmidt B, Ndiaye D, Korte M, Pothion S, Arbibe L, Prullage M *et al* (2006). NF-kappaB regulates spatial memory formation and synaptic plasticity through protein kinase A/CREB signaling. *Mol Cell Biol* 26: 2936–2946.
- Kaltschmidt B, Widera D, Kaltschmidt C (2005). Signaling via NF-kappaB in the nervous system. *Biochim Biophys Acta* 1745: 287–299.
- Kawahara Y, Zinshteyn B, Sethupathy P, Iizasa H, Hatzigeorgiou AG, Nishikura K (2007). Redirection of silencing targets by adenosine-to-inosine editing of miRNAs. *Science* 315: 1137–1140.
- Kelz MB, Chen J, Carlezon Jr WA, Whisler K, Gilden L, Beckmann AM *et al* (1999). Expression of the transcription factor deltaFosB in the brain controls sensitivity to cocaine. *Nature* 401: 272–276.
- King DP, Takahashi JS (2000). Molecular genetics of circadian rhythms in mammals. *Annu Rev Neurosci* 23: 713–742.
- King DP, Zhao Y, Sangoram AM, Wilsbacher LD, Tanaka M, Antoch MP *et al* (1997). Positional cloning of the mouse circadian clock gene. *Cell* 89: 641–653.
- Klugmann M, Lehtlein CB, Symes CW, Klausner BC, Brooks AI, Young D *et al* (2006). A novel role of circadian transcription factor DBP in hippocampal plasticity. *Mol Cell Neurosci* 31: 303–314.
- Knippschild U, Gocht A, Wolff S, Huber N, Lohler J, Stoter M (2005). The casein kinase 1 family: participation in multiple cellular processes in eukaryotes. *Cell Signal* 17: 675–689.

- Konradi C, Cole RL, Heckers S, Hyman SE (1994). Amphetamine regulates gene expression in rat striatum via transcription factor CREB. *J Neurosci* 14: 5623–5634.
- Korzus E, Rosenfeld MG, Mayford M (2004). CBP histone acetyltransferase activity is a critical component of memory consolidation. *Neuron* 42: 961–972.
- Krug M, Lossner B, Ott T (1984). Anisomycin blocks the late phase of long-term potentiation in the dentate gyrus of freely moving rats. *Brain Res Bull* 13: 39–42.
- Kumar A, Choi KH, Renthal W, Tsankova NM, Theobald DE, Truong HT *et al* (2005). Chromatin remodeling is a key mechanism underlying cocaine-induced plasticity in striatum. *Neuron* 48: 303–314.
- Kurabayashi N, Hirota T, Harada Y, Sakai M, Fukada Y (2006). Phosphorylation of mCRY2 at Ser557 in the hypothalamic suprachiasmatic nucleus of the mouse. *Chronobiol Int* 23: 129–134.
- Lenox RH, Gould TD, Manji HK (2002). Endophenotypes in bipolar disorder. *Am J Med Genet* 114: 391–406.
- Levenson JM, O’Riordan KJ, Brown KD, Trinh MA, Molfese DL, Sweatt JD (2004). Regulation of histone acetylation during memory formation in the hippocampus. *J Biol Chem* 279: 40545–40559.
- Levenson JM, Roth TL, Lubin FD, Miller CA, Huang IC, Desai P *et al* (2006). Evidence that DNA (cytosine-5) methyltransferase regulates synaptic plasticity in the hippocampus. *J Biol Chem* 281: 15763–15773.
- Levine AA, Guan Z, Barco A, Xu S, Kandel ER, Schwartz JH (2005). CREB-binding protein controls response to cocaine by acetylating histones at the fosB promoter in the mouse striatum. *Proc Natl Acad Sci USA* 102: 19186–19191.
- Liu Y, Wang Y, Wan C, Zhou W, Peng T, Wang Z *et al* (2005). The role of mPer1 in morphine dependence in mice. *Neuroscience* 130: 383–388.
- Lonze BE, Ginty DD (2002). Function and regulation of CREB family transcription factors in the nervous system. *Neuron* 35: 605–623.
- Luki WJ (2007). Micro-RNA speciation in fetal, adult and Alzheimer’s disease hippocampus. *Neuroreport* 18: 297–300.
- Lyons LC, Collado MS, Khabour O, Green CL, Eskin A (2006). The circadian clock modulates core steps in long-term memory formation in Aplysia. *J Neurosci* 26: 8662–8671.
- Lyons LC, Rawashdeh O, Katzoff A, Susswein AJ, Eskin A (2005). Circadian modulation of complex learning in diurnal and nocturnal Aplysia. *Proc Natl Acad Sci USA* 102: 12589–12594.
- Mahadevan LC, Willis AC, Barratt MJ (1991). Rapid histone H3 phosphorylation in response to growth factors, phorbol esters, okadaic acid, and protein synthesis inhibitors. *Cell* 65: 775–783.
- Malenka RC, Bear MF (2004). LTP and LTD: an embarrassment of riches. *Neuron* 44: 5–21.
- Manev H, Uz T (2006). Clock genes: influencing and being influenced by psychoactive drugs. *Trends Pharmacol Sci* 27: 186–189.
- Mansour HA, Monk TH, Nimgaonkar VL (2005). Circadian genes and bipolar disorder. *Ann Med* 37: 196–205.
- Marie H, Morishita W, Yu X, Calakos N, Malenka RC (2005). Generation of silent synapses by acute *in vivo* expression of CaMKIV and CREB. *Neuron* 45: 741–752.
- Martinowich K, Hattori D, Wu H, Fouse S, He F, Hu Y *et al* (2003). DNA methylation-related chromatin remodeling in activity-dependent BDNF gene regulation. *Science* 302: 890–893.
- Mattson MP (2005). NF-kappaB in the survival and plasticity of neurons. *Neurochem Res* 30: 883–893.
- Mayford M, Kandel ER (1999). Genetic approaches to memory storage. *Trends Genet* 15: 463–470.
- Mayr B, Montminy M (2001). Transcriptional regulation by the phosphorylation-dependent factor CREB. *Nat Rev Mol Cell Biol* 2: 599–609.
- McClung CA (2007). Circadian genes, rhythms and the biology of mood disorders. *Pharmacol Ther* 114: 222–232.
- McClung CA, Nestler EJ (2003). Regulation of gene expression and cocaine reward by CREB and DeltaFosB. *Nat Neurosci* 6: 1208–1215.
- McClung CA, Sidiropoulou K, Vitaterna M, Takahashi JS, White FJ, Cooper DC *et al* (2005). Regulation of dopaminergic transmission and cocaine reward by the Clock gene. *Proc Natl Acad Sci USA* 102: 9377–9381.
- McClung CA, Ulevy PG, Perrotti LI, Zachariou V, Berton O, Nestler EJ (2004). DeltaFosB: a molecular switch for long-term adaptation in the brain. *Brain Res Mol Brain Res* 132: 146–154.
- McDearmon EL, Patel KN, Ko CH, Walisser JA, Schook AC, Chong JL *et al* (2006). Dissecting the functions of the mammalian clock protein BMAL1 by tissue-specific rescue in mice. *Science* 314: 1304–1308.
- Mieda M, Williams SC, Richardson JA, Tanaka K, Yanagisawa M (2006). The dorsomedial hypothalamic nucleus as a putative food-entrainable circadian pacemaker. *Proc Natl Acad Sci USA* 103: 12150–12155.
- Miller CA, Sweatt JD (2007). Covalent modification of DNA regulates memory formation. *Neuron* 53: 857–869.
- Miller JC, Jimenez P, Mathe AA (2006). Restraint stress influences AP-1 and CREB DNA-binding activity induced by chronic lithium treatment in the rat frontal cortex and hippocampus. *Int J Neuropsychopharmacol* 12: 1–11.
- Mioduszevska B, Jaworski J, Kaczmarek L (2003). Inducible cAMP early repressor (ICER) in the nervous system—a transcriptional regulator of neuronal plasticity and programmed cell death. *J Neurochem* 87: 1313–1320.
- Montarolo PG, Kandel ER, Schacher S (1988). Long-term heterosynaptic inhibition in Aplysia. *Nature* 333: 171–174.
- Morgan HD, Santos F, Green K, Dean W, Reik W (2005). Epigenetic reprogramming in mammals. *Hum Mol Genet* 14(Spec No 1): R47–R58.
- Mouravlev A, Dunning J, Young D, During MJ (2006). Somatic gene transfer of cAMP response element-binding protein attenuates memory impairment in aging rats. *Proc Natl Acad Sci USA* 103: 4705–4710.
- Nathan D, Sterner DE, Berger SL (2003). Histone modifications: now summoning sumoylation. *Proc Natl Acad Sci USA* 100: 13118–13120.
- Naylor E, Bergmann BM, Krauski K, Zee PC, Takahashi JS, Vitaterna MH *et al* (2000). The circadian clock mutation alters sleep homeostasis in the mouse. *J Neurosci* 20: 8138–8143.
- Nelson ED, Kavalali ET, Monteggia LM (2006). MeCP2-dependent transcriptional repression regulates excitatory neurotransmission. *Curr Biol* 16: 710–716.
- Nestler EJ (2005). Is there a common molecular pathway for addiction? *Nat Neurosci* 8: 1445–1449.
- Nestler EJ, Carlezon Jr WA (2006). The mesolimbic dopamine reward circuit in depression. *Biol Psychiatry* 59: 1151–1159.
- Nestler EJ, Barrot M, DiLeone RJ, Eisch AJ, Gold SJ, Monteggia LM (2002). Neurobiology of depression. *Neuron* 34: 13–25.
- Newton SS, Thome J, Wallace TL, Shirayama Y, Schlesinger L, Sakai N *et al* (2002). Inhibition of cAMP response element-binding protein or dynorphin in the nucleus accumbens produces an antidepressant-like effect. *J Neurosci* 22: 10883–10890.
- Nikaido T, Akiyama M, Moriya T, Shibata S (2001). Sensitized increase of period gene expression in the mouse caudate/putamen caused by repeated injection of methamphetamine. *Mol Pharmacol* 59: 894–900.
- Ogden CA, Rich ME, Schork NJ, Paulus MP, Geyer MA, Lohr JB *et al* (2004). Candidate genes, pathways and mechanisms for bipolar (manic-depressive) and related disorders: an expanded convergent functional genomics approach. *Mol Psychiatry* 9: 1007–1029.

- Olausson P, Jentsch JD, Tronson N, Neve RL, Nestler EJ, Taylor JR (2006). DeltaFosB in the nucleus accumbens regulates food-reinforced instrumental behavior and motivation. *J Neurosci* 26: 9196–9204.
- Olson VG, Green TA, Neve RL, Nestler EJ (2007). Regulation of morphine reward and feeding by CREB in the lateral hypothalamus. *Synapse* 61: 110–113.
- Olson VG, Zabetian CP, Bolanos CA, Edwards S, Barrot M, Eisch AJ *et al* (2005). Regulation of drug reward by cAMP response element-binding protein: evidence for two functionally distinct subregions of the ventral tegmental area. *J Neurosci* 25: 5553–5562.
- O'Riordan KJ, Huang IC, Pizzi M, Spano P, Boroni F, Egli R *et al* (2006). Regulation of nuclear factor kappaB in the hippocampus by group I metabotropic glutamate receptors. *J Neurosci* 26: 4870–4879.
- Otani S, Roisin-Lallemant MP, Ben-Ari Y (1992). Enhancement of extracellular protein concentrations during long-term potentiation in the rat hippocampal slice. *Neuroscience* 47: 265–272.
- Peakman MC, Colby C, Perrotti LI, Tekumalla P, Carle T, Ulery P *et al* (2003). Inducible, brain region-specific expression of a dominant negative mutant of c-Jun in transgenic mice decreases sensitivity to cocaine. *Brain Res* 970: 73–86.
- Perrotti LI, Hadeishi Y, Ulery PG, Barrot M, Monteggia L, Duman RS *et al* (2004). Induction of deltaFosB in reward-related brain structures after chronic stress. *J Neurosci* 24: 10594–10602.
- Pliakas AM, Carlson RR, Neve RL, Konradi C, Nestler EJ, Carlezon Jr WA (2001). Altered responsiveness to cocaine and increased immobility in the forced swim test associated with elevated cAMP response element-binding protein expression in nucleus accumbens. *J Neurosci* 21: 7397–7403.
- Presutti C, Rosati J, Vincenti S, Nasi S (2006). Non coding RNA and brain. *BMC Neurosci* 7(Suppl 1): S5.
- Prickaerts J, Moechars D, Cryns K, Lenaerts I, van Craenendonck H, Goris I *et al* (2006). Transgenic mice overexpressing glycogen synthase kinase 3beta: a putative model of hyperactivity and mania. *J Neurosci* 26: 9022–9029.
- Reick M, Garcia JA, Dudley C, McKnight SL (2001). NPAS2: an analog of clock operative in the mammalian forebrain. *Science* 293: 506–509.
- Reppert SM, Weaver DR (2001). Molecular analysis of mammalian circadian rhythms. *Annu Rev Physiol* 63: 647–676.
- Romano A, Freudenthal R, Merlo E, Routtenberg A (2006). Evolutionarily-conserved role of the NF-kappaB transcription factor in neural plasticity and memory. *Eur J Neurosci* 24: 1507–1516.
- Roybal K, Theobald D, Graham A, Dinieri JA, Russo SJ, Krishnan V *et al* (2007). Mania-like behavior induced by disruption of CLOCK. *Proc Natl Acad Sci USA* 104: 6406–6411.
- Rudy JW, Biedenkapp JC, Moineau J, Bolding K (2006). Anisomycin and the reconsolidation hypothesis. *Learn Mem* 13: 1–3.
- Sakai T, Tamura T, Kitamoto T, Kidokoro Y (2004). A clock gene, period, plays a key role in long-term memory formation in *Drosophila*. *Proc Natl Acad Sci USA* 101: 16058–16063.
- Salas TR, Reddy SA, Clifford JL, Davis RJ, Kikuchi A, Lippman SM *et al* (2003). Alleviating the suppression of glycogen synthase kinase-3beta by Akt leads to the phosphorylation of cAMP-response element-binding protein and its transactivation in intact cell nuclei. *J Biol Chem* 278: 41338–41346.
- Sassone-Corsi P, Mizzen CA, Cheung P, Crosio C, Monaco L, Jacquot S *et al* (1999). Requirement of Rsk-2 for epidermal growth factor-activated phosphorylation of histone H3. *Science* 285: 886–891.
- Sato TK, Panda S, Miraglia LJ, Reyes TM, Rudic RD, McNamara P *et al* (2004). A functional genomics strategy reveals Rora as a component of the mammalian circadian clock. *Neuron* 43: 527–537.
- Schafe GE, Atkins CM, Swank MW, Bauer EP, Sweatt JD, LeDoux JE (2000). Activation of ERK/MAP kinase in the amygdala is required for memory consolidation of pavlovian fear conditioning. *J Neurosci* 20: 8177–8187.
- Schratt GM, Tuebing F, Nigh EA, Kane CG, Sabatini ME, Kiebler M *et al* (2006). A brain-specific microRNA regulates dendritic spine development. *Nature* 439: 283–289.
- Schroeder FA, Lin CL, Crusio WE, Akbarian S (2006). Anti-depressant-like effects of the histone deacetylase inhibitor, sodium butyrate, in the mouse. *Biol Psychiatry* 62: 55–64.
- Shaw-Lutchman TZ, Barrot M, Wallace T, Gilden L, Zachariou V, Impey S *et al* (2002). Regional and cellular mapping of cAMP response element-mediated transcription during naltrexone-precipitated morphine withdrawal. *J Neurosci* 22: 3663–3672.
- Shaw-Lutchman TZ, Impey S, Storm D, Nestler EJ (2003). Regulation of CRE-mediated transcription in mouse brain by amphetamine. *Synapse* 48: 10–17.
- Si K, Giustetto M, Etkin A, Hsu R, Janisiewicz AM, Miniaci MC *et al* (2003). A neuronal isoform of CPEB regulates local protein synthesis and stabilizes synapse-specific long-term facilitation in aplysia. *Cell* 115: 893–904.
- Spanagel R, Pendyala G, Abarca C, Zghoul T, Sanchis-Segura C, Magnone MC *et al* (2005). The clock gene Per2 influences the glutamatergic system and modulates alcohol consumption. *Nat Med* 11: 35–42.
- Stanton PK, Sarvey JM (1984). Blockade of long-term potentiation in rat hippocampal CA1 region by inhibitors of protein synthesis. *J Neurosci* 4: 3080–3088.
- Stokkan KA, Yamazaki S, Tei H, Sakaki Y, Menaker M (2001). Entrainment of the circadian clock in the liver by feeding. *Science* 291: 490–493.
- Taubenfeld SM, Wiig KA, Bear MF, Alberini CM (1999). A molecular correlate of memory and amnesia in the hippocampus. *Nat Neurosci* 2: 309–310.
- Tsankova NM, Berton O, Renthall W, Kumar A, Neve RL, Nestler EJ (2006). Sustained hippocampal chromatin regulation in a mouse model of depression and antidepressant action. *Nat Neurosci* 9: 519–525.
- Tsankova NM, Kumar A, Nestler EJ (2004). Histone modifications at gene promoter regions in rat hippocampus after acute and chronic electroconvulsive seizures. *J Neurosci* 24: 5603–5610.
- Ulery PG, Nestler EJ (2007). Regulation of DeltaFosB transcriptional activity by Ser27 phosphorylation. *Eur J Neurosci* 25: 224–230.
- Ulery PG, Rudenko G, Nestler EJ (2006). Regulation of DeltaFosB stability by phosphorylation. *J Neurosci* 26: 5131–5142.
- Uz T, Ahmed R, Akhisaroglu M, Kurtuncu M, Imbesi M, Dirim Arslan A *et al* (2005). Effect of fluoxetine and cocaine on the expression of clock genes in the mouse hippocampus and striatum. *Neuroscience* 134: 1309–1316.
- Valentinuzzi VS, Menna-Barreto L, Xavier GF (2004). Effect of circadian phase on performance of rats in the Morris water maze task. *J Biol Rhythms* 19: 312–324.
- Vo N, Klein ME, Varlamova O, Keller DM, Yamamoto T, Goodman RH *et al* (2005). A cAMP-response element binding protein-induced microRNA regulates neuronal morphogenesis. *Proc Natl Acad Sci USA* 102: 16426–16431.
- Wagatsuma A, Sugai R, Chono K, Azami S, Hatakeyama D, Sadamoto H *et al* (2004). The early snail acquires the learning. Comparison of scores for conditioned taste aversion between morning and afternoon. *Acta Biol Hung* 55: 149–155.
- Walker MP, Stickgold R (2006). Sleep, memory, and plasticity. *Annu Rev Psychol* 57: 139–166.
- Wallace TL, Stellitano KE, Neve RL, Duman RS (2004). Effects of cyclic adenosine monophosphate response element binding protein overexpression in the basolateral amygdala on behavioral models of depression and anxiety. *Biol Psychiatry* 56: 151–160.

- Walters CL, Blendy JA (2001). Different requirements for cAMP response element binding protein in positive and negative reinforcing properties of drugs of abuse. *J Neurosci* 21: 9438–9444.
- Walters CL, Cleck JN, Kuo YC, Blendy JA (2005). Mu-opioid receptor and CREB activation are required for nicotine reward. *Neuron* 46: 933–943.
- Walters CL, Kuo YC, Blendy JA (2003). Differential distribution of CREB in the mesolimbic dopamine reward pathway. *J Neurochem* 87: 1237–1244.
- Wang LM, Suthana NA, Chaudhury D, Weaver DR, Colwell CS (2005). Melatonin inhibits hippocampal long-term potentiation. *Eur J Neurosci* 22: 2231–2237.
- Weaver IC, Cervoni N, Champagne FA, D'Alessio AC, Sharma S, Seckl JR *et al* (2004). Epigenetic programming by maternal behavior. *Nat Neurosci* 7: 847–854.
- Weaver IC, Champagne FA, Brown SE, Dymov S, Sharma S, Meaney MJ *et al* (2005). Reversal of maternal programming of stress responses in adult offspring through methyl supplementation: altering epigenetic marking later in life. *J Neurosci* 25: 11045–11054.
- Werme M, Messer C, Olson L, Gilden L, Thoren P, Nestler EJ *et al* (2002). Delta FosB regulates wheel running. *J Neurosci* 22: 8133–8138.
- Williams JM, Beckmann AM, Mason-Parker SE, Abraham WC, Wilce PA, Tate WP (2000). Sequential increase in Egr-1 and AP-1 DNA binding activity in the dentate gyrus following the induction of long-term potentiation. *Brain Res Mol Brain Res* 77: 258–266.
- Williamson SL, Christodoulou J (2006). Rett syndrome: new clinical and molecular insights. *Eur J Hum Genet* 14: 896–903.
- Wirz-Justice A (2006). Biological rhythm disturbances in mood disorders. *Int Clin Psychopharmacol* 21(Suppl 1): S11–S15.
- Wood MA, Kaplan MP, Park A, Blanchard EJ, Oliveira AM, Lombardi TL *et al* (2005). Transgenic mice expressing a truncated form of CREB-binding protein (CBP) exhibit deficits in hippocampal synaptic plasticity and memory storage. *Learn Mem* 12: 111–119.
- Wu GY, Deisseroth K, Tsien RW (2001). Activity-dependent CREB phosphorylation: convergence of a fast, sensitive calmodulin kinase pathway and a slow, less sensitive mitogen-activated protein kinase pathway. *Proc Natl Acad Sci USA* 98: 2808–2813.
- Wu J, Xie X (2006). Comparative sequence analysis reveals an intricate network among REST, CREB and miRNA in mediating neuronal gene expression. *Genome Biol* 7: R85.
- Wu L, Wells D, Tay J, Mendis D, Abbott MA, Barnitt A *et al* (1998). CPEB-mediated cytoplasmic polyadenylation and the regulation of experience-dependent translation of alpha-CaMKII mRNA at synapses. *Neuron* 21: 1129–1139.
- Xu Y, Ku B, Tie L, Yao H, Jiang W, Ma X *et al* (2006). Curcumin reverses the effects of chronic stress on behavior, the HPA axis, BDNF expression and phosphorylation of CREB. *Brain Res* 1122: 56–64.
- Yeh SH, Lin CH, Gean PW (2004). Acetylation of nuclear factor-kappaB in rat amygdala improves long-term but not short-term retention of fear memory. *Mol Pharmacol* 65: 1286–1292.
- Yeh SH, Lin CH, Lee CF, Gean PW (2002). A requirement of nuclear factor-kappaB activation in fear-potentiated startle. *J Biol Chem* 277: 46720–46729.
- Yin JC, Tully T (1996). CREB and the formation of long-term memory. *Curr Opin Neurobiol* 6: 264–268.
- Yin JC, Del Vecchio M, Zhou H, Tully T (1995). CREB as a memory modulator: induced expression of a dCREB2 activator isoform enhances long-term memory in *Drosophila*. *Cell* 81: 107–115.
- Yuferov V, Butelman ER, Kreek MJ (2005). Biological clock: biological clocks may modulate drug addiction. *Eur J Hum Genet* 13: 1101–1103.
- Yuferov V, Krosiak T, Laforge KS, Zhou Y, Ho A, Kreek MJ (2003). Differential gene expression in the rat caudate putamen after 'binge' cocaine administration: advantage of triplicate microarray analysis. *Synapse* 48: 157–169.
- Zachariou V, Bolanos CA, Selley DE, Theobald D, Cassidy MP, Kelz MB *et al* (2006). An essential role for DeltaFosB in the nucleus accumbens in morphine action. *Nat Neurosci* 9: 205–211.
- Zghoul T, Abarca C, Sanchis-Segura C, Albrecht U, Schumann G, Spanagel R (2007). Ethanol self-administration and reinstatement of ethanol-seeking behavior in Per1(Brdm1) mutant mice. *Psychopharmacology (Berlin)* 190: 13–19.
- Zhang H, Darwanto A, Linkhart TA, Sowers LC, Zhang L (2007). Maternal cocaine administration causes an epigenetic modification of PKC{epsilon} gene expression in fetal rat heart. *Mol Pharmacol* 71: 1319–1328.
- Zhu H, Lee M, Agatsuma S, Hiroi N (2007). Pleiotropic impact of constitutive fosB inactivation on nicotine-induced behavioral alterations and stress-related traits in mice. *Hum Mol Genet* 16: 820–836.
- Zueger M, Urani A, Chourbaji S, Zacher C, Lipp HP, Albrecht U *et al* (2006). mPer1 and mPer2 mutant mice show regular spatial and contextual learning in standardized tests for hippocampus-dependent learning. *J Neural Transm* 113: 347–356.