lower than the lowest excitation potential of neon (16.6 volts). Therefore the hypothesis seems to be justified that the acceleration of the electrons occurs for the greater part by the steady field, due to the differences in space potential.

A recently observed experimental fact relating to the distribution of the space potential may throw light on the observations. When a high-frequency glow in a neon discharge tube of the type already described above is operated in the region of the moderate frequencies, say, 3×10^6 cycles, sometimes there are present near each of the two electrodes a series of two or three luminous layers, divided one from the other by darker spaces; the space between the electrode and the first layer is perfectly dark.

The appearance of these layers is very similar to that of the luminous layers in the case of a direct current discharge in neon at small currents, as first described by Holst and Oosterhuis¹.

Fig. 1 (a) is a photograph of the layers in the case of a direct current of 8 micro-amperes. The electrodes are coated with a caesium film on caesium oxide, the pressure of the neon is about 45 mm. As there were from cathode to anode 13 luminous layers and the total voltage was 256, the potential difference between two layers is about 20 volts. Fig. 1 (b) shows a high-frequency discharge in the same tube. In front of the two electrodes three luminous layers are visible, which fact leads to the conclusion that there is a potential drop of about 80 volts near the electrodes.

A. C. VAN DORSTEN.

Laboratorium voor Technische Physica,
Delft.
Sept. 26.

¹ G. Holst and E. Oosterhuis, *Physica*, **1**, 78; 1921. *Phil. Mag.*, **46**, 1117; 1923. See also M. J. Druyvesteyn, *Z. Phys.*, **3**, 33; 1932.

Ionisation of the Ionosphere

IN a recent letter¹ to NATURE, L. Tonks has suggested that the relation between ionisation density N and critical frequency f should be

$$N = (\pi m/e^2) f^2,$$

rather than

$$N = (3/2) (\pi m/e^2) f^2,$$

which has been used in some recent papers. This appears to be correct; for the radio application, the matter may be presented in the following way.

Following Lorentz, we may express the field inside an ionised medium as

$$\vec{E} = \vec{E}_1 + \frac{1}{3}\vec{P} + \vec{E}_2,$$

where $\vec{E}_1 + \frac{1}{3}\vec{P}$ is the portion of the field due to all the charges outside a small sphere Δr of radius large compared to the average spacing of the charges and $\vec{E}_2$ is the field due to the charges inside the Δr sphere. The difficulty in evaluating $\vec{E}$ is evidently that of evaluating $\vec{E}_2$, since $\vec{E}_1$ may be evaluated by the usual potential theory. However, for the radio case it is sufficient to know the value of the space average field $\vec{E}^0$ which may be defined as

$$\vec{E}^0 = \frac{1}{\Delta r} \int_{\Delta r} \vec{E} dr.$$

For any distribution of charges, it may be shown that $\vec{E}_2^0 = -\frac{1}{3}\vec{P}$ so that $\vec{E}^0 = \vec{E}_1^0$.

We may assume for the sake of illustration that we are concerned with the E layer and a frequency of 3,000 kc./s. The mean thermal velocity of the electrons $\approx 10^7$ cm./sec., so that the electron on which

the force is to be calculated moves in a random direction about 3 cm. during a single period of the radio wave. Since the average electron spacing in the upper atmosphere is about 10^{-2} cm., it is thus evident that the space average field is the one which applies to the present problem. Using the space average field we arrive at the value

$$N = (\pi m/e^2) f^2.$$

As an alternative way of viewing the problem, consider the time average field for a period short compared to that of the radio wave. Such a time average field will be sufficient for this problem. During $1/30$ of the period of the radio wave, the electron moves in a random direction about 10^{-1} cm. and since the electron spacing is about 10^{-2} cm., the ionised medium over this short period of time becomes essentially continuous, so that we can derive the expression for the field rigorously from potential theory, obtaining as before $\vec{E} = \vec{E}_1$.

K. A. NORTON.

Bureau of Standards,
Washington, D.C.
Sept. 21.

¹ NATURE, **131**, 101, July 15, 1933.

Linear Transformations of Hierarchical Systems

AS is well known, each of a number of variables which constitute a hierarchical system can be expressed as a linear function of a single general factor g and a specific factor. Each of these specific factors is independent of each of the other specific factors and of g . According to Prof. Spearman's theory of the single general factor, the same g ought to be the general factor of the scores in each of two or more series of mental tests, when the scores in each case tend to constitute a hierarchical system.

But E. B. Wilson has claimed¹ that if a new set of variables is created by linear transformation—"a system of weighting which leads to new scores"—"we can, while keeping to hierarchical systems, do pretty much what we want with g "; or, as he put it in a later paper, g was "relative to the set-up". If that were the whole story, Prof. Spearman's theory would be in serious trouble.

I have recently investigated this matter, with the following results:

(1) There is but an infinitely small chance that a linear transformation (made at random) of a hierarchical system would itself be hierarchical: no new general factor will turn up by accident.

(2) It is nevertheless possible to design linear transformations of a hierarchical system so as to produce new systems which are themselves hierarchical.

(3) It is even possible to make such new systems conserve g or, in other words, to make the new hierarchical system have a general factor which has the same correlations as the original g with the new variables.

(4) It is not possible to conserve g except (a) by using some negative weights in the transformation, or (b) by using so many zero weights that, in the linear expressions for the new scores in terms of the original, the product of any two coefficients of the same original variable is zero.

As regards (a), I understand that negative weights would never be employed by psychologists in handling scores made in mental tests. On the other hand, (b) is satisfied when tests are pooled or, in other