petrol; a good dense deposit is made round the india-ink junctions, and over the rest of the quartz a fairly even deposit not too dense is formed. After the rod is allowed to cool, the resistance is measured and can then be very easily and quickly adjusted in value by evenly wiping off some of the deposit with clean dry cotton wool or by adding a little more soot. The resistance may then be sealed into a clean quartz tube or left bare or coated with paraffin wax, of good white quality, by simple dipping.

A bare 10^{14} ohm resistance of this type has been used satisfactorily for five months, and two wax-covered 10^{10} ohm resistances have functioned well for two months as coupling resistances in counter apparatus. The dipping in molten paraffin wax alters the resistance, usually increasing it considerably, but there is little difficulty in allowing for this and in making a resistance of the approximate desired value. The method of construction obviously gives the possibility of a considerable range of resistance and carrying capacity and it is intended to investigate the properties of the resistance at a later date. The good results so far obtained combined with the ease of adjustment and the cheapness of the construction would seem to warrant the present note.

P. W. BURBIDGE.

Cavendish Laboratory,
Cambridge.

The Origin of Tektites

IN his contributions to NATURE on the "Origin of Tektites" Dr. L. J. Spencer¹ has put forward the view "that tektites are not meteoric, though they are connected with the fall of large meteoric masses, but that they have resulted from the fusion of terrestrial rocks, especially in sandy deserts, by the heat so developed".

The radioactivity of eight specimens of tektites and of some pieces of glass of different kinds was determined with the view of deciding whether they were meteoric or terrestrial in origin. I am obliged to Dr. H. Michel of the Natural History Museum of Vienna, and to the Trustees of the British Museum, for these specimens. The problem was suggested by Prof. H. Mache, of the Technical High School of Vienna, and the work was carried out in his laboratory. The results of determinations are given in the accompanying table.

		Ra $\times 10^{-12}$ per gm.	Th $\times 10^{-4}$ per gm.
1.	Moldavite	1.07	1.08
2.	Moldavite (Habř, Bohemia) ..	1.02	1.60
3.	Moldavite (Probsch) ..	0.78	1.60
4.	Moldavite (Radomolice, Bohemia)	0.99	1.86
5.	Billitonite ..	0.96	0.96
6.	Australite (Lake Eyre district) ..	0.96	0.50
7.	Australite (Victoria) ..	0.85	1.84
8.	Darwin glass (Tasmania) ..	0.50	1.13
9.	Glass (Old beads) ..	0.48	

These results show that the radium content of all the tektites except Darwin glass falls between 0.9×10^{-12} and 1.00×10^{-12} . A difference of 0.1×10^{-12} is found even in different parts of the same rock and has no significance.

This constancy in the radioactive contents of tektites coming from such distant parts of the world as Europe, Asia and Australia clearly suggests some kind of genetic relationship. It is difficult to imagine

that glass formed in three different continents from different raw materials should have the same radium content.

The radioactive contents of several kinds of glass, including glass beads of prehistoric age, was determined, but the values found were low and very different, the highest being 0.48×10^{-12} per gm. There is no possibility of these tektites deriving their radioactive contents from iron meteorites in any way, as these meteorites are very poor in radium, the amount being of the order of 10^{-14} .

The radioactivity of Darwin glass differs from all other tektites considerably and agrees more with glass than with tektites. In this connexion the similarity of silica glass from Wabar and Henbury to the Darwin glass as observed by Dr. Spencer is noteworthy.

The value 1.00×10^{-12} per gm. found for the tektites seems fairly in accord with several such determinations made for granites, which represent the salic part of the earth's crust², and show a similar chemical composition. This strongly suggests that these tektites are derived from some mass which agrees in chemical composition as well as in radioactivity with the granitic layer of the earth.

It will be interesting to determine the radioactive content of the tektites recently discovered in French Indo-China and described by Prof. Lacroix, and of those discovered in the Philippine Islands¹. The determination of the radioactivity of the glasses discovered at Rub' al Khali in Arabia, and at Henbury in Central Australia, will throw further light on the origin of tektites.

V. S. DUBEY.

Department of Geology,
Benares Hindu University.

¹ L. J. Spencer, "Origin of Tektites", NATURE, 131, 117, Jan. 28; 876, June 17, 1933.

² C. S. Piggot, Amer. J. Sci., 5, 21, 35, January 1931.

WITH reference to Dr. L. J. Spencer's letter in NATURE of October 7, p. 571, on the origin of tektites, the strongest argument against his suggestion that they were formed by the fusion of terrestrial material in meteorite craters is that no tektite has yet been described, so far as I am aware, containing partially fused rock or sand. A further possible objection that no tektite hitherto described contains so many vesicles as glass from meteorite craters is discounted by Prof. A. Lacroix's description of 95 kgm. of tektites from the island of Hainan and the neighbouring mainland, among which glass that originally enclosed vesicles of more than a decimetre in diameter is of frequent occurrence ("Les Tectites de l'Indo-chine", Paris, 1932, appendix following p. 235). On the other hand, Dr. Spencer's suggestion finds support in the presence of metallic spheres, resembling those of the Wabar glass, in Darwin glass and in tektites from Indo-China, as described in his letter; and also in the presence of nickel in three Malayan tektites, proved by analyses by Mr. J. C. Shenton. As Dr. Spencer remarks, the evidence for tektites being meteoric is entirely negative; and although his suggestion cannot yet be accepted unreservedly, it appeals to me as the best yet proposed for the origin of these bodies.

J. B. SCRIVENOR.

68 Chaucer Road,
Bedford.
Oct. 9.