

Letters to the Editor

[The Editor does not hold himself responsible for opinions expressed by his correspondents. Neither can he undertake to return, nor to correspond with the writers of, rejected manuscripts intended for this or any other part of NATURE. No notice is taken of anonymous communications.]

Latitude Effect of Cosmic Radiation

It was found for the first time by Clay¹ on voyages between Holland and Java that the intensity of cosmic radiation has a minimum in the neighbourhood of the magnetic equator. The extensive survey directed by Compton² confirmed the existence of this 'latitude effect' and showed it to be more pronounced at higher altitude. More accurate results at sea-level are due to an investigation of Clay and Berlage³. As this again refers to the line from Holland to Java, I thought it would be worth while to perform analogous measurements on a trip from Holland to South Africa. During this investigation Hoerlin⁴ published results he obtained on the line Peru—Strait of Magellan—Hamburg. These results and those of the other authors as given by Clay are represented in Fig. 1 by continuous curves, my own results by open circles. Clearly the latter lie somewhat closer to Clay's curve than to Hoerlin's.

ions/cm² sec.




Fig. 1. Latitude effect of cosmic radiation. Circles indicate results of author. Vertical scale indicates number of pairs of ions in normal air.

Unfortunately, my apparatus broke down in the tropics, so I have not been able to get evidence on the remarkable difference between the southern and northern hemispheres as indicated by Hoerlin's results. Though we may feel satisfied that an equatorial minimum of the same order of magnitude is found by all investigators (indicating that the cosmic radiation consists largely of a cosmic rain of charged particles) it would seem that an accurate repetition of this kind of measurement to obtain the exact shape of the curve is not superfluous.

Regarding my observations, the following particulars may be given. The ionisation chamber had a volume of 3 litres and contained argon at a pressure of 30 atm. It was shielded by 8 cm. of iron and was placed in a hut on board the S.S. *Springfontein* of the Holland Africa line, the deck over it being of negligible thickness. The wall of the ionisation chamber was brought to 120 v. and the ionisation current collected on an insulated rod connected to a Lindemann electrometer and to a small capacity

(4 cm.). To start an observation, the earthing key of the rod was opened and a stop-watch set running at the same moment. The electrometer was kept at zero by gradually applying a potential to the capacity so as to compensate the charge due to the ionisation current. After some time (about 6 min.) the potential (about 3 v.) was read on a voltmeter. From this the number of ions produced in the chamber per cm.³ per sec. may be deduced, assuming saturation. This number is called the 'intensity' of cosmic radiation. A small correction for barometric pressure was applied to it (2.4 per cent for 1 cm. mercury). In the graph in Fig. 1 these values (like those of Hoerlin) have been multiplied by such a factor (1/33.2) as to make the value at 50° coincide with the value given by Clay for normal air.

I wish to thank the Groninger Universiteitsfonds for a grant of money, Prof. Coster for allowing the apparatus to be made in the workshop of his laboratory, Prof. Clay for some kind advice and finally the directors of the Holland Africa line and the crew of the S.S. *Springfontein* for their kind collaboration.

J. A. PRINS.

Natuurkundig Laboratorium
der Rijks-Universiteit,
Groningen.
Oct. 19.

¹ J. Clay, *Proc. Amsterdam*, **30**, 1115, 1927; **31**, 1091, 1928.
² A. H. Compton, *Phys. Rev.*, **43**, 387; 1933.
³ J. Clay and H. P. Berlage, *Naturwiss.*, **20**, 687; 1932. J. Clay, *Naturwiss.*, **21**, 43; 1933.
⁴ H. Hoerlin, *NATURE*, **132**, 61, July 8, 1933.

Nuclear Moments of Xenon

FROM an investigation of the hyperfine structures of the lines of neutral xenon (XeI) which fall in the region $\lambda\lambda 4200-8500$, I have been able to show that the majority of the *s-p* transitions possess fairly complicated structures. These lines, therefore, are unsuitable as wave-length standards.

In a recent note, Kopfermann¹ reports that he has analysed a number of XeI line structures and finds that the isotope Xe¹²⁹ has the nuclear moment $I = \frac{1}{2}$, while that of Xe¹³¹ is probably $\frac{3}{2}$, the *g(I)*-factors being of opposite sign. My analysis confirms this result for Xe¹²⁹, but nothing precise can be stated for Xe¹³¹, except that $I > \frac{1}{2}$. The analogy between the structures of certain xenon lines and those of corresponding mercury lines is, however, so close as to favour the assignment of $I = \frac{3}{2}$ to Xe¹³¹.

Hyperfine separations of the terms of Xe¹²⁹ ($I = \frac{1}{2}$).

Term	Hyperfine separation	Term	Hyperfine separation
1s ₂	- 0.290 cm. ⁻¹	2p ₇	- 0.065 cm. ⁻¹
1s ₄	- 0.057	3p ₆	- 0.075
1s ₄	- 0.198	3p ₈	- 0.134
2p ₂	+ 0.098	3p ₈	- 0.078
2p ₆	- 0.250	3p ₁₀	- 0.076
2p ₈	- 0.070	4Y	+ 0.069

The "-" sign indicates that the hyperfine levels are inverted.

The results of my measurements are summarised in the accompanying table of term structures, in which the hyperfine separations of Xe¹²⁹ are given. Since Kopfermann gave no experimental results, nor any indication of the scope of his observations, a comparison of our data is unfortunately not possible.

E. GWYNNE JONES.

Imperial College of Science,
London, S.W.7.
Oct. 24.

¹ H. Kopfermann, *Naturwiss.*, **39**, 704; 1933.