

ground, but without any of the unpleasant feelings associated with the word giddiness.

In people who are particularly susceptible, vertigo accompanied even by nausea and vomiting may easily be brought about by kinæsthetic impulses or by moving visual stimuli. But that the labyrinth is by far the most usual and important sense organ from which vertigo is aroused, and probably the only one concerned in sea-sickness, is now firmly established. James, Kreidl and Myginol have all commented on the immunity from sea-sickness of deaf-mutes in whom the labyrinths were deficient. The experiments of Dreyfuss on guinea-pigs and of

Kreidl on dogs, cats and pigeons pointed in the same direction. Decerebration, decerebellation, or section of both vagi leaves sea-sickness unaffected, but removal of both labyrinths or section of both eighth nerves results in complete immunity. The most likely explanation of the vomiting seems to be that it is caused by spread of excitation from the vestibular nuclei to the neighbouring medullary 'vomiting centre'.

As a result of the interest aroused in the discussion, a joint research committee of the Sections of Physiology and Psychology has been set up to investigate the conditions of vertigo and its relation to dis-orientation.

Fishes of Mountain Streams

DR. SUNDER LAL HORA, of the Zoological Survey of India, has for many years devoted special attention to the study of the fauna of rapidly running waters in the hill streams of India. His knowledge of this difficult and interesting branch of zoology is unrivalled. No one, therefore, could be better qualified than he to undertake a detailed investigation* of that remarkable group of cyprinoid fishes, the Homalopteridæ. These fishes, inhabitants of swiftly running mountain streams of southern Asia, have undergone a great variety of adaptive modifications induced by the peculiar environmental conditions typical of their habitat.

In the first part of his report the author deals with the taxonomy of the group. It is divided into 2 sub-families—the Homalopterinae, comprising 6 genera and 31 species, and the Gastromyzoninae which is represented by 11 genera having in all about 16 species. In the Homalopterinae 4 new species belonging to the genera *Homaloptera* and *Lepturichthys*, and 2 new varieties of *Balitora brucei* are described. Among the Gastromyzoninae no new species have been found; but in order properly to classify the existing species 5 new genera are proposed in this paper. No attempt has been made to describe in full each and every species, but wherever an amplification of the already existing description seemed to the author to be desirable the species is

* *Memoirs of the Indian Museum*, vol. 12, No. 2, pp. 263-330. 'Classification, Bionomics and Evolution of Homalopterid Fishes'. By Sunder Lal Hora. Calcutta; December, 1932.

either redescribed or a note is inserted concerning some of its most important features.

In the second part of the paper the bionomics and evolution of the Homalopteridæ are discussed at some length. The most characteristic features of its members such as flattened shape, insertion of the pectoral fins (which are used for adhesion) far forward below or even in front of the eyes, possession of a peculiar rostral groove in front of and continued along each side of the mouth, the peculiar structure of the hard and strong lower jaw, are shown to be definitely correlated with the three most important factors in the environment—strong current, high oxygen content, and nature of the food supply available. Throughout the paper attention is repeatedly directed to the 'communal convergence' that is exhibited by these fishes and to the series of characters showing parallel development in the members of the two sub-families. From all the evidence which he has acquired the author believes that the Homalopteridæ are probably a polyphyletic family the members of which are derived from the Cyprinidæ and Cobitidæ and have come to resemble one another superficially under the influence of the same environmental conditions.

Dr. Hora is to be congratulated on having produced a paper which is not only a valuable addition to the literature on the taxonomy of the Homalopteridæ but is also of great theoretical interest.

G. A. S.

Geological Reconnaissance by Aeroplane in Australia*

IN 1932 the Royal Australian Air Force made flights over many of those areas in Australia which are deemed to be worthy of investigation from the point of view of the discovery of oil. Dr. W. G. Woolnough, who was present as observer and geologist, has now detailed the results obtained in a report which gives valuable information regarding the function and importance of aerial work in assisting and expediting geological survey. The object was to determine the disposition of strata and especially to locate dome structures, the investigation being made partly by visual observation and partly by the study of the photographs taken from the air. Much experience is required before the utmost can be achieved by these methods, and Dr. Woolnough states that he scarcely began to appreciate the significance of details seen from above until he had completed one hundred hours of flying.

* Commonwealth of Australia. Report on Aerial Survey Operations in Australia during 1932. By Dr. W. G. Woolnough. (Canberra: Government Printers.)

From July until September, a circuit of Australia was made—also a visit to Tasmania, atmospheric conditions on the whole being favourable. Over Melville Island observation was hindered by a tribal fight, as part of the strategy consisted in the lighting of extensive bush fires, the smoke of which provided nuclei for the condensation of moisture and the development of clouds. The orientation of the photographs and the elevation of hills and scarps above the surrounding country can be determined by noting the direction and lengths of shadows—provided the time of exposure is accurately known.

In those regions where the rocks are well exposed and where topographical forms are the direct result of the differential erosion of beds, the main tectonic features are easily seen, and examination of the photographs reveals where the detailed ground investigation which is essential to the full elucidation of the structure should be undertaken. Most valuable is the clearness with which the true disposition and