dependent upon any changes in the physical configuration of this circuit due to changes in atmospheric conditions. For example, changes in temperature of the inductance coils and condensers in the circuit will produce changes in the electrical values of these components to an extent depending upon the coefficients of expansion of the materials used in their construction. The limited experimental evidence so far available indicates that changes of frequency resulting from temperature variations may exceed fifty parts in a million per degree centigrade. To a lesser, but by no means negligible, extent, changes in atmospheric pressure will produce a change in capacitance of an air condenser by virtue of the alteration in the dielectric constant of air. A further factor to be taken into account in a complete study of the subject is the effect of the load circuit, which is coupled to the oscillator and by means of which the oscillations generated are put to practical use. A brief consideration of the relevant conditions indicates that in order to minimise the effect of

the load circuit on the oscillation frequency, this circuit should be slightly detuned by an amount which depends upon its effective resistance.

In conclusion, the possibility of securing frequency stability in radio transmitting stations by the use of an automatic monitoring arrangement is discussed briefly in the report recently published. The scheme provides for the frequency of the transmitter to be adjusted directly to agree with that of a small power valve oscillator designed for a high-degree of frequency stability under no-load conditions. If such a scheme can be developed successfully without undue complication of equipment, it may provide one solution to the problem of stabilising the frequency of a simple transmitter which has to be operated over a wide range of frequencies. The whole subject is, however, being investigated in a comprehensive manner, since more than one solution may ultimately be necessary to meet the conditions of practical radio communication.

Physiology of the Blue Whale

By Prof. AUGUST KROGH, Laboratory of Zoophysiology, University of Copenhagen

RATE OF GROWTH

IN the paper by Mackintosh and Wheeler, "Southern Blue and Fin Whales" ("Discovery Reports", I, 1929), a graph is given showing the growth in length of the Blue whale from the foetal state to maturity. In Laurie's paper, "Some Aspects of Respiration in Blue and Fin Whales" ("Discovery Reports", VII, 1933), graphs are given showing the relation between length and weight of Blue whales, and since this relation is remarkably constant, it becomes possible to calculate the weight of the young whales at different stages.

Such a calculation shows the new-born Blue whale of 7 m. length to weigh about 2,000 kgm., while at weaning seven months later, the length has increased to 16 m. and the weight to 23,000 kgm. When the whales become sexually mature at two years of age, the females are on an average 23.7 m. in length and weigh 79,000 kgm. The period from weaning to maturity includes two summer seasons (about twelve months) in antarctic waters with abundant food, and one winter in more northern waters where food is scarce. The increase from 23,000 kgm. to 79,000 kgm. therefore takes place mainly or exclusively during the twelve months when they are in the antarctic. Laurie gives a table showing the composition of a 20 m. Blue whale. Assuming the same composition during the period of growth, the 56,000 kgm. increase should be distributed as follows:

	per cent	kgm.	per cent	kgm.	per cent	kgm.
Meat and entrails	59	33,000	6.5	2,150	23	7,600
Blubber	18.6	10,400	88	9,200	4	400
Bones	19	10,600	45	4,750		
		54,000		16,100		8,000
Calories: 150 mill. + 44 mill. = 194 mill.						
Calories per day in 365 days = 540,000						
" " " " " 518 " = 380,000						

It may be of some interest to compare this rate of growth with that of a pig. The new-born pig weighs about 1 kgm. After seven weeks' lactation, the weight is about 6 kgm. and, feeding with about 200 kgm. of skim milk and 300 kgm. of grain, it attains the weight of 90 kgm., at which it is converted to bacon at the age of six months. The food consumed corresponds to a little more than 1 million calories, and the increase in weight from weaning to 320,000 calories, or 2,600 cal. per day. Assuming an average weight for the growing pig of $\frac{6 + 90}{2}$ = about 50 kgm. and for the growing whale of $\frac{23,000 + 79,000}{2}$ = about 50,000

kgm., it is seen that the growth of the pig per unit weight is about five times as rapid as that of the whale. The comparison should not, however, be made on the weight but on the surface basis, corresponding to the cube root of the square of the weight ($W^{2/3}$). When the whale weighs just 1,000 times the pig, the increase per day should be $1,000^{2/3}$ = 100 times as large, but it is found to be 540,000 calories instead of 260,000, or just double that of the pig, in spite of the fact that the pig has its food served regularly, while the young whale is left to its own resources. It is a pity that the chemical composition of the crustacean *Euphausia superba* constituting the whales' staple diet is not known. It would be extremely interesting to know if whale fat is mainly derived directly from the food or mainly built up by synthesis.

METABOLISM, CIRCULATION AND RESPIRATION

On the assumption that the surface law is valid for whales and that the normal metabolism is

1,000 calories per sq. m. per day, Laurie arrives at the figure 275,000 calories as the resting metabolism of a 122,000 kgm. Blue whale. It is worthy of note that the assumption, for which good reasons are given by Kleiber ("Die Tierernährung", 5, 1933), that metabolism is proportional to $W^{3/4}$ instead of $W^{2/3}$ with a unit value of 72 cal. per day, would raise the calculated metabolism to 460,000 cal. It might, I think, be possible to arrive at a value for the metabolism by calculating the heat loss from an internal temperature of 35° C., a water temperature of, say, 5° C., measurements of the thickness of blubber and its properties as a conductor of heat. A series of temperature determinations through the thickness of blubber of a freshly killed whale would enhance the value of such a determination.

The metabolism of a whale is increased by its muscular movements in swimming. Prof. Carl Hansen, of the Danish Technical High School, has kindly calculated for me the towing resistance at various speeds of a 122,000 kgm. whale, the

FIG. 1.

surface of which he calculates as 230 m². He finds the resistance and horse power necessary at different speeds to be as shown in the accompanying graph (Fig. 1) where *R* represents the resistance in kgm. and *N* the necessary horse power. At 3 knots the necessary power is 1.5 h.p., at 5 knots it is 6.5 h.p. and at 10 knots 46.8 h.p. He assumes 100 per cent efficiency as a propeller for the tail, which he believes to be superior to any screw. If I assume a 22 per cent efficiency for the muscular engine working the propeller, I find as a good approximation 10 litres of oxygen or 50 calories consumed per h.p. per minute. It would be of great interest to have actual determinations of the towing resistance of whales at varying rates of speed, and it should not be very difficult to obtain such figures.

The assumed resting metabolism of 275,000 calories per day corresponds to 38 litres of oxygen per minute. Assuming that the whale swims at an average rate of only 3 knots, we have a total metabolism of 53 litres of oxygen per minute. We may suppose in the case of this whale, in accordance with the findings on man and other mammals

that, at rest, about 40 per cent of the oxygen capacity (14 volumes per cent) of the blood is utilised, which gives us a circulation rate of very nearly 1,000 litres per minute. A Blue whale can swim at the rate of 10 knots when harpooned. This means a metabolism of a little more than 500 litres of oxygen per minute. With such heavy work the oxygen utilisation may reach 80 per cent and the circulation rate would become 4,500 litres per minute.

The capacity of the lungs is estimated by Laurie from their weight at the very small figure of 3,050 litres. A better approximation is obtained by measurements of the thorax volume, subtracting the heart, and the weight of the lungs. For a whale of 22 m. this was estimated at 7,000 litres and I arrive at 14,000 litres as the most probable figure for the 27.2 m. whale. Careful measurements of the thorax volume of whales are among the chief desiderata for physiological calculation. 14,000 litres of air would provide the whale with approximately 2,800 litres of oxygen, or enough for 50 minutes at the estimated metabolism of 53 litres per minute.

Because the possible thorax volume is evidently approximately proportional to total volume or weight of animal, while metabolism is proportional to some fractional power of the weight (probably between $W^{2/3}$ and $W^{3/4}$), large size is essential for the capability of prolonged diving.

LIABILITY OF WHALES TO CAISSON DISEASE

In deep diving, the blood passing through the lungs becomes supersaturated with nitrogen. About 1 volume per cent is taken up for each atmosphere of excess pressure. Supposing the whale to stay 5 minutes at 100 m., the 1,000 litres of blood passing per minute would take up an extra amount of 100 litres, or 500 litres in all. This is, I believe, unavoidable. It seems to me inconceivable that the circulation should stop during deep dives, and any blood passing through the lungs could certainly not avoid taking up the corresponding volume of nitrogen.

A slow circulation will reduce the rate at which nitrogen is absorbed, and to that extent large size is advantageous, but it will reduce also the rate at which the gas is given off at lower pressures. Supposing the whale to take a series of dives to 100 m., staying at the surface each time only for the few seconds necessary to take one or two breaths, a progressive supersaturation of the blood and the tissues seems unavoidable, and, should the whale choose to stay at the surface for some time afterwards, nitrogen bubbles should appear both in the blood and in the tissues—in other words, the whale would be liable to a severe attack of caisson disease.

The striking evidence presented by Laurie shows (to my mind conclusively) that serious supersaturation of tissue fluids with nitrogen does not supervene, but just how it is avoided is by no means clear. Laurie has made the remarkable discovery of the nitrogen-binding *X*-organisms in the blood of whales, but, admitting everything he claims for them, their nitrogen fixation is far too slow to be an essential factor, the more so as the fixation requires oxygen, and in all probability not less and probably more than 1 volume for each volume of nitrogen. The 100 litres assumed to be absorbed per minute at 100 m. would therefore require 100 litres of oxygen, or almost double

the volume necessary for the whale's metabolism proper. This, I think, is outside practical possibilities.

I suspect that the retia mirabilia of blood vessels present in all deep-diving mammals may have something to do with the mechanism for escaping caisson disease, but I must confess my inability to see how. I suggest, however, that our knowledge of these structures, in spite of the valuable anatomical research done upon them, is too incomplete to make speculation worth while, and would point out that a detailed and also quantitative investigation of the retia is highly desirable.

Obituary

MR. WILLIAM BARLOW, F.R.S.

WILLIAM BARLOW was born in Islington, London, on August 8, 1845, and inherited from his father, Frederick Barlow, a business dealing with estate and building property; by the exercise of notable acumen in affairs he realised the business and thus found himself early in life possessed of ample means. Barlow was educated privately; he had a taste for physical science and marked mathematical talent, but cultivated the latter unsystematically and perhaps almost exclusively.

Barlow thus found himself in his early thirties with an independence, with a genius for handling geometrical problems of a particular kind, and with ample leisure to devote to the study of crystal structure, which had become the subject of his choice. He had not, however, received that rigid disciplinary training through which most students of physics and chemistry acquire a broad sense of contemporary knowledge of the physical universe. In some respects this was a hindrance but in others an advantage; it left a powerful intellect unhampered by authority and led a logical mind to pursue its inquiries into difficult and obscure paths which might intimidate the more conventionally trained. Towards 1888 Barlow came into contact with Prof. H. E. Armstrong, from whom he received much encouragement and help; he also met Mr. (now Sir) H. A. Miers and the writer and acquired from them most of his knowledge of formal crystallography. He then spent some time with his family in Germany and made the acquaintance of Paul Groth, the crystallographer, occupying himself with the geometry of crystal structure; on returning to England he devoted himself to crystallographic work of a theoretical kind until his death at Stanmore on February 28, 1934.

In summarising the work of an unconventional genius, it is not easy to proceed chronologically; Barlow did not attack problems in the order which they would naturally take in a textbook or in the present notice. His work on the homogeneous partitioning of space may therefore be first reviewed.

For a century past the view has prevailed that crystal structure consists in the similar repetition throughout space of identical units without regard to their shape or constitution. Continuing earlier work by Bravais and others, Sohncke in 1879 introduced the idea of a regular point-system as one in which the pencils of lines drawn from each point of the system to all the remainder are congruent with each other; the regular point-systems, if classified according to the position and nature of their axes of symmetry (whether screw-axes or axes of rotation), are 65 in number. The 65 Sohncke systems, if built up of mathematical points, do not account for all the types of symmetry represented by the 32 crystal systems; it will be seen at once that the structure of hemimorphous crystals, in which a polar axis is present, cannot be described by a Sohncke system without some further assumption, such as polarity of the points or of the component atoms or molecules.

The development of Sohncke's work needed to provide a complete geometrical theory of crystal structure was undertaken independently by Schönflies, Fedorow and Barlow; all three solved the problem but by different methods, and the line of attack adopted by Barlow may be now briefly indicated. Each Sohncke system is characterised by certain coincidence movements, these being translations and rotations about an axis of symmetry, which leave its appearance unchanged; further, a number of the Sohncke systems are enantiomorphous, that is, not identical with their mirror-images. Barlow duplicated the enantiomorphous Sohncke systems by intercalating the mirror-image in such a way that the coincidence movements of the two component point-systems coincide; he worked out the geometrical methods, three in number, by which this duplication can be effected. The 65 Sohncke systems thus became increased by another 165 to a total of 230; these are known as the 230 space-groups and represent all the types of symmetry possible in crystal structures. Each of the 32 crystal systems corresponds to one or more of these space-groups. With the proof that the space-groups number 230, the geometrical theory of crystal structure becomes