Letters to the Editor

The Editor does not hold himself responsible for opinions expressed by his correspondents. He cannot undertake to return, or to correspond with the writers of, rejected manuscripts intended for this or any other part of NATURE. No notice is taken of anonymous communications.

NOTES ON POINTS IN SOME OF THIS WEEK'S LETTERS APPEAR ON P. 189.

CORRESPONDENTS ARE INVITED TO ATTACH SIMILAR SUMMARIES TO THEIR COMMUNICATIONS.

Origin of the Cosmic Rays

FROM the properties of the kinematic world-models which I have been investigating during the past two and a half years, it can be shown that any unimpeded free particle, at large in inter-galactic space, undergoes acceleration as reckoned by an observer located on any arbitrary nebula, and attains the speed of light at some finite epoch in the experience of that observer. It then decelerates. It can also be shown that at any arbitrary epoch, in any arbitrary domain of inter-galactic space, there will occur some particles possessing velocities arbitrarily close to that of light. If such a particle, of atomic dimensions, happens to undergo a collision during this phase of its trajectory, it will give rise to effects similar to those observed in cosmic ray experiments. I therefore identify the primary agency responsible for cosmic rays with high-speed particles accelerated to the vicinity of the speed of light by the gravitational pull of the rest of the universe¹. The arguments required are purely kinematical, and involve no appeal to any specific theory of gravitation, or any arbitrary hypotheses.

This identification is compatible with the corpuscular character assigned to the primary agency by many authorities. The identification accounts for the observed isotropy; and it provides the origin of the high energies, predicting indeed that there is no upper limit to the energy of a single 'ray'. The energy is drawn from the infinite energy associated with the infinitely many particles constituting the universe. Lastly, the identification removes the old impasse to which other theories of the origin of cosmic radiation have appeared to lead: that if the primary rays were born in the interiors of stars, it is difficult to see how they could ever get out; yet if they were born as a result of multiple collisions in inter-galactic space, it is difficult to see how the inter-galactic density of matter could be high enough.

A full account of these investigations will appear shortly in a volume to be published by the Clarendon Press. The results were first described at Dr. N. V. Sidgwick's colloquium at Oxford on February 19, 1934, and have been discussed in numerous lectures elsewhere.

E. A. MILNE.

19 Northmoor Road,
Oxford.
Jan. 11.

¹ cf. Bothe and Kolhörster, *Z. Phys.*, **56**, 777; 1929.

'Extra' Rings and Bands in Electron Diffraction Patterns

SO-CALLED 'extra' or forbidden rings have been observed from time to time in electron diffraction patterns, and their origin has been attributed to half-order diffractions or impurities. Cases in which more than one such ring have occurred in a single pattern have been rare. We have obtained patterns from platinum, gold, silver, cobalt, nickel, chromium,

iron, tin, graphite and zinc oxide exhibiting not only up to as many as 14 'extra' rings, but also remarkable circular bands with well-defined heads.

The main facts relating to these rings and bands, so far as they have been ascertained, are as follows: (1) The method of production of the specimen is immaterial, but the crystals must be orientated; (2) specimens of similar thickness and method of preparation show much less background if the crystals are orientated than when randomly disposed; (3) for a given orientation the 'extra' ring and band pattern is characteristic of the crystal lattice; for

Fig. 1.

example, all body-centred cubic polycrystalline films with (111) face orientation yield 'extra' rings in the same relative positions; (4) no 'extra' spots have been observed in single crystal patterns; (5) with a suitable film, the thickness being rather critical, a pattern consisting almost wholly of 'extra' rings can be obtained; finally, (6) one head of each band coincides with a normal ring or with the central spot, while the other consists of an 'extra' ring. Fact (1) excludes impurity as a contributory factor, and (5) suggests that secondary scattering plays no material rôle in the formation of 'extra' rings; indeed, it can be shown that a primarily diffracted ray, hkl , on undergoing a second scattering, $h'k'l'$ in the same crystal will give diffracted rays falling on normal pattern rings.

It seems to us that the 'extra' rings and bands owe their origin to the external crystal shape, that is, to the boundary faces through which the electron beam leaves the crystals. For example, the pattern, Fig. 1, was obtained from a face-centred cubic polycrystalline silver film orientated with 110 planes normal to the beam, that is, equivalent to a single crystal rotating about an axis in the beam parallel to a cube-face diagonal. Four distinct 'extra' rings lie within the very intense 111 ring, and two conspicuous bands are