

simultaneous demonstrations of the two systems is temporarily possible with neutral red and Janus green.

The lipin nature of the chondriome makes it disappear in acetic acid or alcohol fixatives unless unsaturated fatty acids are first oxidised to hydroxy acids, less soluble in alcohol and xylol, as by treatment with chromic acid or osmic acid, etc. The

vexed questions of the relation of the chondriome system to the plastids; of the methods of multiplication in embryonic cells of both vacuome and chondriome, and of the relation of the vacuome system to the Golgi network or the Holmgren canals are discussed very concisely and with a full citation of the work upon animal cytology which will be very valuable to the botanist.

Zooplankton of the Great Barrier Reef*

FOLLOWING the general report on the zooplankton of the Great Barrier Reef (noticed in NATURE of August 3, 1935) more detailed work on the different groups is now in progress. The pelagic animals belonging to the Tunicata, Mollusca and Siphonophora are here discussed. Comparison with other expeditions is emphasised, particularly the *Siboga* Expedition, but in this year's work on the Barrier Reef a continuous series of samples is available, giving valuable information as to seasonal distribution and life-histories which has never before been obtained to such an extent. Identification of species has already been made by Miss A. B. Hastings (Tunicata), Mr. A. K. Totton (Siphonophora) and Miss Anne L. Massy (Pteropoda), whilst Prof. Paul Pelseneer has looked through the larval gastropods. Identification was not possible in the last group and in the larval lamellibranchs, but reports on the Tunicata and Siphonophora have already been published (in Vol. 4 of these reports).

The larger tunicates were abundant in the stramin net, but among the great numbers of small animals retained by the silk net they were less important. On the average, the tunicates formed one half of the total number of animals caught in the stramin net, while in the coarse and fine nets, oblique hauls, these proportions were only 5.7 per cent and 2.7 per cent respectively. The total collections comprised three species of Doliolids, six species of Salps and eight species of Appendicularians; but the only species which played an important part in the composition of the plankton at the weekly stations were *Salpa democratica*, *Oikopleura rufescens* and *O. longicauda*. On the average for the year, the Thaliacea far outnumbered the Appendicularians in the stramin catches; but there were violent fluctuations in the composition of the Tunicata due to swarming of certain species, especially *Salpa democratica* and *Megalocercus huxleyi*. In the coarse silk net small appendicularians preponderated, except in months when swarms of salps were met with. In both nets the Doliolids were on the whole of small significance.

Easily the most abundant salp was *Salpa democratica*, with two maxima—September–January, and April–May; during the remaining months it was almost completely absent. On only a few days really large catches were made when the salps were often to be seen in the sea in enormous numbers. The spring increases occurred at full moon, the later increases at or after new moon. The authors are of opinion, however, that the salp increases do not really indicate a lunar periodicity in breeding, but that they indicate successive outbursts of reproduction in the

waters round Low Island, the sudden peaks being due to swarms drifting up the Barrier Reef Lagoon. The chief increases occur when the temperature is between 24° and 28° C. It is probable that salinity has no effect on their presence, whilst temperature is important. The comparatively few salps in the collection may indicate that those caught are the better able to live in shallow coastal water, and the same may hold good for appendicularians, or it may be that in some years other species will be found which, owing to sparse distribution, were not brought near the coast in this one year.

Of the Appendicularians treated as a whole, *Oikopleura rufescens* and *O. longicauda* are the most abundant, the maximum occurring in January and February, coinciding with the highest temperature and decreasing salinity. The Barrier Reef collections are so far the only records giving a full picture of a year's course in a region in the Pacific where warm tropical water conditions exist throughout the year. The absence of *Oikopleura dioica* is striking. Of the Mollusca, the pteropods are much the most important, although there were numbers of larval gastropods and lamellibranchs at times. Here again the pteropod records contain the first contribution towards their seasonal study in the tropics. Only three are important in the coastal waters of the Barrier Reef, *Creseis acicula*, *C. virgula* and *Cavolinia longirostris*, agreeing with the results of the *Siboga* Expedition. The creseids extend normally nearer the coast than the other genera.

Siphonophores were much more abundant than medusae and only a few coral planulae were caught. Of the thirty-two species of Siphonophores, twenty from the Barrier Reef Lagoon, two species predominated, *Lensia subtiloides* and *Diphyes chamissonis*, and these were responsible for the main fluctuations in abundance during the year. Many species were absent because of low salinity during the heavy rains, but they were probably present outside the reef. That the three species *Diphyes chamissonis*, *Lensia subtiloides* and *Enneagonum hyalinus* were present in the lagoon during the period of low salinity points to the possibility that these species are specially adapted to life in shallow coastal waters. This seems to be the first time that any siphonophores have been definitely recognised as coastal species in a predominantly oceanic group.

* British Museum (Natural History). Great Barrier Reef Expedition, 1928–29. Scientific Reports. Vol. 2, No. 7: The Zooplankton. 4: The Occurrence and Seasonal Distribution of the Tunicata, Mollusca and Coelenterata (Siphonophora). By F. S. Russell and J. S. Colman. Pp. 203–276. (London: British Museum (Natural History), 1935.) 5s.