Synaptosomal T₃ Content in Cerebral Cortex of Adult Rat in Different Thyroidal States

Pradip Kumar Sarkar, M.Sc., and Arun Kumar Ray, Ph.D.

The quantitation of thyroid hormone levels in cerebrocortical synaptosomes of the adult rat at altered thyroid states was studied because of the recent idea of the involvement of thyroid hormone in the adult mammalian brain. In contrast to a marked fall in serum T₃ levels in PTU-induced hypothyroid rats, the synaptosomal T₃ content was raised by 9.5 fold in this hypothyroid situation compared to the euthyroid control rats. A single injection of T₃ (2 µg/g) to the hypothyroid

rats lowered the T₃ level to about half of those values obtained in only hypothyroid condition. The T₄ levels remained below the detectable range in all the groups tested. This may be because 70% to 80% of the intraneuronal T₃ comes through the local deiodination of T₄. The present investigation indicates a tendency for synaptosomal reorientation of T₃ level for vital neurophysiologic function in altered thyroid conditions. [Neuropsychopharmacology 11:151–155, 1994]

KEY WORDS: Rat brain; Cerebral cortex; Synaptosomes; Hypothyroid; L-Thyroxine; L-Triiodothyronine.

Although high affinity, low capacity nuclear thyroid hormone receptors have been localized in the adult rat brain, no physiological parameter has been identified that may be accepted as an expression of the thyroid hormone–receptor interaction in this organ (Eberhardt et al. 1978; Schwartz and Oppenheimer 1978; Valcana and Timiras 1978; Dozin-Van Roye and De Nayer 1979; Oppenheimer et al. 1980; Gullo et al. 1987a,b). Recent evidences have shown selective uptake of thyroxine (T₄) and triiodothyronine (T₃), rapid conversion of T₄ to T₃, and specific ¹²⁵I-T₃-binding sites in synaptosomal fraction of nervous tissue in the adult rat brain. This favors direct nonnuclear action of thyroid hormone related to neurotransmission in the adult mammalian brain (Dratman et al. 1976; Dratman and Crutchfield

1978; Kastellakis and Valcana 1989; Hashimoto et al. 1991).

To further support that thyroid hormone is important in the normal neural function of the adult mammalian brain, thyroid hormone concentrations in synaptosome from the adult rat cerebral cortex in normal and different thyroid states have been determined.

MATERIALS AND METHODS

Animal Treatment

Adult male Charles Foster rats (180–200 g body weight [BW]) were maintained in a temperature controlled room (24 ± 1°C) with a 12 h light–dark cycle, and fed *ad libitum* with standard rat diet. Some of the rats were made hypothyroid by daily intraperitoneal (IP) injection of n-propylthiouracil (PTU) (Sigma Chemical Co., USA) at a dose of 2 mg/100 g BW for 14 days. Some of the hypothyroid and euthyroid control rats were injected IP (single) with T₃ (Sigma Chemical Co., USA) at a dose of 2 µg/g BW. Simultaneously, a group of euthyroid control rats were injected with the same volume of vehicle (alkaline 0.9% saline). All the rats were sacrificed after 24 hours of the last injection of T₃, or PTU, or PTU + T₃.

From the Department of Animal Physiology, Bose Institute, Calcutta, India.

Address reprint requests to: Prof. Arun Kumar Ray, Ph.D., Department of Animal Physiology, Bose Institute, P-1/12 CIT Scheme VII-M, Calcutta 700054 India.

Received September 17, 1993; revised April 13, 1994; accepted April 29, 1994.

Table 1. T₄ and T₃ Concentrations in Rat Serum

Group	T ₄ (μg/dl)	T ₃ (ng/ml)
Euthyroid	4.30 ± 0.20	0.69 ± 0.04
Hypothyroid	0.41 ± 0.03 ^a	0.14 ± 0.02 ^a

Results are expressed as mean ± SEM of 3 separate observations. Each observation was made from 6 rats.

^a Significantly different from corresponding value, $p < .001$.

The serum samples were collected and stored at -70°C until use.

Preparation of Synaptosomes

The synaptosomes from the cerebral cortex were prepared according to the method of Sarkar and Ray (1992a). Briefly, the 10% cortical homogenate (in 0.32 M sucrose) was layered in 1.2 M sucrose, after removal of the cell debris and nuclei, and centrifuged at $34,000 \times g$ for 50 min at 4°C . The fraction collected between the 0.32 M sucrose layer and the 1.2 M sucrose layer was diluted at 1:1.5 with ice-cold, double-distilled water, further layered on 0.8 M sucrose, and again centrifuged at $34,000 \times g$ for 30 min. The pellet obtained was washed, repelleted at $20,000 \times g$, for 20 min, and ruptured hypo-osmotically with 5 mM imidazole-HCl buffer at pH 7.4 at 4°C . The ruptured synaptosomal suspension was used for T₄ and T₃ radioimmunoassay (RIA).

Radioimmunoassay

RIA of T₄ and T₃ concentrations in both serum and synaptosomes were measured using RIA kits supplied by the Board of Radiation and Isotope Technology, Bombay, India. All the samples were analysed in duplicate. The sensitivity of the T₃ and T₄ assay was 0.24 ng/ml and 0.5 μg/dl, respectively, of the samples based on 90% B/Bo intercept where B is the corrected average counts of standard/sample, and Bo is the corrected average counts of zero standard. The T₄-assay kit showed

5% cross-reactivity and 0.1% cross-reactivity with T₃ and T₂, respectively. The T₃-assay kit demonstrated a cross-reactivity at 0.1% with both T₄ and T₂. Eighty percent to eighty-five percent of the hormones added to the ruptured synaptosomal suspension were recovered with the RIA kits.

Protein concentration was measured by the method of Vera (1988), using bovine serum albumin as standard.

Statistical Analysis

Results are expressed as mean ± SEM of three separate observations. Each observation was made from six rats. Statistical analysis of the data were made by performing Student's *t* test with $p < .05$ as the significance cutoff point. The multiple groups were also checked with Duncan Multiple Range Test.

RESULTS

Serum T₄ and T₃ Levels

An appreciable fall in the serum T₄ and T₃ levels was observed in hypothyroid animals compared to the control euthyroid values (Table 1).

Synaptosomal Concentrations of T₄ and T₃

In synaptosome, opposite nature of changes were noticed (Table 2). Hypothyroid rats showed a 9.5 fold higher ($p < .001$) amount of T₃ in the cerebrocortical synaptosomes compared to that found in euthyroid synaptosomes. A single injection of T₃ (2 μg/g) to the hypothyroid rats reduced (1.6 fold, $p < .001$) the synaptosomal T₃ level in comparison to the hypothyroid rats. An elevation (2.5 fold, $p < .001$) of the synaptosomal T₃ level was found in euthyroid control rats after a single injection of T₃ (2 μg/g).

The T₄ level remained undetectable in synaptosomes in all the experimental conditions in the present studies.

Table 2. T₃ Concentrations in Rat Cerebrocortical Synaptosomes in Different Thyroid States

T ₃ (ng/mg protein)				F Values (one way ANOVA)
(A) Euthyroid Control	(B) Hypothyroid	(C) Hypothyroid + T ₃ (2 μg/g)	(D) T ₃ (2 μg/g)	
0.43 ± 0.06	4.10 ± 0.06	2.56 ± 0.14	1.07 ± 0.24	163.60 $p < 0.001$

T₃ results are expressed as mean ± SEM of 3 separate observations. Each observation was made from 6 rats. The values between the groups A-B, A-C, A-D, B-C, B-D, and C-D are statistically significant within $r < 0.05$ in Duncan Multiple Range Test.

DISCUSSION

Despite a number of recent evidences that indicated uptake and concentration of ¹²⁵I-T₃ in rat cerebral cortex synaptosomal fraction, and favored a possible role of the thyroid hormone associated with the process of neurotransmission (Heninger and Albright 1975; Dratman et al. 1976; Dratman and Crutchfield 1978; Kastellakis and Valcana 1989; Serrano-Lozano et al. 1991), direct quantitation of the hormone in synaptosomes in different thyroid states was lacking. In the present study, an attempt was made to demonstrate how the T₃ level in synaptosomes alters in response to different thyroidal condition.

Hypothyroid condition showed an appreciable decline in both serum T₄ and T₃ level in rats in a usual way as found by other investigators (Larsen and Frumess 1977; Leonard et al. 1981; Cooper et al. 1983). Although it has been shown earlier that in hypothyroid condition, the whole brain, or different regions of the brain, maintain similar levels of T₃ compared to the euthyroid control rats through increased activity of 5'-deiodinase type II (5'D-II), and corresponding high fractional rate of T₄ to T₃ conversion (Obregon et al. 1978; Vigoroux et al. 1979; Crantz and Larsen 1980; Kaplan and Yaskoski 1980; Leonard et al. 1981; Leonard et al. 1984; Morreale de Escobar et al. 1988; St Germain et al. 1988; Escobar del Rey 1989; Serrano-Lozano et al. 1991), insufficient evidence is available except for a few recent reports (Sarkar and Ray 1992b; Mason et al. 1993) to quantitate the synaptosomal concentration of thyroid hormone. Approximately 8-fold higher concentration of T₃ has been found in synaptosome by Mason et al. (1993) compared to the whole brain in euthyroid rats. Our present observation of approximately 9.5-fold higher T₃ content in synaptosome of hypothyroid rats compared to the euthyroid controls may be the result of a higher fractional rate of T₃ production by increased activity of 5'D-II, and a correspondingly higher selective uptake and concentration of T₃ molecules in the synaptosomes to cope up with the physiological need of thyroid hormone in this tissue at this condition (Crantz and Larsen 1980; Kaplan and Yaskoski 1980). In euthyroid rat brain, selective uptake of ¹²⁵I-T₃ and its concentration in synaptosomal compartment have been demonstrated (Dratman et al. 1976; Dratman and Crutchfield 1978). In addition, the use of hypothyroid animals only after 14 days of PTU treatment, where some adaptive mechanisms still unknown in nature, do not reach the equilibrium as compared to the animals kept in chronic hypothyroid condition for a much longer duration as used by other workers (Kaplan and Yaskoski 1980; Leonard et al. 1981; Morreale de Escobar et al. 1988). This may be another reason for maintaining a high level of synaptosomal T₃ in our hypothyroid rats. Expression of the data in different

forms such as per g organ (brain) basis, or per mg compartmental (synaptosomal) protein basis, as presented in our experiment, also becomes an additive factor for discrepancies among different groups of workers regarding the quantitative aspects of T₃ or T₄ in the brain (Morreale de Escobar et al. 1988; Escobar del Rey et al. 1989; Mason et al. 1993).

The fall in T₃ concentration in synaptosome prepared from T₃-treated hypothyroid rat cerebral cortex may be the result of inhibition of 5'D-II activity after 24 hours of the T₃ administration, in the presence of the considerable amount of exogenous T₃. An inhibition in the activity of 5'D-II has been observed within 4 hours of T₃ treatment to the thyroidectomized rats (Leonard et al. 1981; Leonard et al. 1984). A rise in the synaptosomal T₃ level in the hypothyroid rats, and a fall in the same in the T₃-treated hypothyroid animals after 24 hours of T₃-treatment, also reflects the tendency for a compensatory regulatory mechanism of thyroid hormone metabolism in the adult rat brain in altered thyroid conditions, although the nature of the mechanism remains unknown. T₃-treated control rats have shown higher levels of synaptosomal T₃, compared to the control values. This may be a result of the extra T₃ transport influenced by a high dose of exogenously administered T₃ (2 µg/g) in a similar way as observed by Dratman et al. (1976, 1978), and Kastellakis and Valcana (1989).

In our experiment, the T₄ level in synaptosomes remained undetected in untreated control groups and other experimental rat groups. This may reflect a state of rapid conversion of T₄ to T₃ in the brain by 5'D-II enzyme (Dratman and Crutchfield 1978; Obregon et al. 1978; Vigoroux et al. 1979; Crantz and Larsen 1980; Kaplan and Yaskoski 1980). Other researchers (Dratman et al. 1976; Dratman and Crutchfield 1978) have already shown that after IV administration of radiolabeled T₄ and T₃, the hormone is concentrated as T₃ in a synaptosomal fraction of the whole rat brain, and T₄ to T₃ conversion occurs very rapidly within the nerve cells. T₃ formed in the neuronal cell body then may be translocated down the axon to the synaptic ends. Kastellakis and Valcana (1989) also demonstrated a saturable and nonsaturable uptake of T₃ and T₄, respectively, in isolated synaptosomes in an *in vitro* model indicative of a two-component T₃-uptake system.

The procedure we developed for the direct radioimmunoassay of T₃ and T₄ in synaptosomes using a kit also appears as an alternative method for the thyroid hormone assay without prior extraction of the hormone as followed in other conventional RIA methods. As the RIA kit reagents also contained 8-anilino-naphthosulphonic acid that makes T₃ and T₄ free from the protein bound form, the chances for nonspecific binding of the hormone with synaptosomal protein materials are minimum. In addition, the data regarding cross-reactivity,

sensitivity, and recovery in our RIA system may support it to be a viable alternate, direct assay method.

The data emerged from our study reveal the quantitative aspects of involvement of T_3 in synaptosomes in different thyroid states, and favors its role in neuronal functions as formerly described (Dratman et al. 1976, 1978; Hashimoto et al. 1991). A stimulation of synthesis of synapsin-I protein (related to neurotransmission) by T_3 in the developing brain has been reported (Salemi et al. 1990). Although, the synaptosomal T_3 levels varied widely with different treatments, our results illustrate a unique, but unknown regulatory mechanism of the thyroid hormone metabolism in the mature mammalian brain.

ACKNOWLEDGMENT

Thanks are due to Council of Scientific and Industrial Research, New Delhi, India, for financial assistance to P.K.S.

REFERENCES

- Cooper DS, Kieffer JD, Halpern R, Saxe V, Mover H, Maloof F, Ridgway EC (1983): Propylthiouracil (PTU) pharmacology in the rat. I. Effect of PTU on thyroid function. *Endocrinol* 113(3):921-928
- Crantz FR, Larsen PR (1980): Rapid thyroxine to 3,5,3'-triiodothyronine conversion and nuclear 3,5,3'-triiodothyronine binding in rat cerebral cortex and cerebellum. *J Clin Invest* 65:935-938
- Dozin-van Roye B, De Nayer P (1979): Nuclear triiodothyronine receptors in rat brain during maturation. *Brain Res* 177:551-554
- Dratman MB, Crutchfield FL, Axelrod J, Colburn RW, Thoa N (1976): Localization of triiodothyronine in nerve ending fractions of rat brain. *Proc Natl Acad Sci (USA)* 73:941-944
- Dratman MB, Crutchfield FL (1978): Synaptosomal [125 I]triiodothyronine after intravenous [125 I]thyroxine. *Am J Physiol* 235(6):E638-E647
- Eberhardt NL, Valcana T, Timiras PS (1978): Triiodothyronine nuclear receptors: An *in vitro* comparison of the binding of triiodothyronine to nuclei of adult rat liver, cerebral hemisphere, and anterior pituitary. *Endocrinol* 102:556-563
- Escobar del Rey F, Ruiz de Ona C, Bernal J, Obregon MJ, Morreale de Escobar G (1989): Generalized deficiency of 3,3',5'-triiodo-L-thyronine (T_3) in tissues from rats on a low iodine intake, despite normal circulatory T_3 levels. *Acta Endocrinol (Copenh)* 120:490-498
- Gullo D, Sinha A, Woods R, Pervin K, Ekins R (1987a): Triiodothyronine binding in adult rat brain: Compartmentation of receptor populations in purified neuronal and glial nuclei. *Endocrinol* 120:325-331
- Gullo D, Sinha AK, Bashir A, Hubank M, Ekins RP (1987b): Differences in nuclear triiodothyronine binding in rat brain cells suggest phylogenetic specialization of neuronal functions. *Endocrinol* 120:2398-2403
- Hashimoto H, Walker CH, Prange Jr AJ, Mason GA (1991): The effects of thyroid hormones on potassium stimulated release of 3 H-GABA by synaptosomes of rat cerebral cortex. *Neuropsychopharmacol* 5(11):49-54
- Heninger RW, Albright EC (1975): Alteration in tissues and serum concentrations of TSH, iodide, T_4 , and T_3 induced by various dietary iodide levels. *Proc Soc Exptl Biol Med* 150:137-141
- Kaplan MM, Yaskoski KA (1980): Phenolic and tyrosyl ring deiodination of iodothyronines in rat brain homogenates. *J Clin Invest* 66:551-562
- Kastellakis A, Valcana T (1989): Characterization of thyroid hormone transport in synaptosomes from rat brain. *Mol Cell Endocrinol* 67:231-241
- Larsen PR, Frumess RD (1977): Comparison of the biological effects of thyroxine and triiodothyronine in the rat. *Endocrinol* 100:980-988
- Leonard JL, Kaplan MM, Visser TJ, Silva PR (1981): Cerebral cortex responds rapidly to thyroid hormones. *Science* 214:571-573
- Leonard JL, Silva JE, Kaplan MM, Mellon SA, Visser TJ, Larsen PR (1984): Acute post-transcriptional regulation of cerebrocortical and pituitary iodothyronine 5'-deiodinase by thyroid hormone. *Endocrinol* 114:998-1004
- Morreale de Escobar G, Obregon MJ, Ruiz de Ona C, Escobar del Rey F (1988): Transfer of thyroxine from the mother to the rat fetus near term: Effects on brain 3,5,3'-triiodothyronine deficiency. *Endocrinol* 122:1521-1531
- Mason GA, Walker CH, Prange Jr AJ (1993): L-triiodothyronine: Is this peripheral hormone a central neurotransmitter? *Neuropsychopharmacol* 8:253-258
- Obregon MJ, Morreale de Escobar G, Escobar del Rey F (1978): Concentrations of triiodothyronine in the plasma and tissue of normal rats as determined by radioimmunoassay: Comparison with results obtained by an isotopic equilibration technique. *Endocrinol* 103:2145-2153
- Oppenheimer JH, Schwartz HL (1980): Thyroid hormone action at the nuclear level: In Hennemann G (ed), *Thyroid Hormone Metabolism*, New York, Basel, Marcel Dekker pp. 383-415
- Salemi G, Ferraro D, Savettieri G (1990): Triiodothyronine accelerates the synthesis of synapsin I in developing neurones from fetal rat brain cultured in a synthetic medium. *Neurochem Res* 15(8):827-831
- Sarkar PK and Ray AK (1992a): A simple biochemical approach to differentiate synaptosomes and non-synaptic mitochondria from rat brain. *Meth Find Exptl Clin Pharmacol* 14(7):493-497
- Sarkar PK and Ray AK (1992b): Synaptosomal T_3 concentration in altered thyroid states in rat, 61st Annual Conference of Society of Biological Chemists (India), Section Endocrinology and Reproductive Biology, Abstract No 174
- Schwartz HL, Oppenheimer JH (1978): Nuclear triiodothyronine receptor sites in brain: Probable identity with hepatic receptors and regional distribution. *Endocrinol* 103:267-273
- Serrano-Lozano A, Morata P, Montiel M, Perez-Aguiler JP,

- Morell M (1991): L-thyroxine and L-triiodothyronine in the cerebral tissues of the adult rat: Effect of hypothyroidism. *Rev Esp Fisiol (Spain)* 47(3):141-145
- St Germain DL (1988): The effects and interaction of substances, inhibitors and the cellular thiol-disulfide balance on the regulation of type II iodothyronine 5'-deiodinase. *Endocrinol* 122(5):1860-1868
- Valcana T, Timiras PS (1978): Nuclear triiodothyronine receptors in the developing rat brain. *Mol Cell Endocrinol* 11:31-41
- Vera JC (1988): Measurement of microgram quantities protein by a generally applicable turbidimetric procedure. *Anal Biochem* 174:187-196
- Vigorous E, Clos J, Legrand J (1979): Uptake and metabolism of exogenous and endogenous thyroxine in brain of young rats. *Horm Metab Res* 11:228-232