

Enhancement of Postsynaptic Sensitivity to Dopaminergic Agonists Induced by Neonatal Hippocampal Lesions

Rui-Qian Wan, Ph.D., and Roy Corbett, Ph.D.

The effects of neonatal hippocampal lesions on behavioral responsiveness to dopaminergic agonists and antagonists were examined. The ventral hippocampus was damaged bilaterally using ibotenic acid on postnatal day 7, and locomotor responses to dopaminergic agonists and antagonists were evaluated on postnatal day 35 (PD35), 56 (PD56), and 70 (PD70). Quinpirole (0.06, 0.125, 0.25, and 0.5 mg/kg SC), but not SKF38393 (5 and 10 mg/kg SC), increased locomotion in a dose-dependent manner in control and lesioned groups on PD35 and PD56. However, lesioned rats displayed a greater behavioral response to quinpirole than controls at the doses of 0.25 and 0.5 mg/kg on both

PD35 and PD56. Amphetamine (1.5 mg/kg IP) increased locomotor activity in both groups on PD70, but this effect was greater in lesioned rats than in controls. Raclopride (0.25 and 0.5 mg/kg SC) and SCH23390 (0.01 and 0.02 mg/kg SC) blocked the amphetamine-induced hyperlocomotion in the lesioned and control groups. These results suggest that neonatal hippocampal lesion-induced behavioral hyperresponsiveness to amphetamine is likely related to an increased postsynaptic sensitivity of the D₂ subtype of receptors. [Neuropsychopharmacology 16:259–268, 1997] © 1997 American College of Neuropsychopharmacology

KEY WORDS: Amphetamine; Dopamine D₁; Dopamine D₂; Hippocampus; Locomotion; Neonatal lesions; Neurotoxin

Abnormal neurodevelopmental processes may be involved in some aspects of schizophrenia (Weinberger 1987; Mednick et al. 1991; Kerwin and Murray 1992; Bloom 1993). Clinical studies have suggested that regional neuropathology during early development may result in behavioral abnormalities in later life (Weinberger 1987; Bloom 1993; Kotrla and Weinberger 1994). Neuropathological changes in the hippocampus have been described in schizophrenic brains (Brown et al. 1986; Falkai and Bogerts 1986; Beckmann et al. 1987; Bo-

gerts 1993; Randolph et al. 1993; Hall et al. 1994). The hippocampus is anatomically interconnected with many structures in the mesolimbic and mesocortical dopamine (DA) systems (Groenewegen et al. 1991). Hippocampal glutamatergic efferents and dopaminergic efferents from the ventral tegmental area (VTA) converge onto the same neurons in the nucleus accumbens (NAc) (Totterdell and Smith 1989; Groenewegen et al. 1991). This hippocampal-accumbens interaction has been postulated to play an important role in modulating motor behaviors. Rats with hippocampal lesions become highly susceptible to environmental stimulation and psychomotor stimulants such as amphetamine. Thus, hippocampal lesions have been used to produce an animal model for the study of schizophrenia (McKinney and Moran 1981; Schmajuk 1987; Grey et al. 1991; Whishaw and Mittleman 1991).

A series of studies examining the effects of neonatal hippocampal lesions on behavior at different ages in rats has been reported (Lipska and Weinberger 1993;

From Neuroscience PGU, Hoechst Marion Roussel, Inc., Route 202-206, P.O. Box 2500, Somerville, NJ.

Address correspondence to: Dr. Rui-Qian Wan, Biological Psychiatry Branch, National Institute of Mental Health, Bldg. 10/3N212, 9000 Rockville Pike, Bethesda, MD 20892.

Received September 11, 1995; revised August 5, 1996; accepted August 14, 1996.

Lipska et al. 1993, 1995a, 1995b). Neonatal lesions of the hippocampus were found to induce delayed behavioral impairments that resulted in a hyperresponsiveness to stress and amphetamine at postnatal day 56 (PD56), but not at postnatal day 35 (PD35; Lipska et al. 1993). In rats PD35 and PD56 correspond to puberty and early adulthood, respectively (Ojeda et al. 1980). A previous *in vivo* microdialysis study of freely moving rats in our laboratory indicated that although amphetamine induced greater locomotion in rats with neonatal hippocampal lesions than in controls, the level of extracellular DA in the NAc was similar in both lesioned and control rats (Wan et al. 1996). This dissociation between an enhanced behavioral responsiveness and a lack of increase in DA release from presynaptic terminals suggested that an increased postsynaptic sensitivity may play an important role in these behavioral impairments. In addition, another study indicated that rats with neonatal hippocampal lesions were behaviorally hyperresponsive to apomorphine, a nonselective DA receptor agonist, at puberty and early adulthood (Lipska and Weinberger 1993). However, the role of the specific subtype of DA receptors in this lesion-induced hypersensitivity has not been identified. Thus, the present study was designed to identify the specific DA receptor subtype that is involved in the mediation of behavioral hyperresponsiveness induced by neonatal hippocampal lesions. Our experiments examined the lesion-induced effect on locomotor responsiveness to D_1 and D_2 receptor agonists and antagonists at different ages.

MATERIALS AND METHODS

Subjects

Pregnant Sprague-Dawley rats (Charles River) were obtained at 12 to 15 days of gestation. The rats were housed individually and maintained under the standard laboratory conditions as outlined in the "NIH Guide for the Care and Use of Laboratory Animals" (National Institute of Health Publications, No. 85-23, revised 1985) with 12-hour dark-light cycle. Food and water were available *ad libitum*. Male pups in each litter were used for the experiments. Pups were weaned on postnatal day 25 and grouped two to three per cage according to their surgical status. To maintain a mixed sexual environment for growing pups, two female pups were kept in each litter until weaning. There were 128 male rats included in this study, which were from approximately 24 litters.

Surgery

Neonatal hippocampal lesions were produced in male pups on postnatal day 7 (PD7). Litters that had four to six male pups were selected for the experiments. The

half of the pups in each litter was randomly assigned to the lesioned group. The other half of the pups was the control group.

Bilateral lesions of the ventral hippocampus were performed using a method described previously (Lipska et al. 1993; Wan et al. 1996). Briefly, each of the anesthetized pups (hypothermia on wet ice for about 15 min) was placed on a platform that was fastened on a stereotaxic apparatus. The head of the pup was loosely fixed on the platform and the skull was kept flat. After an incision was made in the skin, two injectors made of 32-gauge hypodermic tubing were used to deliver drug solution or vehicle. Ibotenic acid (IBO, Sigma, St. Louis, MO) was prepared in 0.1 M phosphate buffered saline (PBS, pH 7.4) in the concentration of 10 $\mu\text{g}/\mu\text{l}$. A volume of 0.3 $\mu\text{l}/\text{side}$ of IBO (lesions) or the same volume of PBS (controls) was infused bilaterally into the ventral hippocampus using the stereotaxic coordinates AP -3.0 , ML ± 3.5 , and DV -4.0 mm relative to bregma. The flow rate of infusion was 0.15 $\mu\text{l}/\text{minute}$. After completion of infusion the injectors remained in the places for another 4 minutes. The incision was closed with two clips. The pup was then placed on a warm water heating pad (American Medical Systems, Cincinnati, OH) at 37° C and returned to its mother after its body temperature recovered.

Locomotor Apparatus and Behavioral Assessments

Each locomotor chamber (clear polycarbonate cage, 45 \times 22 \times 20 cm) was equipped with three photoelectric light sources spaced at 18-cm intervals and 1 cm above the grid floor to record horizontal movements. Consecutive interruptions of two separate light beams were registered as activity counts and summarized every 5 minutes by the microprocessor-based control system.

The behavioral tests examined spontaneous exploration, drug-influenced locomotor activity, and stereotyped behavior. For each rat, spontaneous locomotion was recorded during the first 60 minutes. At the end of this period, the rat was injected with either drug or vehicle and returned immediately to the chamber for another 60 minutes. All behavioral assessments took place between 9:00 A.M. and 2:00 P.M.. Stereotyped behaviors were assessed for 30 s every 10 minutes following injection. Their overall stereotypy was rated on a modified version of the scale that was originally described elsewhere (Maclennan and Maier 1983; Mittleman et al. 1993): 0, no stereotypy; 1, intermittent stereotypy; 2, continuous stereotypy over a wide area; and 3, continuous stereotypy in a restricted area.

Drugs and Experimental Groups

d-Amphetamine (Sigma, St. Louis, MO), (–)-Quinpirole [Research Biochemicals International (RBI), Nat-

ick, MA], (\pm)-SKF-38393 (RBI), SCH-23390 (RBI), and Raclopride (ASTRA, Sodertalje, Sweden) were prepared in saline and administered in an injection volume of 1 ml/kg. The doses of drugs used in the present study were selected based on the previous studies and our own preliminary experiments (Mittleman et al. 1993; Petry et al. 1993; Wan et al. 1996).

On PD35, rats were examined for their behavioral responses to quinpirole (0 (saline), 0.06, 0.125, 0.25, and 0.5 mg/kg SC) and SKF (5 and 10 mg/kg SC). Rats in the same lesion status were randomized to each group of the different drugs and doses ($n = 8$ at each of the doses for every drug). Rats from the same litter were randomly distributed to different groups.

On PD56, all of the rats were examined again for their behavioral response to quinpirole and SKF at the same doses used on PD35. Groups of the different drugs and doses ($n = 8$ /group) were formed randomly from rats in the same lesion status. However, the new assignment was made so that each of the rats was tested with a drug or a dose different from that it had received on PD35. Again, rats from the same litter were assigned to different groups. In addition, a new group of controls and a group of lesioned rats ($n = 8$ /group) were added to examine behavioral responses to quinpirole (0.125 mg/kg) in combination with SKF (10 mg/kg). SKF was injected 10 minutes prior to quinpirole. These two groups were not tested previously.

On PD70, six groups of controls and six groups of lesioned rats ($n = 8$ /group) were randomly formed from rats that had been tested on PD35 and PD56. This experiment examined the effects of raclopride (0.25 and 0.5 mg/kg SC) and SCH (0.01 and 0.02 mg/kg SC) on amphetamine-induced (1.5 mg/kg IP) locomotion. Raclopride or SCH was injected 10 minutes prior to the injection of amphetamine.

Histology

At the end of the behavioral assessments, each rat was anesthetized with Nembutal (pentobarbital sodium injection, USP) and perfused transcardially with 4% formalin-PBS solution. The brain was removed from the skull. After further fixation in formalin solution, the brain was sliced with a freezing cryostat. The sections through the damaged area were saved, mounted, and stained with cresyl violet. Each section was examined under a microscope to determine the location and extent of lesions.

Statistics

Repeated-measures analysis of variance (ANOVA) was used to analyze behavioral data. Neuman-Keuls post-hoc comparisons and Dunnett's *t*-tests were used when

appropriate to identify a significant difference between groups.

RESULTS

Histology

Bilateral damage to the hippocampus occurred consistently in all rats. The place and size of lesions were similar to that of our previous study (Wan et al. 1996). Figure 1 provides an example of neonatal hippocampal damage. Substantial lesions were observed in the ventral hippocampus that included the CA1, CA3, and dentate gyrus. The lesions were indicated by cavitation, cell loss, and neuroglial proliferation (Figure 2). The dorsal part of the hippocampus, including the CA1, CA3, and dentate gyrus, was generally spared in most of the rats. Isolated small pieces of the ventral dentate, CA3, or CA1 were spared in some rats. Damage to the subiculum was indicated in the posterior portion of the hippocampus in most of the rats. Small damage to the entorhinal and piriform cortices was present in a few rats. However, damage to the adjacent thalamic areas was not detected.

Behavior

Spontaneous Exploration. Spontaneous exploration as indicated by locomotor activity before injections was not different between control and lesioned groups on PD35 and PD56. An example of similar spontaneous locomotion between two groups is illustrated in Figure 3.




Figure 1. Coronal sections that indicate the place and size of neonatal hippocampal lesions produced by ibotenic acid. Sections were stained with cresyl violet. Anteroposterior atlas coordinates for these coronal sections (from left to right) correspond roughly to -4.30 to -5.30 mm posterior to bregma (Paxinos and Watson 1986).




Figure 2. Cresyl violet-stained coronal section that indicates ibotenic acid-induced lesions in the hippocampus. Damage is indicated by cavitation, cell loss, and neuroglial proliferation. *Arrows*, focal neuronal loss and neuroglial proliferation in the CA2 and CA3 areas. The major part of the dorsal hippocampus is generally spared.

This similar spontaneous exploration between the control and lesioned groups was indicated in the experiments that examined the effects of raclopride and SCH on amphetamine-induced locomotor activity on PD70 as well.

Effect of Quinpirole. Administration of saline did not influence locomotor activity in either group on PD35 and PD56 (Figure 3). Quinpirole increased the locomotor activity for both groups in a dose-dependent manner (dose, PD35: $F_{4,70} = 16.73$, $p < .001$; PD56: $F_{4,70} = 10.25$, $p < .001$). At the lowest dose (0.06 mg/kg), quinpirole suppressed locomotion in both groups relative to saline on both PD35 and PD56 (drug, PD35: $F_{1,28} = 21.54$, $p < .01$; PD56: $F_{1,28} = 27.66$, $p < .01$). Hippocampal lesions influenced the locomotor response to quinpirole (lesion, PD35: $F_{1,70} = 11.78$, $p < .01$; PD56: $F_{1,70} = 8.86$, $p < .01$; lesion \times dose ($F_{4,70} = 3.32$, $p < .05$). At the doses of 0.25 and 0.5 mg/kg, the lesioned group had greater locomotor activity than the control group on PD35 and PD 56 (Figure 3).

Intermittent stereotyped behavior occurred only following the highest dose of quinpirole (0.5 mg/kg). The frequency of sniffing and rearing behaviors increased

on both PD35 and PD56. However, the scores of stereotyped behavior in the lesioned group did not differ from the control group. Both groups exhibited a significant effect of time for the occurrence of stereotyped behavior during the 60 minutes following injection (time, $F_{5,140} = 6.03$, $p < .01$; Figure 3, inserts).

Effect of SKF-38393. SKF alone at two doses (5 and 10 mg/kg) did not significantly alter locomotor activity or stereotyped behavior in either group on PD35 and PD56. Lesioned rats did not display a behavioral hypersensitivity to SKF as compared to controls (data not shown).

Effect of Quinpirole in Combination with SKF-38393. Quinpirole (0.125 mg/kg) in combination with SKF (10 mg/kg) markedly enhanced locomotor activity in the lesioned group but not the control group (lesion, $F_{1,28} = 4.99$, $p < .05$; lesion \times drug, $F_{1,28} = 5.02$, $p < .05$; lesion \times drug \times time, $F_{11,308} = 2.35$, $p < .01$, Figure 4). For the lesioned group, quinpirole (0.125 mg/kg) or SKF (10 mg/kg) alone did not significantly increase locomotion compared to saline, although the quinpirole-induced locomotor activity did show a significant effect over time compared to saline (time, $F_{11,154} = 3.41$, $p < .001$; drug \times time, $F_{11,154} = 4.22$, $p < .001$). Following combined administration of quinpirole and SKF, the locomotor activity of lesioned rats was higher than saline (drug, $F_{1,14} = 5.74$, $p < .05$; drug \times time, $F_{11,154} = 6.06$, $p < .001$) and SKF (drug, $F_{1,14} = 4.78$, $p < .05$; drug \times time, $F_{11,154} = 5.01$, $p < .001$). However, the locomotion induced by the combined injection did not significantly differ from that induced by quinpirole alone.

Stereotyped behavior was observed only following injections of quinpirole in combination with SKF (drug, $F_{3,56} = 38.04$, $p < .001$; time, $F_{5,280} = 14.65$, $p < .01$; drug \times time, $F_{15,280} = 14.66$, $p < .01$, Figure 4, inserts). Stereotypy occurred more frequently in the lesioned group than the control group (lesion, $F_{1,56} = 26.84$, $p < .01$; lesion \times drug \times time, $F_{15,280} = 9.41$, $p < .01$).

Effect of Raclopride on Amphetamine-Induced Locomotion. Amphetamine (1.5 mg/kg) enhanced locomotor activity in both groups (drug, $F_{1,28} = 50.55$, $p < .001$, Figure 5). The lesioned group had a higher locomotor response to amphetamine than the control group (lesion, $F_{1,28} = 4.78$, $p < .05$; lesion \times drug, $F_{1,28} = 6.12$, $p < 0.05$; lesion \times drug \times time, $F_{11,308} = 5.02$, $p < .001$). Raclopride at both doses (0.25 and 0.5 mg/kg) abolished amphetamine-enhanced locomotion in both groups. For both groups, the locomotor activity following injections of raclopride in combination with amphetamine was significantly lower than that treated with amphetamine alone (p 's $< .001$), and no difference was observed compared to saline. In lesioned rats, the locomotor activity following combined amphetamine and raclopride injection at the lower dose (0.25 mg/kg) increased during the later test session, which was indicated by a significant time effect during the




Figure 3. Effect of neonatal hippocampal lesions on quinpirole-induced locomotion (*main graphs*) and stereotyped behavior (*inserts*) on PD35 and PD56. Left panel, behavioral responses examined on PD35; right panel, behavioral responses examined on PD56; symbols (Open circles, controls; solid circles, lesions), mean; vertical bars, SEM. Vertical arrows, time of injection. ** $p < .01$; * $p < .05$ versus controls.

60-min period (time, $F_{11,154} = 2.83$, $p < .01$; drug \times time, $F_{11,154} = 5.92$, $p < .001$, Figure 5, right panel).

Stereotyped behavior was found only following administration of amphetamine alone in both groups (drug, $F_{3,56} = 115.71$, $p < .001$; time, $F_{5,280} = 25.73$, $p < .001$; drug \times time, $F_{15,280} = 25.73$, $p < .001$, Figure 5, inserts). The score of stereotypy was higher in the lesioned group than the control group (lesion, $F_{1,56} = 20.23$, $p < .001$; lesion \times drug \times time, $F_{15,280} = 6.25$, $p < .01$). Raclopride at both doses suppressed stereotyped behavior induced by amphetamine.

Effect of SCH-23390 on Amphetamine-Induced Locomotion. SCH at both doses (0.01 and 0.02 mg/kg) also abolished amphetamine-induced hyperlocomotion in both groups (p 's $< .001$, Figure 6). The locomotor activity following SCH in combination with amphetamine did not differ from saline.

SCH at both doses also completely suppressed the amphetamine-induced stereotyped behavior (Figure 6, inserts). Rats injected with the higher dose of SCH (0.02 mg/kg) were inactive during almost the entire test session.

DISCUSSION

Enhancement of Postsynaptic Sensitivity Induced by Neonatal Hippocampal Lesions

Hippocampal-accumbens interaction has been identified as important in modulating psychomotor activity. The effects of hippocampal lesions on DA release from presynaptic terminals in the NAc have been examined using *in vivo* microdialysis. Bilateral hippocampal lesions made in adult rats induced locomotor hyperresponsiveness to amphetamine, and this behavioral impairment was related to an increased amphetamine-stimulated DA release in the NAc (Wilkinson et al. 1993). However, our previous microdialysis study in freely moving rats indicated that neonatal hippocampal lesions had no effect on amphetamine-induced DA release in the NAc at PD56, although lesioned rats did show a higher locomotor response to amphetamine than did controls (Wan et al. 1996). These results suggest that neonatal hippocampal lesion-induced behavioral hyperresponsiveness to amphetamine may be related to a postsynaptic DA receptor supersensitivity.




Figure 4. Effect of neonatal hippocampal lesions on locomotion (*main graphs*) and stereotyped behavior (*inserts*) induced by quinpirole (triangles, 0.125 mg/kg) or SKF38393 (squares, 10 mg/kg) alone versus coadministration of quinpirole and SKF38393 on PD56. *Left panel*, behavioral responses in control rats; *right panel*, behavioral responses in lesioned rats; (open circles, saline) symbols, mean; vertical bars, SEM. ***p* < .01; **p* < .05 versus control rats that received coadministration. See text for other comparisons among treatment.

The present study provides evidence for enhanced postsynaptic DA receptor sensitivity following neonatal hippocampal lesions. The results indicated that quinpirole at the low dose (0.06 mg/kg) decreased locomotion,

which was likely due to stimulation of presynaptic autoreceptors (Imperato et al. 1988; Eilam and Szechtman 1989; Barnes et al. 1990; Mogenson and Wu 1991), whereas higher doses (0.25 and 0.5 mg/kg) increased




Figure 5. Effect of neonatal hippocampal lesions on locomotion (*main graphs*) and stereotyped behavior (*inserts*) induced by amphetamine (solid circles, 1.5 mg/kg) alone versus coadministration of amphetamine (1.5 mg/kg) and raclopride (squares, 0.25 mg/kg; triangles, 0.5 mg/kg) on PD70. *Left panel*, behavioral responses in control rats; *right panel*, behavioral responses in lesioned rats; (open circles, saline) symbols, mean; vertical bars, SEM. ***p* < .01; **p* < .05 versus control rats that received amphetamine. See text for other comparisons among treatments.




Figure 6. Effect of neonatal hippocampal lesions on locomotion (*main graphs*) and stereotyped behavior (*inserts*) induced by amphetamine (*solid circles*, 1.5 mg/kg) alone versus coadministration of amphetamine and SCH23390 (*squares*, 0.01 mg/kg and *triangles*, 0.02 mg/kg) on PD70. *Left panel*, behavioral responses in control rats; *right panel*, behavioral responses in lesioned rats; *open circles*, saline; symbols, mean; vertical bars, SEM. ****p** < .01; ***p** < .05 versus control rats that received amphetamine. See text for other comparisons among treatments.

locomotor activity. These results were consistent with previous reports that quinpirole produced a biphasic effect on locomotion with respect to dose and time course (Eilam and Szechtman 1989; Van Hartesveldt et al. 1992). Of particular importance, however, was the fact that rats with neonatal hippocampal lesions had a behavioral hypersensitivity to quinpirole. This neonatal lesion-induced behavioral impairment was observed in rats at both PD35 and PD56.

The increase in locomotion induced by higher doses of quinpirole may be attributable to its postsynaptic stimulation (Van Hartesveldt et al. 1992; Braun et al. 1993; Yue et al. 1994). One explanation for quinpirole-induced hyperlocomotion could be the production of an active metabolite based on the late onset of hyperactivity following systemic injection of quinpirole (Gonzalez-Lima et al. 1987; Eilam and Szechtman 1989). Previous studies, however, have suggested that quinpirole at higher doses may directly stimulate postsynaptic D₂ receptors. The NAc is one of the regions that has a high density of D₂ receptors in the rat brain (Gehlert and Wamsley 1985; Gehlert et al. 1992). Intraaccumbens injections of quinpirole produced an early decrease followed quickly by an increase in locomotor activity (Van Hartesveldt et al. 1992). This behavioral biphasic effect induced by quinpirole was similar to that produced by intraaccumbens injections of DA with respect to the time course of the behavioral response (Wachtel et al.

1979). The similarity in the behavioral profiles suggests that quinpirole and DA are alike in their direct stimulation of pre- and postsynaptic DA receptors.

Although our results from microdialysis and behavioral studies suggest that neonatal hippocampal lesions increase the sensitivity of postsynaptic DA receptors, it remains unclear whether this hypersensitivity is the result of an increase in the density of DA receptors. Two independent studies of DA receptor binding using similar neonatal hippocampal lesions have reported inconsistent results. One study indicated that the density of D₃ receptors was lower in the NAc and that the density of D₁ receptors was higher in the caudate putamen, with no change in D₂ receptors in lesioned rats (Flores et al. 1996). The other study showed that the lesions did not alter D₂/D₃ and, D₄ receptor densities in the dorsolateral and ventrolateral striatum and the NAc (Knable et al. 1994). However, this latter study might not rule out an alteration in D₃ receptors because the study did not distinguish between the D₂ and D₃ receptor subtypes. A decrease in D₃ receptors could have been masked by changes in D₂ receptors. Postsynaptic D₃ receptors have been shown to be involved in the modulation of locomotor activity. U99194A, a D₃ antagonist, increases locomotion, suggesting that D₃ receptors may exert an inhibitory influence on psychomotor activity (Waters et al. 1993, 1994). Moreover, previous binding studies have shown that quinpirole and raclopride bind to the

D₃ receptor subtype in addition to the D₂ receptor subtype (Gehlert et al. 1992; Seeman and Van Tol 1994). These data suggest that quinpirole-induced hyperlocomotion in neonatal hippocampally lesioned rats may be related not only to a stimulation of D₂ receptors but also to a decrease in the inhibitory effect of D₃ receptors.

Roles of D₁ and D₂ Receptors in Lesion-Induced Hypersensitivity

Increased sensitivity of the D₂ subtype of DA receptors may play an important role in neonatal hippocampal lesion-induced behavioral impairments. Two results in this study support this conclusion. First, the lesioned group showed a behavioral hypersensitivity to quinpirole indicated by hyperlocomotion induced by quinpirole alone in a dose-dependent manner in lesioned rats relative to controls. In contrast, rats with neonatal hippocampal lesions did not show behavioral hypersensitivity to SKF that alone had no significant effect on locomotion in either group. The dosages of SKF (5 and 10 mg/kg) were found in previous studies to be sufficient to produce effects of SKF on locomotor or stereotyped activities (Eilam et al. 1992; Mittleman et al. 1993). However, we cannot rule out the possibility that SKF alone at a higher dose than 10 mg/kg might induce differential effects on behavioral responses in lesioned rats compared to controls. Second, raclopride diminished the amphetamine-induced hyperlocomotion and stereotyped behavior. Evidence has suggested that D₂ receptors may play a primary role in mediating activity of hippocampal accumbens neurons. A previous study indicated that iontophoretic applications of quinpirole or the D₂ antagonist sulpiride, but not the D₁ antagonist SCH23390, altered excitability of hippocampal accumbens neurons (Yang and Mogenson 1986). Thus, it is likely that the sensitivity of D₂ receptors in the hippocampal accumbens system may be altered because of a degeneration of hippocampal projections to the NAc following neonatal hippocampal lesions. Similar impairments of D₂-mediated behaviors have been demonstrated in rats with hippocampal lesions made in adulthood (Mittleman et al. 1993).

Even though D₁ and D₂ receptors can produce opposite effects on some behavioral and neurochemical responses (Eilam et al. 1991, 1992), activation of both D₁ and D₂ receptors has been shown to be necessary to achieve the full range of expression of DA-mediated behaviors (Clark and White 1987; Waddington 1989; Walters et al. 1987). Previous studies have indicated that coadministration of D₁ and D₂ agonists produced synergistic effects on behavioral responses, effects that could not be induced by the activation of either receptor alone (Amalric et al. 1986; Barone et al. 1986; Molly et al. 1986; Robertson and Robertson 1986; White et al. 1988; Dreher and Jackson 1989). The results of the present

study indicate that activation of D₁ receptors is necessary for amphetamine-induced behaviors because SCH abolished amphetamine-induced hyperlocomotion in both control and lesioned groups. These results are consistent with those of the previous studies (Molly et al. 1986; Petry et al. 1993). In the present study, however, coadministration of quinpirole and SKF did not significantly increase locomotion in control rats. This lack of effect may be attributed to the doses of the two drugs administered and to the differences in experimental methodology between the present and previous studies (Dreher and Jackson 1989; White et al. 1988). Of importance was that coadministration of both quinpirole and SKF significantly increased locomotion only in lesioned rats compared to the treatment with saline or SKF alone. Although this effect was not significantly higher than that produced by quinpirole alone, the locomotor response following coadministration of the two drugs had a more rapid onset and a longer duration than that produced by quinpirole alone. This result provides additional evidence for the presence of neonatal hippocampal lesion-induced postsynaptic hypersensitivity of DA receptors.

In summary, the results of the present study extend previous findings indicating that there was no neonatal hippocampal lesion-induced effect on DA release in the NAc, although lesioned rats displayed locomotor hyperresponsiveness to amphetamine. The data from the present study indicate that behavioral hyperresponsiveness in lesioned rats is likely due to the enhanced postsynaptic sensitivity of D₂ receptors.

ACKNOWLEDGMENT

The authors thank Dona Cramer, Ellen Soly, Doug Rush, and Melinda Cilio for their excellent help in preparing the histological samples for this study and Susan Weiss, Ann Woods, and Chris Hough for their help in preparing this manuscript.

REFERENCES

- Amalric M, Koob GF, Creese I, Swerdlow NR (1986): "Selective" D₁ and D₂ receptor antagonists fail to differentially alter supersensitive locomotor behavior in the rat. *Life Sci* 39:1985–1993
- Barnes JM, Barnes NM, Costall B, Naylor RJ (1990): The actions of (-)-N-propylnorapomorphine and selective dopamine D₁ and D₂ receptor agonists to modify the release of [³H] dopamine from the rat nucleus accumbens. *Neuropharmacology* 29:327–336
- Barone P, Davis TA, Braun A, Chase TN (1986): Dopaminergic mechanisms and motor function: Characterization of D₁ and D₂ dopamine receptor interactions. *Eur J Pharmacol* 123:109–114
- Beckmann H, Cattaz WF, Jakob H (1987): Biochemical and

- neuropathological indices for the aetiology of schizophrenia. In Haffner H, Gattaz WF, Janzarik W (eds), *Search for the Causes of Schizophrenia*. Springer-Verlag, New York
- Bloom FE (1993): Advancing a neurodevelopmental origin for schizophrenia. *Arch Gen Psychiatry* 50:224–227
- Bogerts B (1993): Recent advances in the neuropathology of schizophrenia. *Schizophr Bull* 19:431–445
- Braun AR, Jaskiw GE, Vladar K, Sexton RH, Kolachana BS, Weinberger DR (1993): Effects of ibotenic acid lesions of the medial prefrontal cortex on dopamine agonist-related behaviors in the rat. *Pharmacol Biochem Behav* 46:51–60
- Brown R, Colter N, Corsellis JAN, Crow TJ, Frith CD, Jagoe R, Johnstone EC, March L (1986): Postmortem evidence of structural brain changes in schizophrenia: Differences in brain weight, temporal horn area, and parahippocampal gyrus compared with affective disorder. *Arch Gen Psychiatry* 43:36–42
- Clark D, White FJ (1987): D₁ dopamine receptor—The search for a function: A critical evaluation of the D₁/D₂ dopamine receptor classification and its functional implications. *Synapse* 1:347–388
- Dreher JK, Jackson DM (1989): Role of D₁ and D₂ dopamine receptors in mediating locomotor activity elicited from the nucleus accumbens of rats. *Brain Res* 487:267–277
- Eilam D, Szechtman H (1989): Biphasic effect of D₂ agonist quinpirole on locomotion and movements. *Eur J Pharmacol* 161:151–157
- Eilam D, Clements KVA, Szechtman H (1991): Differential effects of D₁ and D₂ dopamine agonists on stereotyped locomotion in rats. *Behav Brain Res* 45:117–124
- Eilam D, Talangbayan H, Canaran G, Szechtman H (1992): Dopaminergic control of locomotion, mouthing, snout contact, and grooming: Opposing roles of D₁ and D₂ receptors. *Psychopharmacology* 106:447–454
- Falkai P, Bogerts B (1986): Cell loss in the hippocampus of schizophrenics. *Eur Arch Psychiatry Neurol Sci* 236:154–161
- Flores G, Barbeau D, Quirion R, Srivastava LK (1996): Decreased binding of dopamine D₃ receptors in limbic subregions after neonatal bilateral lesion of rat hippocampus. *J Neurosci* 16:2020–2026
- Gehlert DR, Wamsley JK (1985): Dopamine receptors in the rat brain: Quantitative autoradiographic localization using [³H]sulpiride. *Neurochem Int* 7:717–723
- Gehlert DR, Gackenheim SL, Seeman P, Schaus J (1992): Autoradiographic localization of ³H-quinpirole binding to D₂ and D₃ receptors in rat brain. *Eur J Pharmacol* 211:189–194
- Gonzalez-Lima F, Hart WT, Rivera-Quinones C (1987): Metabolite involvement in the behavioral effects of bromocriptine in cats. *Eur J Pharmacol* 141:109–115
- Gray JA, Feldon J, Rawlins JNP, Hemsley DR, Smith AD (1991): The neuropsychology of schizophrenia. *Behav Brain Sci* 14:1–84
- Groenewegen HJ, Berebde HW, Meredith GE, Haber SN, Voorn P, Wolters JG, Lohman AHM (1991): Functional anatomy of the ventral, limbic system-innervated striatum. In Willner P, Scheel-Kruger J (eds), *The Mesolimbic Dopamine System: From Motivation to Action*. Wiley, Chichester, pp 19–59
- Hall LD, Herrod JJ, Carpenter TA, McKenna PJ (1994): Magnetic resonance imaging in schizophrenia: A review of clinical and methodological issues. In Ancill RJ, Holliday S, Hienbottam J (eds), *Schizophrenia: Exploring the Spectrum of Psychosis*. Wiley, Chichester, pp 115–135
- Imperato A, Tanda G, Frau R, DiChira G (1988): Pharmacological profile of dopamine receptor agonists as studied by brain dialysis in behaving rats. *J Pharmacol Exp Ther* 245:257–264
- Kerwin RW, Murray RM (1992): A developmental perspective on the pathology and neurochemistry of the temporal lobe in schizophrenia. *Schizophr Res* 7:1–12
- Knable MB, Murray AM, Lipska BK, Karoum F, Weinberger DR (1994): D₂/D₃ and D₄ receptor densities are not altered in rats with neonatal hippocampal damage. *Soc Neurosci Abstr* 513.3
- Kotrla KJ, Weinberger DR (1994): A neurodevelopmental perspective on normal brain functioning, plasticity and schizophrenia. In Ancill RJ, Holliday S, Hienbottam J (eds), *Schizophrenia: Exploring the Spectrum of Psychosis*. Wiley, Chichester, pp 67–93
- Lipska BK, Weinberger DR (1993): Delayed effects of neonatal hippocampal damage on haloperidol-induced catalepsy and apomorphine-induced stereotypic behaviors in the rat. *Dev Brain Res* 75:213–222
- Lipska BK, Jaskiw GE, Weinberger DR (1993): Postpubertal emergence of hyperresponsiveness to stress and to amphetamine after neonatal excitotoxic hippocampal damage: A potential animal model of schizophrenia. *Neuropsychopharmacology* 9:67–75
- Lipska BK, Chrapusta SJ, Egan MF, Weinberger DR (1995a): Neonatal excitotoxic ventral hippocampal damage alters dopamine response to mild repeated stress and to chronic haloperidol. *Synapse* 20:125–130
- Lipska BK, Swerdlow NR, Geyer MA, Jaskiw GE, Braff DL, Weinberger DR (1995b): Neonatal excitotoxic hippocampal damage in rats causes post-pubertal changes in prepulse inhibition of startle and its disruption by apomorphine. *Psychopharmacology* 122:35–43
- MacLennan AJ, Maier SF (1983): Coping and the stress-induced potentiation of stimulant stereotypy in the rat. *Science* 219:1091–1093
- McKinney WT, Moran EC (1981): Animal models of schizophrenia. *Am J Psychiatry* 138:478–483
- Mednick SA, Cannon TD, Barr CE, Lyon M (1991): *Fetal Neural Development and Adult Schizophrenia*. Cambridge, Cambridge University Press
- Mittleman G, LeDuc PA, Whishaw IQ (1993): The role of D₁ and D₂ receptors in the heightened locomotion induced by direct and indirect dopamine agonists in rats with hippocampal damage: An animal analogue of schizophrenia. *Behav Brain Res* 55:253–267
- Mogenson GJ, Wu M (1991): Effects of administration of dopamine D₂ agonist quinpirole on exploratory locomotion. *Brain Res* 551:216–220
- Molly AG, O'Boyle KM, Pugh MT, Waddington AL (1986): Locomotor behaviors in response to new selective D₁ and D₂ dopamine receptor agonists, and the influence of

- selective antagonists. *Pharmacol Biochem Behav* 25: 249–253
- Ojeda SR, Andrews WW, Advis JP, Smith White S (1980): Recent advances in the endocrinology of puberty. *Endocrine Rev* 1:228–257
- Paxinos G, Watson C (1986): *The rat brain in stereotaxic coordinates*, 2nd ed. New York, Academic Press
- Petry N, Furnidge L, Tong Z-Y, Martin C, Clark D (1993): Time sampling observation procedure for studying drug effects: Interaction between d-amphetamine and selective dopamine receptor antagonists in the rat. *Pharmacol Biochem Behav* 44:167–180
- Randolph C, Goldberg TE, Weinberger DR (1993): The neuropsychology of schizophrenia. In Heilman KM, Valenstein E (eds) *Clinical Neuropsychology*, 3rd ed. Oxford, New York, pp 499–522
- Robertson GS, Robertson HA (1986): Synergistic effects of D₁ and D₂ dopamine agonists on turning behavior in rats. *Brain Res* 384:387–390
- Schmajuk NA (1987): Animal models for schizophrenia: The hippocampally lesioned animal. *Schizophr Bull* 13:317–327
- Seeman P, Van Tol HHM (1994): Dopamine receptor pharmacology. *TIPS* 15:264–270
- Totterdell S, Smith AD (1989): Convergence of hippocampal and dopaminergic input onto identified neurons of the nucleus accumbens of the rat. *J Chem Neuroanat* 2:285–298
- Van Hartesveldt C, Cottrell GA, Potter T, Meyer ME (1992): Effects of intracerebral quinpirole on locomotion in rats. *Eur J Pharmacol* 214:27–32
- Wachtel H, Ahlenius S, Andén N-E (1979): Effects of locally applied dopamine to the nucleus accumbens on the motor activity of normal rats and following α -methyltyrosine reserpine. *Psychopharmacology* 63:203–206
- Waddington JL (1989): Functional interactions between D₁ and D₂ dopamine receptor systems: Their role in the regulation of psychomotor behavior, putative mechanisms, and clinical relevance. *J Psychopharmacol* 3:54–63
- Walters JR, Bergstrom DA, Carlson JH, Chase TN, Braun AR (1987): D₁ dopamine receptor activation required for postsynaptic expression of D₂ agonist effects. *Science* 236:719–722
- Wan RQ, Giovanni A, Kafka SH, Corbett R (1996): Neonatal hippocampal lesions induced hyperresponsiveness to amphetamine: Behavioral and microdialysis studies. *Behav Brain Res* 78:211–213
- Waters N, Svensson K, Haadsma-Svensson Smith MW, Carlsson (1993): The dopamine D₃-receptor: A postsynaptic receptor inhibitory on rat locomotor activity. *J Neural Transm* 94:11–19
- Waters N, Lofberg L, Haadsma-Svensson Svensson K, Sonesson C, Carlsson A (1994): Differential effects of dopamine D₂ and D₃ receptor antagonists in regard to dopamine release, in vivo receptor displacement and behavior. *J Neural Transm* 98:39–55
- Weinberger DR (1987): Implication of normal brain development for the pathogenesis of schizophrenia. *Arch Gen Psychiatry* 44:660–669
- Whishaw IQ, Mittleman G (1991): Hippocampal modulation of nucleus accumbens: Behavioral evidence from amphetamine-induced activity profiles. *Behav Neural Biol* 55: 289–306
- White FJ, Bednars LM, Wachtel SR, Hjorth S, Brooderson RJ (1988): Is stimulation of both D₁ and D₂ receptors necessary for the expression of dopamine-mediated behaviors? *Pharmacol Biochem Behav* 30:189–193
- Wilkinson LS, Mittleman G, Torres E, Humby T, Hall FS, Robbins TW (1993): Enhancement of amphetamine-induced locomotor activity and dopamine release in nucleus accumbens following excitotoxic lesions of the hippocampus. *Behav Brain Res* 55:143–150
- Yang CR, Mogenson GJ (1986): Dopamine enhances terminal excitability of hippocampal-accumbens neurons via D₂ receptor: Role of dopamine in presynaptic inhibition. *J Neurosci* 6:2470–2478
- Yue J-L, Nakamura S, Ueda H, Misu Y (1994): Endogenously released L-dopa itself tonically functions to potentiate postsynaptic D₂ receptor-mediated locomotor activity of conscious rats. *Neurosci Lett* 170:107–110