sterone only by the absence of the fourth oxygen, and can therefore also be called 'desox-corticosterone'. It shows striking resemblance with corticosterone. Its reducing capacity and the cholestenone-like ultra-violet absorption spectrum are in accordance with the function of the three oxygen atoms. But the chief point is that it has definite cortical hormone activity on adrenalectomized dogs and rats as shown by Prof. E. Laqueur in Amsterdam*. The minimal necessary daily dose is not yet exactly established, but it is certain that the activity is not less than one third of that of corticosterone. The crystals have 136°–138°, which possibly may be raised a few m.p. degrees when greater amounts are prepared. The important fact, however, is that this compound represents the first substance with high cortical hormone activity prepared from inactive material.

In a paper which arrived here a few days ago, Kendall *et al.*³ state that corticosterone is identical with their compound *B*⁴. They now give the m.p. of the latter as 177°–179° (uncorr.) which is in excellent agreement with our value for corticosterone. Also the biological activity is quite of the same order as that reported by us. Kendall states that this substance has a higher cortical hormone activity than that of any other pure compound so far isolated. The empirical composition is given as C₂₁H₃₀O₄, and the structure is expressed in formula (III). While Kendall is thus the first to have published an appropriate structure for corticosterone, it must be pointed out that in his original paper⁴, Kendall stated that compound *B* had a melting point 135°–139°, which is more than 40° lower than that now given, and an empirical composition of C₂₄H₃₄O₅. Nothing was said of its biological activity, while activity for compound *E* was claimed⁵. It is therefore scarcely possible that his original compound *B* is identical with corticosterone; it may have been a mixture.

On the other hand, we agree with Kendall, that his new results bring considerable evidence for his suggestion that the unknown fourth oxygen is contained in a secondary hydroxy group. If this suggestion can be proved conclusively, there will remain very few positions in which this hydroxy group can be placed. In the meantime, there exists evidence which would suggest its presence at position 11, rather than at position 12.

MARGUERITE STEIGER.
T. REICHSTEIN.

Chem. Institut,
Eidg. Tech. Hochschule,
Zürich.
April 6.

* The biological tests were made with the acetate.

¹ de Fremery, Laqueur, Reichstein, Spanhoff, Uylert, *NATURE*, **139**, 26 (1937).

² This formula was not published, but deposited in a Swiss patent filed December 1936.

³ Kendall, Mason, Hoehn, McKenzie, *Proc. Staff Meet. Mayo Clinic*, **12**, 136 (1937).

⁴ Mason, Myers, Kendall, *J. biol. Chem.*, **114**, 613 (1936).

⁵ Kendall, Mason, Myers, *Proc. Staff Meet. Mayo Clinic*, **11**, 361 (1936).

Variation of Wind with Height

IN the Professional Note of the Meteorological Office, Air Ministry, referred to in *NATURE* of April 24 in a Research Item (p. 721) on the "Variation of Wind with Height" at Wadi Halfa, Mr. J. Durward remarks upon an apparent discrepancy of nearly 70° between observation and theory when the results of observations of the winds by means of pilot balloons between 10,000 ft. and 13,000 ft. are compared with the run of the isobars at 4,000 metres as shown on Teisserenc de Bort's map.

I should like to remark that the discrepancy really emphasizes the necessity for further investigation of the meteorology of the equatorial region. Our general ideas of the relation of wind to the distribution of pressure, initiated by Buys Ballot's law, are based on the experience of temperate latitudes. As the equator is approached, the term in the equation depending on the earth's rotation gradually loses its significance and at the equator itself is altogether inoperative. In that region, if air 'comes to rest', it is by, somehow, acquiring a velocity of 1,000 miles an hour in order to keep pace with the ground on which it 'rests', a velocity which in temperate or arctic latitudes would be intolerable.

The equatorial region, therefore, cannot be regarded as amenable to the ideas of the relation of air-motion to the gradient of pressure which are illustrated by our weather-maps. Special investigation is needed to put the meteorologists of the equatorial zone in a corresponding position of advantage.

Special maps are certainly required; but what form the maps should take also requires consideration. Monthly means are not likely to be of much service where the geostrophic effect is negligible. Years ago, Mr. Lempfert and I got a good deal of insight into the working of that effect in the temperate zone by the construction of hourly maps compiled from the available records, and now that records of pressure are required for other purposes besides weather-maps, I have been wondering whether there are enough records in existence of pressure over land and sea to make hourly maps of the equatorial regions, if the records for a single week, for example, could be collected. The demonstration of the diurnal variation of pressure might be a test of the reality of the representation, and incidentally the local divergencies from the smooth curve would no doubt be expressed as weather. I have already an offer from Colombo of any information that is available there, and with the aid of ships and sealing-wax the walrus's idea of talking about many things might be realized!

NAPIER SHAW.

10 Moreton Gardens, S.W.5.
April 28

Aldehyde Mutase

WITH reference to the note by M. Dixon and C. Lutwak-Mann, published in *NATURE* of March 27, p. 548, on the non-identity of aldehyde mutase with the aldehyde oxidase, I would point out that this fact was first established so far back as 1931 by Michlin and Severin and published in *Biochemische*