between reactions of aryl radicals in the vapour phase and the corresponding reactions in solution, since in the latter case the free radical invariably reacted with the solvent molecules. The characteristics of the phenyl radical, formed in the presence of a solvent, may be summarized as follows: (a) the free radical is extremely reactive and able to interact with almost the first molecule of any type encountered; (b) polymerization to diphenyl does not take place; (c) reaction with all non-aromatic hydrogen-containing solvents results in the formation of benzene ($C_6H_5\cdot + RH \rightarrow C_6H_6$); (d) reaction with all non-aromatic halogen-containing solvents results in the formation of a halogeno-benzene ($C_6H_5\cdot + RHal \rightarrow C_6H_5Hal$); and (e) reaction with neutral aromatic liquids results in the formation of diphenyl derivatives in which the radical enters the aromatic nucleus at the *ortho* and/or *para* position with respect to any directing group which may be present.

A close similarity is thus revealed between the properties of alkyl radicals and those of aryl radicals in solution, and this correspondence gives striking support to the views originally put forward with reference to a non-ionic mechanism for certain reactions of the diazo hydroxides, the nitrosoacetylaminos, the arylazotriarylmethanes and the diacyl peroxides. In addition, various lines upon which the correspondence may be further extended immediately suggest themselves. The whole subject of reactions involving free radicals in solution will shortly be reviewed elsewhere.

University,
Manchester.

University,
Durham.
Oct. 22.

D. H. HEY.

W. A. WATERS.

¹ Norrish and Bamford, *NATURE*, **140**, 195 (1937).

² Walker and Wild, *J. Chem. Soc.*, 1132 (1937).

³ Grieve and Hey, *J. Chem. Soc.*, 1797 (1934).

⁴ Hey, *J. Chem. Soc.*, 1966 (1934).

⁵ Waters, *J. Chem. Soc.*, 113 (1937).

Behaviour of Cylinders of Inflammable Gas in a Fire

IN a letter to *NATURE*¹, my friend, Prof. K. C. Bailey, refers to the evidence I gave before the Free State Government's inquiry into the Pearse Street fire of October 5, 1936, where, owing to the failure of the water supply, three firemen lost their lives through carbon monoxide poisoning. Prof. Bailey's success in extinguishing a flame with a burst of gas from a high-pressure container will scarcely surprise anyone who has tried to light a gas jet (at only a few inches water-pressure) by holding a kindled match directly in its path. A draught of air, also, will extinguish a candle; though it might perhaps be expected to contribute to the combustion. A draught does not extinguish a blast-furnace.

Prof. Bailey's theory accounts for the two explosions that were heard by so many witnesses as being due (1) to the bursting of the coal-gas cylinder, and (2) to the chemical explosion of the atmosphere then constituted. Actually the evidence is perfectly clear that the coal-gas cylinder yielded plastically, bulging and dividing into a "pair of trousers". All the witnesses were shown this cylinder at the time of the inquiry.

Now the companion oxygen cylinder burst in such a manner as to blow part of its wall away; yet this rupture, though it must have been percussive in the highest degree, is apparently held to have been inaudible, and the discharge of 120 cubic feet of oxygen into a highly super-fuelled atmosphere is not, apparently, held to have been a similar cause of explosion.

There is no doubt, surely, that whatever cooling took place in the body of the liberated coal-gas, its periphery must quickly have attained ignition temperature in the burning inferno which Prof. Bailey postulates. Then it must have moved as a flame-mantled blimp towards and through the great ventage whence a glass roof had fallen in at some time before the two explosions.

22 Tree Root Walk,
Sheffield, 10.
Oct. 29.

OLIVER C. DE C. ELLIS.

¹ *NATURE*, **140**, 503 (Sept. 18, 1937).

Points from Foregoing Letters

THE synthesis of a complex carbohydrate from carbon dioxide and water in the presence of light, by means of a specially prepared nickel catalyst, is reported by Prof. E. C.-C. Baly. The compound, when deposited on a surface, can be hydrolysed by diastase, and hence it may prove to be the parent of a starch-like substance.

Although the spectrum of nitrogen excited by electron bombardment varies greatly with pressure, R. Bernard finds that the spectral composition of auroras having a height of several hundred kilometres remains nearly constant. He considers, therefore, that the energy of the electrons and not pressure is the factor affecting the relative intensity of the radiations emitted.

M. G. E. Cosyns suggests that the anomalous distribution of cosmic rays coming from different directions (zenithal angles) may be explained by taking into account the diffusion of cosmic rays in air.

Dr. H. Grüneberg gives a list of pairs of mutations of the fruit fly, *Drosophila*, which have very nearly related characters, such as chlorotic-yellow, and points

out that the corresponding genes occur as doublets in close proximity within the same chromosome. He suggests that new genes arise by small adjacent duplications followed by differentiations.

Dr. C. D. Darlington states that the genetic action of chromosomes in the resting nucleus on the cell may be inferred from the rapid death of pollen grains and embryo sacs with defective chromosome complements in hybrid plants.

Further to the experiments on the sensitization by certain cancer-producing substances of mice by light, Dr. J. C. Mottram and Dr. I. Doniach find that paramecium is destroyed by benz-pyrene suspensions (1 in 10,000) in the presence of light of wave-length between 3500 Å. and 4100 Å., but not in the dark.

Dr. D. H. Hey and Dr. W. A. Waters point out that the reactions of free alkyl radicals produced photochemically in solution are exactly similar to those of free aryl radicals which they consider to be formed when certain classes of complex organic molecules decompose.