

of the results of recent excavations lies in the fact that they have brought to light remains belonging to the pre-Roman period, as well as evidence pointing to the measure of fusion with native culture.

At Saint-Rémy, which possesses two theatres of the Roman period, both Greek and Roman constructions beside a tile-covered canal have been found near a Roman road, triumphal arch and tomb dating from the first century, and beneath a Roman wall. Fifty-two altars of local stone and marble were dedicated to either the god Sylvanus or the goddess Cybele. The foundations of an entirely Greek house were uncovered, in which twelve rooms are seen to have opened into an *atrium*, in which the bases of four columns are still visible. Baths of the time of Augustus, measuring forty-seven yards by thirty-five yards, were found. Into these, water flowed through a huge sculptured head. A large amount of Greek red-on-black pottery was found with forty-four small votive lamps and a magnificent rock crystal ring, ornamented with a woman's head.

Further discoveries in these excavations are now announced. According to a dispatch from the Paris correspondent of *The Times* in the issue of May 23, while most of the walls which have been found belong to a large Roman construction, the architectural remains of pre-Roman date are also important. A

pillar with egg and astragalus pattern is of second-century Greek style, while column heads, probably pointing to Italian models, are evidently the work of competent masons of Greek training. The gargoyles, however, are the rustic Celtic versions of classical gargoyle lions. It is thought that they belong to a building, possibly a temple, of Greco-Celtic style.

Further evidence of cultural fusion comes from Glanom, a small Celtic *oppidum* near Saint-Rémy, where in a large necropolis graves containing Celtic and local Hellenistic pottery have been excavated by M. de Brun, keeper of the local museum. The tombs, it is stated, are marked by tall pillars, on which Celtic names are inscribed in Greek character. The cemetery is dated in the period of the Roman occupation at the end of the second century B.C. Still earlier evidence comes from Roquepertuse, where the Count de Gerin-Ricard, keeper of the Musée Borely at Marseilles, has excavated a sanctuary of the fourth century B.C. Here he has found what is described as a curious mixture of native and Greek taste. The pillars supporting the entablature are decorated with the heads of slain enemies, the horses on the frieze are of the barbarian type, while the double heads affixed to the architrave are a successful and attractive blend of Celtic and Greek elements.

## Antarctic Plankton\*

EXACT systematic work is essential for oceanographical plankton studies, since we must know the species with which we are dealing before we can use them in any practical way. Mr. Hendey's extensive monograph on the diatoms of the Southern Ocean is most useful and is a careful study entailing a large amount of labour. With so much material it has been possible to review the inside variation present in most groups and in the individual species. The interesting fact is clearly brought out, agreeing with Wimpenny (1936) that the diatom frustules in cold water are smaller, thicker-walled and more vigorous than those from warmer waters, the reverse of what is known to exist in many marine animals. The holoplanktonic flora of the cold water from the oceanic habitats forms the major part of the collection and consists of a small number of discoid and a large number of filamentous pennate forms. The author, however, in his classification discards the usual division into Centricæ and Pennatæ, and his method of regarding many so-called species and varieties as "phases" of one species reduces considerably the number of species recognized. Thus, to take one example, *Corethron* according to Hendey is a monotypic genus, *Corethron criophilium*, much discussed by many workers, being the only species with many phases. This at least tends to simplify matters, and a reduction of the ever-increasing number of species is all to the good.

Most of the diatoms of the Southern Ocean have

a wide distribution and can tolerate a great variation in the conditions of their environment, but it is found that *Rhizosolenia curvata* Zacharias has a very limited range, being a typical sub-antarctic species occasionally spreading into Antarctic waters. Dr. Hart in his paper on this diatom shows that it forms a valuable indicator of the southern limit of sub-antarctic surface water. It has always been recognized that this surface water is frequently forced some small distance south of its normal limit, the Antarctic Convergence, and it has been possible to prove that *R. curvata*, an easily recognizable and characteristic form, is always in the plankton whenever it has been possible to show such southward movement hydrographically.

The seasonal circulation of the Antarctic macroplankton is dealt with by Dr. Mackintosh. Amongst other interesting facts it is shown that certain species which are usually regarded as inhabitants of the surface layers make an annual vertical migration into deep water in winter, and this migration by which they move from one current system to another is probably the means by which they keep within their normal geographical boundaries. The three species which in places make up the bulk of the Antarctic macroplankton, *Rhincalanus gigas*, *Eukrohnia hamata* and *Calanus acutus*, are mainly concentrated in the surface water in summer but descend into very deep water in winter. Since part of the Antarctic surface water moves in a northerly direction and there is a compensating southerly movement of the warm deep water, this vertical migration almost certainly results in the return of the species to the regions from which it originally came. The circulation is on a very large scale, the vertical range being from four to six hundred metres and the horizontal range being presumably some hundreds of miles.

\* Discovery Reports. Issued by the Discovery Committee, Colonial Office, London, on behalf of the Government of the Dependencies of the Falkland Islands. Vol. 16. Pp. 151-364+plates 6-13. The Plankton Diatoms of the Southern Seas. By N. Ingram Hendey. 33s. net. Vol. 16. Pp. 365-412. The Seasonal Circulation of the Antarctic Macroplankton. By N. A. Mackintosh. 6s. net. Vol. 16. Pp. 413-446+plate 14. *Rhizosolenia curvata* Zacharias. By Dr. T. John Hart. 5s. net. (Cambridge: University Press, 1937.)