effects of the depression. If the younger herring have forsaken their spawning grounds and gone elsewhere, we may hope that they will at once return in force when conditions improve, and that the Plymouth fishery will rapidly be re-established. If, however, they have throughout held to their

former migration routes, and the present dearth is due to lack of suitable conditions for the larvæ, they are in the same position as the bottom fish, and a number of years must elapse before the fishery can be resumed.

(To be continued.)

Luminescence

AN important series of discussions arranged by the Faraday Society took place in the Biochemical Theatre of the University of Oxford on September 15-17 on the subject of "Luminescence". A more appropriate time for the discussion of work in this field could scarcely have been chosen, and in spite of recent events, a surprisingly large number of visitors from overseas was present. The discussions were formally divided into three parts: luminescence of vapours, liquids, and solutions; luminescence of solids; chemiluminescence. The subject of luminescence is of particular interest at the present time for two reasons. First, it forms an excellent testing ground for some of the more recent theories, particularly with respect to the physical properties of solids, and the nature of chemical reactions. Secondly, a good deal of stimulation has been given by the technical interest in the subject. The use of fluorescent solids in discharge tubes as a means of modifying the colour of the light or increasing the efficiency of such devices is undoubtedly leading up to new work of interest, and contributions to the meeting were made by more than half a dozen industrial laboratories.

The first two parts of the discussion dealt essentially with the absorption of radiation by matter and its emission in the form of light; the final section was concerned, however, with the radiations emitted as a result of chemical reactions. The now classical experiments of Polanyi and his collaborators on the characteristics of the reaction between streams of halogen and sodium vapour at low pressure are a specially interesting example of chemiluminescence, in that they are providing the basis for theoretical work in this field.

A valuable feature of the earlier parts of the meeting was the way in which ideas concerning the precise meaning of the terms 'fluorescence' and 'phosphorescence' were brought out. If a luminescent compound is first illuminated at *O* (see accompanying graph) with radiation of constant intensity, an appreciable time may elapse before the luminescence reaches a constant value, as along

AB. If the exciting radiation (for example, ultra-violet) is removed at a time corresponding to *B*, a rapidly falling curve of intensity *BC* follows. Many, and in fact most, of those engaged in experimental work in this field have usually referred to the part *AB* as 'fluorescence' and the part *BC* as 'phosphorescence'. This usage is undoubtedly convenient from the experimenter's point of view, since the time constant of the luminescence has usually been measured during the period *BC*. It has of course been realized that the electrons returning to the ground-state during the equilibrium period *AB* do so only after a finite and possibly considerable time after the absorption process. It was suggested in the course of the discussion that definitions based on the type of electron transition would be more fundamental than those already described. On this basis the term 'fluorescence'

should be reserved for all processes in which electrons are returning from *excited* energy states of parent atoms, and the term 'phosphorescence' used only in those cases where the luminescence is dependent on temperature, and is a result of electrons being involved in metastable states or completely removed from the parent atoms. It is likely that the older definitions will continue to be used on account of their convenience, and because of insufficient data concerning processes. Further progress along these lines can only be made by a study of the time-dependent effects of luminescence, and the corresponding photo-conductivity properties.

Very many substances, both liquid and solid, are capable of absorbing some part of the ultra-violet spectrum, but comparatively few are luminescent. It is to be assumed that the

non-luminescent substances convert this absorbed energy into heat in the form of elastic vibrations of much smaller quanta than visible light, and the important theory of Peierls developed some years ago on the problem of radiationless transitions in solids is now receiving much attention. This problem of radiationless transitions is of obvious importance to all branches of the subject. In liquids and solutions, for example, the fluorescence may be 'quenched' by increasing concentration of the luminescent centres, or by the addition of oxygen. The problems associated with the quenching of luminescence by a close coupling between the excited molecules and the surrounding solvent on one hand, and the question of internal quenching on the other, were made clearer by pictorial representation in intersecting curves of potential energy as a function of atomic or molecular separation. The work on liquids is concerned chiefly with organic solutions, and the complexity of the problem involved is shown by the fact that some organic solids show a large drop in the quantum efficiency of fluorescence when put into solution, while others show an equally sharp rise.

Almost a whole day was given up to the discussion of luminescence in solids, and on the whole this was the most successful feature of the meeting. It is now clearly recognized that there are two chief classes of luminescent solids. The more restricted class contains pure solids, and it is sometimes possible to recognize these by the fluorescence of their solutions; notable examples of this kind are the uranyl salts, the platino-cyanides, and certain rare earth compounds. The fluorescence in the first two examples appears to be associated with co-ordination groups; in the rare earths the sharp line spectra are connected with transitions within the $4f$ shell, but it has not been explained why only certain of the rare earths are fluorescent.

Most luminescent solids, however, owe their properties to the deliberate addition of some impurity. But between the two well-defined groups of pure and deliberately impure solids are a number of others, the cause of luminescence in which is still in some doubt. Zinc oxide, for example, fluoresces in the near ultra-violet with an easily recognizable spectrum, and while it is assumed that the luminescence is linked up with the 'self' impurity of excess zinc, no direct proof of this has yet been obtained. The presence of *foreign* metal atoms in a solid may be made a matter of direct spectroscopic or chemical test, but other means will have to be adopted in cases such as that of zinc oxide. Calcium tungstate is also fluorescent when free from other metallic impurities, and it has been suggested that the WO_4 co-ordination group is of fundamental importance to the

luminescence; it may well be, however, that the excess or defect of one of the lattice constituents has a bearing on this problem also. 'Pure' zinc sulphide presents a problem of even greater difficulty. Zinc sulphide has been prepared by several workers during the last few years in a state apparently free from foreign metallic impurity, and the fluorescence is a bright blue. The precipitation of zinc sulphide free from oxide is, however, a matter of great difficulty, if not impossible, and experiments have yet to be carried out to prove whether the luminescence is a property of the pure ZnS lattice or of a lattice containing (a) excess zinc or (b) some oxygen replacing sulphur atoms. These points have been mentioned merely to indicate that the problems of interpreting the luminescence of solids are by no means confined to the theoretical side.

The discussions at Oxford showed also the importance of low temperatures in the investigation of luminescence in solids. The spectra are, apart from a few exceptions, generally of a continuous nature at room temperature. The use of temperatures in the neighbourhood of 90° – 20° K. has shown that many new features may be brought out, either in resolution, or in the comparison of various substances with a common activator. In quite a number of solids, luminescence is apparent only at these low temperatures. The difficulties of measuring the absorption spectra of solids in powder form are well known, but the discussion brought to light a new technique that will doubtless be of value in this field. Single crystals will, when obtainable, be of value, but the distinction between surface and volume luminescence was emphasized as a point of importance.

The theoretical aspects of this part of the discussion were of value in making clear the advance in ideas from the rather restricted zone theory as applied to semi-conductors a few years ago. On this theory the sharp energy levels of an isolated atom were converted in the solid to a series of possible energy bands, separated by 'forbidden' regions. In a pure insulating solid, it might be possible to raise an electron from the highest full band to the next possible empty band. When this happened, the electron would be free to move through the crystal under an applied field; such a case would correspond to the ionization of the parent atom, and the recombination of the electron with its positive 'hole' would give rise to the emission of radiation. The advance in theory is concerned with a possible series of excitation levels lying between the highest full band and the lowest conduction band. The 'forbidden' zone on these ideas no longer exists in simple form, and it is possible, as might be expected, to excite electrons in solids without making them free for photo-conduction. An important feature of the newer

theoretical ideas is that it should be possible to distinguish between excited and ionized atoms in the solid by tests on photo-conduction and on phosphorescence. In the simplest cases the phosphors with a hyperbolic decay law should be photo-conducting, while those with exponential decay laws should not. The well-known high efficiency of alpha-particle excitation of luminescence in zinc sulphide phosphors, it was pointed out, is not due to the absorption of energy by the whole lattice and its subsequent transfer to the activating impurity atoms, but to the production of numbers of fairly high-energy electrons (δ -rays) colliding directly with the impurities.

The last part of the meeting was occupied with chemiluminescence, which is usually concerned with the production of light during chemical reactions; more recently, it has been shown that some reactions produce feeble radiations in the far ultra-violet. Bioluminescence is now generally regarded as a chemiluminescence in which the luciferin is oxidized in the presence of molecular oxygen and the enzyme luciferase. The work of Polanyi and Evans on sodium and chlorine vapour reactions is typical of one type of chemiluminescence and indicates that the production of light is a consequence of the following steps:

where X is Cl, Br or I, rather than one involving a neutral halide molecule and an excited sodium atom. Bawn has recently extended this type of work to reactions involving organic halides such as CH_2Cl_2 , $\text{CH}_3\text{CCl}_2\text{CH}_3$, etc.

The other type of chemiluminescence to be discussed was the oxidation process appearing in the luciferin reaction already mentioned, the oxidation of the Grignard reagents, the dimethyldiacridylum salts, and many others. Two methods of approach were apparent at the discussions. Weiss adopted the more physical method, and discussed the cases where the chemiluminescence can be regarded as the reverse of photochemical dissociation ($A + B \rightleftharpoons AB + h\nu$) and the quantum conditions associated with this type of reaction. Drew, on the other hand, has attacked the problem of chemiluminescence in the phthalic hydrazides from the purely chemical point of view of investigating the reactions that lead to their formation, the effects of the nature and position of substituents, and the interaction of the hydrazides with hydrogen peroxide.

The discussions as a whole were of value in making clear the precise nature of some of the important problems to be solved, and both the official meetings, and the more informal exchanges of views that inevitably take place on occasions of this kind, played their part.

J. T. RANDALL.

The State and Medical Research in Great Britain*

THE annual Harveian Oration was delivered by Sir Edward Mellanby, secretary of the Medical Research Council, on October 18 at the Royal College of Physicians, London, the subject of his discourse being "The State and Medical Research".

Sir Edward first paid a tribute to Harvey, who "stands out as one of the greatest investigators, and his work—a wonderful combination of observation and experiment—is an example of the finest achievement of the human mind". Harvey's election in 1630 as physician-in-ordinary to Charles I brought him into close association with the head of the State, and one result of this association was permission to make use of deer and other animals in the royal demesnes for study—probably the first instance, with certain legendary exceptions, of State support for medical research. Harvey's example has induced more and more individuals to take up research, and the revolu-

tionary changes in medicine during the past sixty years only increase the appreciation of his foresight and skill. These revolutionary changes include the introduction of new methods for the diagnosis, treatment and prevention of disease; the enormous increase in hospital accommodation; the formation and development of the clinical laboratory and radiological department; and the expansion and multiplication of the public health services.

Increase of knowledge has been mainly responsible for these developments, and it has come to be realized that health and disease are not conditions conferred upon man for his humble acceptance, but that by a certain method (Harvey's method) disease may be controlled and death itself warded off, and more and more men and women proceed to investigate and to accumulate knowledge.

The acceptance in Great Britain of the new urge for research has been comparatively recent. The late Earl Balfour said in 1896, "I have all my life

* Substance of the Harveian Oration before the Royal College of Physicians, delivered by Sir Edward Mellanby, K.C.B., F.R.S., on October 18.