probably the most important component of the respiratory system in insects and other organisms¹², is similarly influenced, and it is hoped to investigate this aspect by the use of spectroscopic techniques.

When we come to examine more complex insecticidal systems, it is clear that similarity in biological response may merely indicate a coincidence in the dynamic chain of metabolic changes initiated at different links in the chain. For example, pyrethrins I and II are complex unsaturated ketonic esters which differ completely in molecular structure from the relatively simple inert fat-soluble D.D.T.¹. Yet similarity in insecticidal action is shown by a dispersant action on the cuticle and internal tissue lipoids, and the lethal stages of poisoning reveal an increase in the phenoloxidase activity of tissue and haemolymph. The minute nature of the lethal dosages is consistent with an adsorption process on the tissue receptors; but it is unlikely that narcosis or death is directly related to primary insecticide-tissue combination. A contributory factor may be the accumulation of reactive *o*-quinones which would block essential substrate access to the whole complex of enzyme systems normally associated with the tissue components. This narcotic stage would be followed by death when the further dispersant action of the insecticides resulted in the accumulation of lethal concentrations of toxic quinonoid metabolites in the tissues. Richter¹³ has shown the general similarity between the catechol oxidases in plants, fungi and insects, and has found that oxygen uptake is greatly reduced by the *o*-quinone reaction products, which combine with the available catechol oxidase. However, these more specific factors in insecticidal action are beyond the scope of the present article and will be considered elsewhere.

This account is based on an investigation into the fundamental mode of action of insecticides and drugs which is being carried out by me at the Department of Colloid Science, Cambridge. I wish to express my deep indebtedness to Prof. E. K. Rideal and to Dr. J. H. Schulman for much helpful advice and stimulating discussion, especially on the physico-chemical aspects of the work.

¹ Luger, P., Martin, H., and Muller, P., *Helv. Chim. Acta*, **27**, 892 (1944).

² Hurst, H., *Nature*, **145**, 462 (1940).

³ Hurst, H., *Trans. Farad. Soc.*, **39**, 390 (1943).

⁴ Pryor, M. G. M., *Proc. Roy. Soc., B*, **128**, 378 (1940).

⁵ Wigglesworth, V. B., *Bull. Ent. Res.*, **33**, 205 (1942).

⁶ Fraenkel, G., and Rudall, K. M., *Proc. Roy. Soc., B*, **129**, 1 (1940).

⁷ Chargaff, E., and Ziff, J., *J. Biol. Chem.*, **131**, 25 (1939).

⁸ Dennell, R., *Nature*, **154**, 57 (1944).

⁹ Keilin, D., and Mann, T., *Proc. Roy. Soc., B*, **122**, 119 (1937).

¹⁰ Wigglesworth, V. B., *Nature*, **153**, 493 (1944); **154**, 333 (1944).

¹¹ Raper, H. S., *Biochem. J.*, **20**, 735 (1926); **21**, 89 (1927).

¹² Keilin, D., *Ergeb. Enzymforsch.*, **2**, 239 (1933).

¹³ Richter, D., *Biochem. J.*, **28**, 901 (1934).

organization and training of many generations of science students at Girton and Newnham Colleges.

In the early days of the higher education of women at Cambridge, the University lectures were not open to women and the teaching of science was left entirely to the women's colleges to provide as best they might. Friends of the women helped when they could, but there was no help from the University as such. Very soon after Miss Saunders had taken her Tripos she was appointed demonstrator in biology at Newnham College and taught the students of both Girton and Newnham Colleges; later she became a lecturer, and together with the late Mrs. Bidder (Marion Greenwood) helped to organize all the practical work in biology and botany given in what came to be known as the Balfour Laboratory. This building, a former Nonconformist chapel given to Newnham College by the generosity of Mrs. Sidgwick and Miss Balfour in memory of their brother Francis Maitland Balfour, the zoologist, was converted into the Balfour Laboratory under the direction of Miss Greenwood and Miss Saunders, and here Miss Saunders taught for many years, becoming its director after Miss Greenwood's marriage to Dr. Bidder.

She always set before her pupils a high standard, and much excellent work was carried out by them there. Even after many University professors and lecturers opened their lectures to women, all the practical work for Part I of the Tripos was still organized and carried out by her, until the new University Laboratories were built and accommodation provided in them for the women students.

There must be many women employed in scientific work in universities, colleges, schools and other places all over the world who know that they owe their present position and success largely to the stimulus of Miss Saunders' teaching and the high quality of the work on which she insisted.

G. L. ELLES.

ETHEL SHAKESPEAR.

THE main writings of Miss E. R. Saunders relate to floral morphology. The most important of them were issued between 1923 and 1936 in the *Annals of Botany*, the *New Phytologist*, the *Journal and the Proceedings of the Linnean Society*, the *Journal of Botany* and the *Journal of Genetics*, and in two volumes published by Messrs. Heffer and Sons. The earliest of them furnished a factual basis for a new and interesting outlook on gynoecial morphology; the later documents elaborated a theory of carpel polymorphism and brought to focus important problems of systematic and evolutionary interest which are still unsolved.

The first of these writings appeared at a time when some rendering of the classical conception of a monomorphic carpellary 'leaf' gave contentment to many inquirers. They looked on such an organ as the basic unit of gynoecial construction in all groups of flowering plants and thought of it as possessed of a median style and a distal stigma. Whatever was their mode of approach to floral matters, their common purpose was to settle long-standing differences of opinion regarding the 'basic numbers', arrangements and forms of such 'leaves' for all types of 'syncarpous' gynoecia and to furnish for each alliance a 'fundamental ground-plan of floral construction'.

Few indeed were prepared to describe precisely the 'foliar' prototypes of which they spoke; but many

OBITUARIES

Miss E. R. Saunders

WHEN Edith Rebecca Saunders died on June 6 at the age of eighty as the result of a bicycling accident in Cambridge, there passed one to whom many scientific women owe a great debt for her labours on their behalf in the early days of the higher education of women. Others have written of the value of her botanical research work, but not the least valuable part of her services was the

acted on the view that such common 'free' carpels as follicles and achenes exemplified their units and demonstrated some of the ways in which still older organs had been transformed, reduced in numbers, aborted or suppressed, and gradually rearranged on some uniform specific plan. Despite their union in pistils, they still showed, per species, a single carpel-form. Nevertheless, although there was constant reference to facts of development and histology and to all manner of final forms of pistil, opinion on gynoecial construction had come no nearer to harmony than it had been from the days of Braun and Eichler to those of Kerner and Klein.

The bearing of Miss Saunders' views may be made plain through a few points of past interest. Thus, while some believed that facts of development were of great value in the interpretation of all organs, others considered that adult form and structure gave the surest guide. There was also a mode of thought which gave prominence to vascular tissue not only as a major index to the status of an organ but also as a sign of its modification, or as a pointer to its prior development in a given position. There was a common language explaining the arrangement of vascular strands, their modes of branching, the remarkable conservatism of a vascular system, and the usefulness of 'vestigial' strands in tracing the old sites of arrested organs. Further, some believed that a leaf may be truly terminal since, as they said, a legume is a unit carpellary leaf and is fully distal. Others rejected this view and regarded a legume as the sole and displaced survival of a lateral group of 'leaves'. They read all 'syncarpous' gynoecia according to their notions of numbers and arrangements of lateral leaves, of those which had been suppressed, and of those which were now displaced.

Miss Saunders entered this field convinced that there could be no agreement on a 'fundamental ground-plan' for any pistillate family so long as the concept of a monomorphic carpel was the starting point in phyletic discussion. She pointed, in particular, to the unsettled controversies over pistil construction in the Cruciferae and other affinities, to the seemingly insoluble problem of the 'commissural' stigma, to the varying interpretations of obdiplostemony, to the wide range of ovule placentation still unexplained, and to the difficulties facing all monomorphists when they turned to families with 'supernumerary' styles. For these and all related matters she claimed the need of a new approach.

Her first proposal was as startling as it was simple, in that it visualized carpel dimorphism as shown to-day by many flowering groups. The heart of her view was that even a single pistil comprises united leaves; some with the form and the function of ovule-bearing organs, others of distinctive form and the function of which is stigmatic. Through this she sought early to rationalize cruciferous and other gynoecia and to explain the presence of commissural stigmas. As did others who opposed her plan for an extension of a 'leafy' view of carpels, she relied largely on the form and vasculature of mature organs. She came to identify and arrange her united carpel forms according to her estimate of branched or unbranched vascular strands, the positions of ovules upon ovarian walls, the arrangement of stigmas, and the older conception of the cyclic construction of a flower. It was inevitable that she should reject the conception of a terminal leaf once she had decided to interpret a distal legume as a union-product of distinctive carpel forms. Also it was natural that, with strands as a

major instrument in the identification of unit organs, she extended her dimorphic view into a doctrine of carpel polymorphism. One cannot attempt to outline the many directions in which she sought later to apply her theory and to settle once and for all the fundamental ground-plans of floral construction on which she had set her heart. But her final volumes are in themselves a glowing testimony to the breadth of her inquiries, to her courage, and to the great array of facts which she disclosed.

It may be long before there is common agreement as to the chief objective of floral study and the valuation of the evidence on which Miss Saunders relied. But it may truly be said that few of her contemporaries in formal morphology stirred imagination more deeply than she, or did so much to stimulate inquiry.

J. McL. THOMPSON.

Mr. H. E. Potts

THE very narrow but nevertheless important field of patent law and practice has lost a great mind by the death of Mr. H. E. Potts on July 4. He was an earnest and original contributor to the literature of patent law. Early in his career as a chartered patent agent, he was struck by the fact that, in the drafting of patent specifications and claims, a patent agent had to call upon powers of analysis and then subsequent synthesis, and that this logical process should have some analogy in mathematics. His first contribution on "An Application of Mathematics to Law" was indeed to *Nature* of April 24, 1913 (p. 187), and this has been followed over the years by a development of the idea there lightly touched upon.

Although possibly never hitherto positively appreciated, it is nevertheless a fact that a patent granted upon an invention when tested by the High Courts of Great Britain is either 'wholly bad' or 'wholly good'. It is believed that Mr. Potts was the first to point out that there should be some method of assessing validity quantitatively, as well as qualitatively; and recently he put forward a number of suggestions in regard to the appreciation of monopolies of invention as granted by the Crown on a graduated basis, as distinct from the 'all good' or 'all bad', developing this proposition side by side with its mathematical exposition and following accepted theories of modern philosophy.

Of many valuable publications, perhaps Potts' best known is his book "Patents and Chemical Research", which has become a standard work in the literature on patents.

WE regret to announce the following deaths:

Mr. A. D. E. Elmer, the distinguished plant collector and author of many papers on the botany of the Philippines, in July 1942.

Dr. Frank Blair Hanson, associate director for the natural sciences at the Rockefeller Foundation, formerly professor of zoology at Washington University, St. Louis, on July 21, aged fifty-nine.

Dr. Gustave M. Meyer, associate in biochemistry during 1913-41 at the Rockefeller Institute for Medical Research, known for his work on carbohydrates and tissue chemistry, on May 9, aged sixty-nine years.

The Ven. Lonsdale Ragg, founder and editor of the *Tree Lover*, and known for his drawings of trees, on July 31, aged seventy-eight.