THE CHEMICAL SOCIETY

By DR. F. SHERWOOD TAYLOR

DURING July 15-17 of this year the Chemical Society is celebrating its centenary, which but for the War would have taken place in 1941.

The early scientific societies extended their activities to all branches of science; it was only at the beginning of the nineteenth century that societies for the study of special branches of science began to be formed. The Astronomical, Linnean and Geological Societies belong to this period. The Chemical Society was the first national body to take chemistry as its sole object of study, for it was not until 1857 that the French Chemical Society was formed, while the Berlin Chemical Society did not originate until 1868.

In the early nineteenth century there was a considerable body of Englishmen eminent in chemistry—the names of Dalton, Davy, Wollaston, Faraday come to mind. In the eighteen-forties, however, the main activity in the science was on the Continent, where Liebig, Wöhler, Bunsen, Dumas, Gerhardt and Berzelius were at the height of their powers. The British chemists of the time were well aware of the importance of the subject, both for pure science and for industry, and the first attempts to form a society met with a warm response. The moving spirit was Robert Warington, who on February 23, 1841, convened a meeting “for the purpose of taking into consideration the formation of a Chemical Society”, which was duly constituted on March 30. Thomas Graham became president; Brande, Daniell, Clark, Daubeny and Grove were among the officers. The purposes of the Society were stated in the following words: “for the advancement of Chemistry and those branches of science immediately connected with it, for the communication and discussion of discoveries and observations relating to such subjects: the formation of a Library of scientific works and of a Museum of Chemical preparations and Standard Instruments are also ulterior objects of the Society”. All these intentions, excepting the last, towards which only a beginning was made, have been carried out in a manner that its founders can scarcely have anticipated.

The Society began work at once and published from time to time *Memoirs* and *Reports of the Proceedings*. Membership increased, and in 1849 a Royal Charter was obtained.

The first meetings were held in the premises of the Society of Arts; in the same year they were removed to the Westminster Literary and Scientific Institution at 47 Leicester Square; but in 1843 returned to the Society of Arts. In 1850 the Society moved to 142 Strand, then to Cavendish Square, where it occupied a part of the premises of the Polytechnic Institution. About 1853 the Government promoted a plan for accommodating the scientific societies at Kensington Gore, but this found little favour. The Burlington House site was proposed instead, but it was not until 1874 that the new buildings were ready and the Society entered upon the premises it has now occupied for nearly three-quarters of a century. In that year its fellows numbered 733; to-day they number more than seven thousand, and in view of the parallel expansion of numbers both of men and books it is not surprising that further expansion has been in question.

The “communication and discussion of discoveries”, set out as the Society’s first intention, has been

carried out by the reading and discussion of papers, and their publication. It is impossible to estimate the value of the thousands of discussions that have taken place at its meetings, opportunities for the sharing of ideas or the no less valuable clash of opinion, which has so often had the result of leading the research worker to re-examine his conclusions, clarify his explanations and reinforce his experimental proofs. We know that much of the work that led to the adoption of the correct views concerning atomic and molecular weights was done at these meetings, and in some instances by men, such as Odling, whose published work was comparatively unimpressive. These meetings have been of especial value to junior research workers, who find in them at once a means of contact with senior workers of reputation and experience and a strong encouragement in an often ill-rewarded pursuit.

The editors of the *Journal* have likewise done much to maintain the high standard of experimental proof for which the chemists of Britain are justly esteemed. The printing of scientific papers was among the first of the Society’s activities. A volume of *Proceedings* and another of *Memoirs* appeared during 1841-43; these were followed by two volumes of *Memoirs and Proceedings of the Chemical Society of London*, and between 1847 and 1864 by fourteen volumes of the *Quarterly Journal*, which contained not only original papers, but also abstracts and official reports. During 1862-71 a *Monthly Journal* was published. Space was always a difficulty, and during this period abstracts were no longer published, but in 1871 “Abstracts of Chemical Papers published in British and Foreign Journals” were included in the *Journal*. From 1878 onward the *Transactions* and *Abstracts* were separated, while from the year 1885 the *Proceedings*, which had previously been scattered through the *Journal*, were separated and form a record of the Society’s business and activities. After 1914 these were published in an abbreviated form with the *Journal* and no longer contained summaries of papers. In 1924 the publication of abstracts became the province of the Bureau of Abstracts and ceased to be part of the *Journal*. Finally, in 1933 the *Journal* adopted that larger format which is familiar to-day. A most valued publication of the Society is the *Annual Reports on the Progress of Chemistry*, first published in 1905; and in the present year a new publication has commenced, namely, *Quarterly Reviews*.

Next to the reception of the publications of the Chemical Society, the most valued privilege of its fellows is the use of the library. In 1842 the secretary was empowered to obtain a bookcase. The Society received many periodicals in exchange for its own, but in 1850 the library consisted only of some 330 volumes and a hundred pamphlets. In 1855 the president, Dr. Williamson, directed the attention of fellows to the library “as a place of deposit for curious works upon chemistry”. In 1860 Mr. Watts took charge of the library, and a Library Committee was first appointed in 1866. He was succeeded by Dr. L. T. Thorne in 1884, who was followed in 1894 by Mr. R. Steele. In 1903 Mr. F. W. Clifford, assistant librarian at University College, was appointed librarian, and held that office until his death in 1942. His kindly and efficient aid and his store of remarkable anecdotes will long be remembered by the present generation of British chemists. He was succeeded by Mr. A. E. Cummins, who had been assistant librarian since 1919.

By 1890 the library, exclusive of pamphlets, amounted to 9,884 volumes; during Mr. Clifford's tenure of office, it greatly expanded and became the finest chemical library in the world. The number of books to-day amounts to some 48,000, and their issue to fellows, which seems to have been customary from the first, is a most valuable service.

In 1853 it was decided to add "a little more life and general interest to the meetings by devoting up to five meetings a year to delivery of discourses on matters connected with the progress of Chemistry". A good many lectures were given, but only a small proportion of them were printed; however, as these lectures became fewer and at the same time more weighty, their publication became commoner. Five lectureships have since been founded, of which the oldest is the Faraday Lectureship, which is normally given every three years and originally had to be delivered by a 'foreigner or someone from a British Colony'. The first lecturer was Dumas; he was followed by Cannizzaro, Hofmann, Wurtz, Helmholtz and Mendeléeff. Since then, Rayleigh, Ostwald, Emil Fischer, Richards, Arrhenius, Millikan, Willstätter, Bohr, Debye, Rutherford and Langmuir have delivered the Faraday Lecture, and the eighteenth of these lectures is to be delivered on July 16 by Sir Robert Robinson on "The Development of Electrochemical Theories of the Course of Reactions of Carbon Compounds". Since 1918 four other lectureships have been founded, namely, the Hugo Müller, the Liversidge, the Pedler and the Tilden Lectureships.

The Society also gives a very practical and direct aid to research through its Research Fund, which expends an annual sum of about £7,000 in grants towards the expenses incurred by research workers.

The accomplishment of the Chemical Society is principally to be measured by the benefit that has accrued to its members' work from the amenities and facilities it has provided. We cannot estimate the intensity of this benefit, but we may form some notion of its extensiveness by considering the work that British chemists have done and are doing. A brief survey of such work must give a false impression, for it can offer no idea of the enormous volume of pieces of work, not always notable by themselves, but in the mass forming the solid structure of the science of chemistry. The principal impression that is given by a survey of British chemistry during the past century is the great increase in its volume and value after the year 1900. Not only was more work done, but also its importance relative to the work being done at the same time in other countries greatly increased. In the nineteenth century there were fine pieces of work, such as Graham's treatment of colloids, Roscoe's researches on vanadium and Perkin's syntheses of aromatic dyestuffs, but from the beginning of the twentieth century the contribution of British chemists rapidly increases. The remoter causes of this expansion are not altogether clear, but there was certainly a wave of opinion near the turn of the century which led to a more active encouragement of research at the universities of Britain, and especially at Oxford and Cambridge. Be this as it may, the facts are sufficiently impressive. Witness, for example, the discovery of the inert gases by Rayleigh, Ramsay and Travers, the elucidation of the phenomena of radioactivity by Rutherford and Soddy, the opening up of the great field of crystal chemistry by the Braggs, the epoch-making researches of Moseley. In the field of organic chemistry the electrochemical theory of the course of organic

reactions is the fruit of British researches, especially those of Robinson, who has contributed so brilliantly to so many departments of organic chemistry. It would be invidious to single out one or two researches from the mass of brilliant organic work that to-day occupies the greater part of the *Journal of the Chemical Society*, but the work of present members of the Chemical Society upon the carbohydrates, the steroids, proteins, and other classes of organic compounds, is well enough known to all. The subject of chemical kinetics has been remarkably developed in Britain, in numerous centres, and is represented in the person of this year's president, Prof. C. N. Hinshelwood.

All this work has contributed to, and has been aided by, the great development of the British chemical industry, which, after a period of stagnation in the latter part of the nineteenth and early twentieth century, has been built up to a position second only to that of the United States. Never has British chemistry seen a more brilliant period than the present, and we may look forward with confidence to newer and greater conquests, both in our knowledge of things, and their utilization for the benefit of mankind.

TECHNIQUE OF X-RAY POWDER PHOTOGRAPHY*

By H. P. ROOKSBY

Research Laboratories of The General Electric Co., Ltd.,
Wembley

THE aspects of technique that influence the quality of the X-ray powder photograph can be classified into two groups: those concerned with the kind and purity of the X-radiation; and those concerned with the camera. A battery of X-ray tube target materials is desirable to give the necessary flexibility in the selection of wave-lengths. Demountable tubes in which targets are interchangeable, or sets in which sealed-off X-ray tubes containing different target elements can be quickly interchanged, provide alternative lines of approach.

In both schemes, some deterioration of the purity of the spectrum emitted by the tube has to be faced, though with the demountable set the provision of efficient evacuating equipment and regular cleaning of the target face minimize the difficulties. Nevertheless, it has to be recognized that a pure characteristic spectrum can only be maintained with certainty by the use of a crystal monochromator.

With Debye-Scherrer cameras mounted in the conventional position with respect to the X-ray tube target, line definition is primarily a function of specimen diameter and focal distance. There is a gain in resolution when the focal distance is lengthened by increasing the camera diameter. Cameras of maximum diameter of about 20 cm. have been adopted during the last few years; these are fitted with knife edges to provide reference shadows on the exposed negative. Accuracy of construction is essential to enable extrapolation methods to be applied for lattice parameter measurements¹. Clarity of the powder photograph is improved by evacuation or hydrogen filling, and in some instances screening will help by the absorption of the secondary fluorescent radiation excited in the specimen.

* Abridgment of a lecture before the X-ray Analysis Group of the Institute of Physics delivered on April 22.