The possible marked extension of the grazing season and the expanded opportunities for conservation of the surplus product of highly valuable protein food-stuff affords hope of a greatly enhanced, even revolutionary, level of animal production. Sir George Stapledon, in a noteworthy address, suggested a ten-year plan for more livestock to utilize his envisaged 9 million acres of leys and 4 million acres of permanent grass against the present 10 million permanent and 3 million leys—and asked for guidance as to numbers, kinds and quality for this target. Yet he, adding his emphasis to that offered by many of the demonstrations, recognizes the extraordinary niceties of reaction of the plant strains and mixtures to all points of management, and implied doubts whether a suitable technique of testing the absolute value of the pastures can be derived unless they are grazed at different intensities of grazing, and admitted that at times, for suitable conditions of conservation of surplus growth, drastic treatment by cutting has to be adopted.

The main use of grassland is for grazing—but that platitude over-simplifies many of the real issues. Experimentally it may justify using the grazing animal to measure pasture productivity in terms of live-weight increase or milk yield, which terms achieve an apparent validity since they are involved in economic or practical returns. However, it has also been clearly shown that grazing in itself is a powerful factor in altering the composition and 'utility' of the grazed pastures. Pasture management has long been realized to involve control of the biotic factor, but from the animal physiologist's point of view the nature of this complex influence is by no means sufficiently understood for it to be used as anything more than a very general and indirect measure of productivity. This is especially the case when there is a dearth of experimental evidence on the important question as to how far the very conditions of restraint, implied in using animals to graze small plots under trial conditions, themselves affect the animals' reactions, behaviour, growth, development and output, as of live-weight.

The ultimate purpose is to translate experimental results into farm practice. This introduces another set of, as yet, imponderable factors. The field is a unit only in so far as it is part of the farm; the grazing animal also is part of a herd or flock. For the farm as a whole—even for the country as a whole—the supply of winter feed is the limit to production; to widen this restriction, forage crops and conserved grassland in the form of silage or dried grass offer prospects of greater development. But a better and more efficient alternative in present circumstances, in Dr. John Hammond's view, might be to utilize the high production of the summer months for cheaper summer fattening, and to conserve and store the animal product rather than the grass. Again, the numbers of animals on a farm are relatively fixed over long periods; under practical conditions, especially in these days of self-contained dairy herds, the farmer cannot arrange the timing or intensity of grazing to meet those niceties of management which the grassland expert dictates as optima for the good of the sward. As Prof. W. C. Miller suggested, the complex situation of stocking and stock-management may require more complex mixtures of grazing constituents to provide pastures which can be practically mismanaged with safety, rather than a multiplicity of mixtures and treatments, each needing specific control to evoke their best response.

So long as conditions vary from farm to farm, field to field, and season to season, the principles of stock and grazing management must be allowed very flexible application. Even those of rotational grazing, with subdivision of the areas into smaller units, particularly in so far as it involves the human factor of judgment of pasture productivity, are not fully understood, as the plots at the College Farm at Nantcellan show. The formal and informal discussions at these meetings proved that a great deal of endeavour remains to be applied to the problems of the pastures we grow, the uses to which they can be put, and our methods of using them—and that any mutually exclusive interpretation of the results of experiment and observation by animal and grassland specialists is no longer admissible.

J. E. NICHOLS

THE RADIATIONS FROM THE EARTH'S ATMOSPHERE

IN the latitudes of London, Paris or Washington the appearance of an aurora borealis is somewhat infrequent and finds few investigators prepared with equipment for detailed observation. Most of the work on the location and spectra of aurora has therefore been done at such places as Tromsø, which are within the main auroral belt: our knowledge of the phenomenon is mainly due to Størmer and Vegard, and their collaborators. The principal progress has been achieved in the present century, and especially in the last ten or fifteen years. It has only been rendered possible by the rapid general advances in physical techniques and equipment.

During the first quarter of this century, it began to be recognized that at all latitudes the night sky is appreciably luminous even in the absence of the moon. This effect is partly due to radiation from the stars, both those individually visible and those much more numerous stars too faint to be detected by the unaided vision; the zodiacal light, which is of the nature of scattered sunlight, also makes a contribution. But the major part of the luminescence originates in the atmosphere itself: its spectrum shows a complex set of emission lines and bands against a continuous background. The intensity is very low, so the observational work is extremely difficult. Slipher, who was among the first to study the spectrum of the night sky light, discovered that a permanent component of it is the famous green line previously known to be prominent in the auroral spectra. Rayleigh, who used characteristically simple methods, devised and carried out, partly with the aid of observers overseas, a many-sided programme of observations. By means of filters he distinguished three different spectral regions and studied their geographical distribution and their temporal variations. He even measured the absolute intensity of the green line: the number of photons he found to be emitted in a square centimetre column of atmosphere is about 2×10^8 per sec. This remarkable pioneer work gave results of lasting value.

The problem of the identification of the green line was solved by McLennan and his colleagues who, in the years 1925–27, showed that it arose from atomic oxygen and is due to what is called a forbidden transition—that is, one that violates the normal selection rules and is only possible because of their approximate nature. Now the emission of the green-

line radiation does not bring the oxygen atoms involved to their ground-state; it instead brings them to the lowest excited state. In final passage from this state, radiation may be emitted corresponding to certain red lines. These have now been found both in the night sky and auroral spectra. It is possible that the forbidden lines of atomic nitrogen are also present.

More recently, in 1938, there came from France the astonishing news that sodium occurs in the high atmosphere in the free atomic form. Proof for this was provided by the identification of its characteristic yellow line (or rather doublet).

In addition to the atomic lines a large number of bands appear. Those in the spectra of auroras can be ascribed mainly to neutral and singly ionized molecular nitrogen. Much work has yet to be done on the interpretation of the night sky spectrum; but it seems probable that the principal contribution to the visible region is from neutral molecular nitrogen, and the principal contribution to the ultra-violet region is from neutral molecular oxygen: the bands of singly ionized molecular nitrogen seem to be normally absent. An interesting discovery, made in 1942 by Herman, Herman and Gauzit, is a remarkably strong emission in the infra-red region: its identification is still in doubt.

A growing number of highly skilled observers have been drawn into this branch of research and have obtained many unexpected and varied results. These have aroused the interest of geophysicists, astrophysicists, laboratory spectroscopists, theoretical physicists, radio scientists, chemists, and mathematicians. One manifestation of this interest was an international conference on the "Emission Spectra of the Night Sky and Auroras" organised under the auspices of the Gassiot Committee of the Royal Society and held in the Society's rooms at Burlington House, London, during July 7-10.

The Gassiot Committee is charged with the supervision and planning of a long-term programme of research on the radiation in the atmosphere and on related phenomena. This commitment was undertaken at the request of the Air Ministry (on the advice of the Meteorological Research Committee). Financial aid is given by the Ministry. The initiative in proposing the conference was taken by Prof. H. S. W. Massey, chairman of one of the sub-committees appointed by the Gassiot Committee to deal with different sections of the general programme.

The conference was attended by guests from Belgium, France, Norway, Sweden and the United States, and by many from the British Isles. Russian scientific workers have done much work in the field concerned, but none was able to be present: the Embassy, however, sent an observer. The papers read at the conference will be published by the Physical Society in the next issue of its "Report on Progress in Physics". Here only an outline of the ground covered will be attempted.

The best recent observations of the night sky spectrum have been made in France and the United States. Papers describing the results were contributed by G. Dejardin, J. Dufay, Teheng Mao-Lin, A. and E. Vassy, R. Grandmontagne, C. T. Elvey and E. O. Hulburt. There was considerable discussion of the altitude of the luminous region, which is still very uncertain. Much attention was given to the special character of the spectrum at twilight, when the negative bands of nitrogen, the red lines of oxygen, the yellow doublet of sodium and possibly other

components are enhanced. A. Kastler described work by J. Bricard and himself on the polarization of the light of various atomic lines using a very simple polariscope due to Savart and improved by Lyot. The green oxygen line was found to show less than 3 per cent polarization—a result to be expected from theory, but contradicted by recent measurements in the U.S.S.R. D. Barbier, R. W. B. Pearse and others discussed the identification of the spectral components. A great deal of time was devoted to the complex and fascinating task of explaining the cause of the emission. D. R. Bates, H. S. W. Massey, and M. Nicolet contributed papers relating to this problem. It seems probable that the emission is due in part to the dispersion of the energy stored up by the dissociated oxygen (and perhaps also nitrogen) formed during the day: recombination of the charged particles in the ionosphere will play a minor part. Consideration has also to be given to the suggestion that incident particles may have to be invoked to explain such effects as the irregularity of the light. Much remains to be elucidated. P. M. S. Blackett presented an interesting paper on the possibility of cosmic rays making an observable contribution to the continuum through Cerenkov radiation.

Recent observational work on auroras was reported by G. Kvitte (on behalf of L. Vegard) and by R. Bernard. Confirmation was given to the suggestion that hydrogen and helium may be present. An important new analysis of the spectrum was given by M. Nicolet. K. G. Emeléus read a paper sent by J. Kaplan describing laboratory experiments on the reproduction of the characteristic emissions.

It is generally accepted that auroras are due to the action (direct or indirect) of charged particles guided pole-wards by the earth's magnetic field. However, the nature, origin and energy of these is still the subject of controversy. Widely different views were expounded in papers by H. Alfvén, S. Chapman, and F. Hoyle.

One session of the conference was devoted to a discussion, led by R. W. Ditchburn, on the instruments and techniques of observation. E. O. Hulburt gave a valuable account of the work being done in the United States on the exploration of the upper atmosphere by means of rockets.

S. CHAPMAN
D. R. BATES

OBITUARY

Prof. J. Perner

THE death occurred at Prague, on June 9, of Dr. Jaroslav Perner, professor of palaeontology at the Charles University. Born at Tyneć-nad-Labem on March 28, 1869, Perner claimed to be a pupil of the French geologist, Joachim Barrande (1799-1883), in whose footsteps he followed by continuing the investigation of the fossil fauna of the Silurian and Devonian rocks of middle Bohemia. His studies were incorporated in various publications, including "Système silurien du centre de la Bohême" and a work of four volumes entitled "Etudes sur les graptolites de Bohême". Most of his other publications were in Czech or German.

Early in life Perner came to England and spent a period in the 'nineties at Mason College (afterwards the University of Birmingham) working under the late Prof. Charles Lapworth. J. G. F. D.