effect was observed, and the Coombs test (with rabbit anti-human serum) was negative when barely agglutinating doses of the serum were mixed with *Rh'*, *Rh''* and *Rh* negative cells which were agglutinated by undiluted serum. In this respect the antibody resembles the α , β and Lewis agglutinins but differs from the *Rh* agglutinins.

A full account of our investigations will be published in Australia.

R. J. WALSH

CARMEL M. MONTGOMERY

Red Cross Blood Transfusion Service,
Sydney. July 29.

Subdivisions of the MN Blood Groups in Man

A SAMPLE of the unique serum described in the preceding letter was kindly sent to us by Dr. Walsh and Miss Montgomery. We found that the antibody was not connected with the *ABO*, *Rh*, *P*, Lutheran, Kell or Lewis systems, but that it did subdivide the *MN* blood groups.

The donor of the serum is group *MN* and her red cells are not agglutinated by her own serum. The agglutinin is active at 37° C., and at lower temperatures.

One hundred and ninety unselected English bloods have been tested for agglutination by this serum, with the following results:

	<i>M</i>		<i>MN</i>				<i>N</i>		Total
	+	-	+	-	+	-	+	-	
	36	14	57	38	15	30	190		
	19%	7%	30%	20%	8%	16%	100%		
Inter-pretation	<i>MS MS</i>	<i>MM</i>	<i>MS N</i>	<i>MN</i>	<i>NS NS</i>	<i>NN</i>			
	<i>MS M</i>		<i>MS NS</i>		<i>NS N</i>				
			<i>M NS</i>						

+ = agglutinated by the new serum.
- = not agglutinated.

It is clear that the antigen identified by the new antibody is not independent of the *MN* system; for of the *M* containing bloods 93 are positive and 52 negative, while of the non-*M* containing bloods 15 are positive and 30 are negative ($\chi^2 = 13$ for 1 d.f.).

These results can be explained on the supposition that there are four allelomorphs at the *MN* locus, *M*, *MS*, *N* and *NS*; *MS* and *NS* being positive and *M* and *N* negative in respect of the new antibody. We may picture *S* as being a change, or mutation, that has happened to some *M* and to some *N* genes, rendering the resulting red cells agglutinable by the new antibody.

On the other hand, there is no proof that *S* is not a gene closely linked to the *MN* locus; such a gene would presumably have an allelomorph *not S* (or *s*). If a serum anti-*not S* (or anti-*s*) is found, the latter explanation will then be the more probable, and the situation will be similar to that of the *C*, *D* and *E* antigens of the *Rh* system.

If this is an example of linkage, the genes must be close together, otherwise crossing-over would presumably have resulted in equal frequencies of *S* in *M* and *N* chromosomes.

Whatever the true interpretation may be, the gene frequencies of the forms *M*, *MS*, *N* and *NS* can be calculated from the above data, and Prof. R. A. Fisher has very kindly applied his maximum likelihood method for *Rh* gene frequencies¹ with the following results:

$$\begin{aligned} M &= 0.262671 \\ MS &= 0.250487 \\ N &= 0.394701 \\ NS &= 0.092121 \end{aligned}$$

From these gene frequencies the expected genotype frequencies are calculated.

	Expected	Observed	χ^2
<i>MS MS</i>	36.9236	36	0.02310
<i>MS M</i>	13.1093	14	0.06052
<i>MM</i>			
<i>MS N</i>	55.5351	57	0.03864
<i>MS NS</i>			
<i>M NS</i>	39.3991	38	0.04970
<i>MN</i>	15.4300	15	0.01198
<i>NS N</i>	29.6029	30	0.00533
<i>NS NS</i>			
<i>NN</i>	190.0000	190	0.18927
			d.f. 2,
			$p = 0.9$

It will be seen from the table that the agreement between the expected and the observed numbers is very close.

A fuller account of this work will be published elsewhere and will include the results of family investigations which further support our interpretation; for example, a family has been found of the type *MS N* $\times$ *MM* showing segregation of *S* with *M*, and one of the type *M NS* $\times$ *NN* showing segregation of *S* with *N*.

RUTH SANGER
R. R. RACE

Medical Research Council,
Blood Group Research Unit,
Lister Institute of Preventive Medicine,
London, S.W.1. Aug. 5.

¹Fisher, R. A., *Ann. Eugen.*, 13, 150 (1946).

The Blood Groups of the Basques

THE linguistic uniqueness of the Basque people and their physical resemblance to some of the late palaeolithic inhabitants of Europe have long been known. As Boyd and Boyd¹ have shown for the Spanish Basques and as Vallois² has shown for French populations with a large Basque element, they have a high frequency of group *O* persons. In this they resemble the Icelanders, the Scots, the Irish, the northern Welsh and the Sardinians. They differ from these and from all the other peoples of Europe yet examined in their very low frequency for the *B* gene. The facts so far cited tend to show that the Basques are a relict population which at least in Spain has suffered no significant admixture of elements akin to the general western European population. It remains to be determined whether, on the other hand, a race akin to the Basques forms a significant element in the population of western Europe.

In spite of considerable variations in the frequencies of the *ABO* blood groups, the frequency of *Rh*-negative persons scarcely varies from 16 per cent in any large western European population yet examined. In Fisher's³ terminology *Rh*-negatives are homozygous *dd*; the frequency of the *d* gene is thus $\sqrt{0.16}$, or 40 per cent, and that of its allelomorph, *D*, 60 per cent. Most non-European populations are almost exclusively *Rh*-positive, and until recently no population was known anywhere with an *Rh*-negative frequency much above 16 per cent.

As Haldane⁴ and Wiener⁵ have pointed out, every infant that dies of haemolytic disease due to the common type of *Rh*-incompatibility is heterozygous *Dd*, and such deaths destroy equal numbers of *D* and *d* genes; thus the rarer of the two genes will in time be eliminated. The present *Rh* ratio in western Europe is thus an unstable one, and one is led to postulate a mixing of populations predominantly *Rh*-positive and *Rh*-negative respectively.