P. Fremontii, *P. Wislizenii* and *P. Eugenioi* showed only very weak fungicidal activity; it seemed first that the activity was restricted to the balsam poplars; but recently *P. regenerata*, *P. serotina erecta* and *P. marilandica* were found also to give active extracts.

The bark of *Alnus glutinosa*, *Fagus sylvatica* and *Quercus pedunculata* showed (in the winter) almost no activity at all.

The leaves of *P. candicans* showed also fungicidal activity, though weaker than the bark of this species.

Fungicidal activity of plant extracts is not uncommon, but in general the leaves show more activity than the bark². In trees of the genus *Populus* the distribution of inhibitors in the plant seems to be different.

J. GROSJEAN

Instituut voor Plantenziektenkundig Onderzoek,
Wageningen.
Jan. 16.

¹ Grosjean, J., *Tijds. Plantenziekten*, **49**, 172 (1943); see p. 176.

² Gilliver, K., *Ann. App. Biol.*, **34**, 136 (1947); see p. 142.

Proliferative Capacity of the Hypophyseal Portal Vessels

THE pathway by which the hypothalamus influences the secretion of the anterior pituitary gland has been the subject of much research. The possibility that the hypophyseal portal vessels may form part of this pathway has been discussed¹, and the anatomy of these structures in the rat and other forms has been described. In 1949, Harris² observed that regeneration of the hypophyseal portal vessels in the rat may follow section of the pituitary stalk, and later³ that post-operative return of reproductive activity can be correlated with such regeneration. In this work the hypophyseal stalk was exposed by the sub-temporal route and either cut with a knife or torn with a hook. In the literature, accounts of stalk section made by a variety of methods show divergent results. Since the conditions for a regeneration of the portal vessels may vary according to the method used for severing the hypothalamo-hypophyseal connexions, we have investigated the regenerative capacity of these vessels in different circumstances.

Regeneration of the hypophyseal portal vessels and the state of the endocrine glands have been studied in rats. The experiments are divided into four groups:

(1) Hypophysectomy and transplantation of anterior lobe tissue under the median eminence (12 experiments).

(2) Hypophysectomy and transplantation of the extirpated anterior lobe into the emptied hypophyseal capsule (16 experiments).

(3) Extirpation of the anterior pole of the hypophysis (13 experiments).

(4) Section of the hypophyseal stalk (25 experiments).

Twelve hypophysectomized rats served as control material. The duration of the experiments varied from 1 to 104 days.

The observations show that a hypophyseal transplant near the hypophyseal stalk in the subarachnoid space becomes richly vascularized from the portal vessels. Regeneration of the portal vessels after stalk section by the sub-temporal route has been confirmed, and previous studies of the activity of the anterior lobe³ extended. It seems clear that in the subarachnoid space the portal vessels can undergo pro-

liferative and reparative changes. Further work is in progress concerning the cytology and activity of these transplants.

The evidence obtained from the experiments on rats with a transplant in the hypophyseal capsule, and on rats with the anterior pole of the hypophysis removed, indicated that capillaries derived from the portal vessels may invade such isolated anterior lobe tissue in the capsule. However, anterior pituitary tissue transplanted into the capsule did not become vascularized or active. Since the possibilities of obtaining living transplants of anterior lobe tissue inside the hypophyseal capsule are highly dependent on technical factors, other techniques are being investigated.

G. W. HARRIS

DORA JACOBSON

Physiological Laboratory,
Cambridge.
Dec. 27.

¹ Harris, G. W., *Physiol. Rev.*, **28**, 139 (1948).

² Harris, G. W., *Nature*, **163**, 70 (1949).

³ Harris, G. W., *J. Endocrinol.*, **6**, xviii (1949a).

Autonomic Nervous System of the Ratfish

DETAILED accounts are available for the autonomic nervous system of various Selachians¹⁻⁴, but little is known of the same system in the Chimaeroid fishes or Holocephali. To remedy this defect a study has been made of the autonomic nervous system of the Chimaeroid *Hydrolagus coliei*. This species is common in British Columbian inshore waters⁵.

The autonomic system of *Hydrolagus* comprises a sympathetic portion in the trunk, and a parasympathetic cranial portion. In the sympathetic system, segmentally arranged vertebral ganglia occur from the level of the heart to the anus. The ganglia are closely associated with small suprarenal bodies, and they are connected together by a loose plexus lying in the dorsal body wall. A regular chain and commissures between ganglia are absent. The largest ganglia are the gastric ganglia; these are joined to large masses of suprarenal tissue on the subclavian arteries. White rami communicantes connect the sympathetic ganglia to the spinal nerves. The sympathetic ganglia give rise to visceral nerves supplying blood vessels, genital ducts, suprarenals, intestine, liver and pancreas. A sympathetic extension into the head and tail is absent, and there is no sympathetic innervation of the heart.

The parasympathetic system lies in a midbrain and a hindbrain outflow. The former passes through the oculomotor nerve to an adherent ciliary ganglion; from the latter, short ciliary nerves proceed to the eyeball. The hindbrain outflow comprises ganglia in the glossopharyngeal and vagal nerves. The vagus also supplies the heart and intestine, and connects with the sympathetic system. A true sacral parasympathetic system is wanting; but motor fibres from posterior abdominal spinal nerves directly innervate smooth muscle of the rectum without the interpolation of post-ganglionic neurones.

The autonomic nervous systems of Selachians and Chimaeroids are remarkably similar, despite the fact that the two groups have preserved an independent evolution since the Palaeozoic⁶. It would appear that this kind of autonomic system represents a simple level of organisation present in the common ancestor