PLANETARY ATMOSPHERES

The Atmospheres of the Earth and Planets

Papers presented at the Fiftieth Anniversary Symposium of the Yerkes Observatory, September 1947. Edited by Gerard P. Kuiper. Pp. vii+366+16 plates. (Chicago: University of Chicago Press; London: Cambridge University Press, 1949.) 60s net.

ASTROPHYSICS has produced some exciting chapters during the past half-century. One of the latest, and certainly not the dullest, concerns the atmospheres of the planets. Twenty-five years ago the whole subject was still very obscure, while even with the atmosphere of most concern to us, that of the earth, our knowledge was comparatively limited. Since then, by brilliant spectroscopic work in the visible and infra-red, important constituents of the atmospheres of Venus, Mars, Jupiter and Saturn have been identified. We know of abundant carbon dioxide on Venus, but can find little trace of oxygen or water vapour; of smaller quantities of carbon dioxide on Mars, with again extremely little oxygen or water vapour; of methane and ammonia in the major planets, with evidence for ice as a major constituent of the rings and inner satellites of Saturn. More than this, it has been discovered that the spectrum of the polar cap of Mars resembles that of hoar frost, and that the greenish areas on Mars do not show the absorption spectrum of chlorophyll, although it is still possible they may be due to lichens or mosses. The spectrum of the browner areas of this planet has been matched with that of a brownish felsite, but has been shown to be quite different from that of various other terrestrial soils and rocks of about the same colour.

Solid, or at least non-gaseous, particles are present in all planetary atmospheres. Those on Venus are not yet identified; on Mars there is a variable haze which may be similar to very fine cirrus cloud, and quite possibly composed of ice crystals. The clouds on Jupiter and Saturn can be explained as ammonia crystals, in an atmosphere containing much hydrogen. The optical properties of such atmospheres are being linked up with those of the terrestrial atmosphere and even of interstellar space.

But the astronomical discoveries are in a sense only a part of the story, which now involves several other sciences—meteorology, radio physics, geology and even ballistics—and this is specially true of the atmosphere of the earth. This we think of as a complicated heat machine, driven by solar radiation, showing great variability in detail, but apparently of great long-term stability. The evidence available suggests little fundamental change since the beginning of geological time, although in view of the great discrepancies in abundance of individual elements (when compared with the major planets, the sun, stars or nebulae) present opinion inclines to the belief that our atmosphere is secondary, perhaps in part produced by life itself. Here a paradox is encountered: the atmospheric oxygen of the earth may be chiefly a product of living matter, yet this same oxygen is the principal agent which filters out the lethal ultra-violet from the solar spectrum.

We realize that our pictures of the atmospheres of other planets are excessively simplified (as are most astronomical pictures) by lack of data. On the earth, however, we are familiar with the bewildering complexity of meteorological phenomena in the lower atmosphere, while the study of meteors, of the night-sky spectrum, of auroras, and of the reflexion or

transmission of radio waves, shows a no less complicated multi-layer upper atmosphere, its properties controlled by the absorption spectra of various gases, notably oxygen and ozone. We link disturbances in these higher layers with events on the sun, the effects of which are transmitted in part by ultra-violet light and in part by corpuscles the identity of which is still uncertain. Our ideas are now being tested by soundings with rockets, the results so far substantially confirming the picture built up from the other sources just named. These rocket flights are permitting an extension of our direct knowledge of the ultra-violet solar spectrum, which at sea-level is cut off by the atmosphere at about 3000 Å. Photographs have been taken of the spectrum down to 2200 Å., showing thus far a rapidly weakening continuous spectrum, heavily blanketed by absorption lines. There is as yet no trace of the strong radiation, probably arising in the chromosphere and corona, and of much shorter wavelengths, believed to be responsible for ionospheric disturbances.

Study of our upper atmosphere leads us to the belief that the temperatures there are high, perhaps up to 1,500° K., and this re-opens questions as to the loss of gases from the top of the atmosphere. What is the nature of the equilibrium when atomic collisions become relatively infrequent, and to what extent are there similarities with the outer atmospheres of the sun and other stars?

All these matters, and a great deal more, are discussed in the book under review, which is a collection of papers given at a symposium on planetary atmospheres held at the University of Chicago, in September 1947, in connexion with the fiftieth anniversary of the Yerkes Observatory. The inspiration for the symposium came from Prof. G. P. Kuiper, who has been responsible for some of the most important developments of the subject, and who has also edited this book. I have myself no great liking for symposia, but if they all were to succeed in producing such a wealth of exciting and significant contributions as this one, I would willingly change my opinion. Much of the work recorded here is of outstanding quality, and the greater part of it is American in origin; one of the pleasing virtues of a good book is that the authors take little notice of the first of these facts and none at all of the second. R. O. REDMAN

CONGRESS OF THE EUROPEAN BREWERY CONVENTION

European Brewery Conference

Congress, Lucerne 1949. Vol. 1. Pp. ix+273. (New York and Amsterdam: Elsevier Publishing Co., Inc.; London: Cleaver-Hume Press, Ltd., 1949.) 37s. 6d.

THE European Brewery Convention was set up in 1947 with the object of "co-ordinating the scientific work done in different countries in the technical fields of Brewing and Malting". This volume contains three symposia—the papers submitted at the second biennial congress of the Convention.

In the first of these symposia, proteins in brewing, E. Sandegren (Sweden) reports recent advances in the work of fractionating barley proteins which has been proceeding in Uppsala and Stockholm in recent years. Salting-out with phosphates was followed by the application of the polarographic method to the