

Physiological Effects of Trihydroxy-N-Methylindol and its Reaction with Copper Ions

TRIHYDROXY-N-METHYLINDOL, a highly fluorescent substance derived from adrenochrome¹⁻⁴, sensitizes the smooth muscles of the cat to the excitatory (nictitating membrane) and inhibitory (non-pregnant uterus) actions of adrenaline⁵.

This substance has a great affinity for copper ions. The fluorescence of a 2×10^{-6} solution disappears in three minutes at laboratory temperature in the presence of 1×10^{-6} copper ions (as copper sulphate); it inhibits the oxidation by copper ions of adrenaline in aqueous solution. A substance has been isolated and analysed which corresponds to the formula $C_9H_7NO_3Cu + H_2O$ or $C_{18}H_{14}N_2O_6Cu_2 + 2 H_2O$ (found: C, 42; H, 3.6; N, 5.1; $C_{18}H_{14}N_2O_6Cu_2$ requires C, 41.5; H, 3.5; N, 5.4 per cent).

The hypothesis may be put forward that the physiological actions of trihydroxy-N-methylindol (decrease of bleeding time⁶ and of capillary permeability, sensitization to adrenaline) are related to its copper-binding activity in accordance with the general concept formulated by Lavollay⁷, Javillier and Lavollay⁸, Clark and Geissman⁹, in order to explain the 'vitamin P'-like actions of flavonoids and related substances.

P. FISCHER
J. LECOMTE
L. DELANDTSHEERE

Laboratory of General Pathology,
Laboratory of Medical Pathology,
University of Liège. Aug. 3.

¹ Harley-Mason, J., *J. Chem. Soc.*, 1276 (May 1950).

² Lund, A., *Acta Pharm. Toxicol.*, 5, 121 (1949).

³ Fischer, P., *Bull. Soc. Chim. Belgique*, 53, 206 (1949).

⁴ Fischer, P., Derouaux, G., Lambot, H., and Lecomte, J., *Bull. Soc. Chim. Belgique*, 59, 72 (1950).

⁵ Fischer, P., and Lecomte, J., *Arch. Int. Physiol.*, 58, 236 (1950).

⁶ Fischer, P., and Trabert, P., *Experientia*, 6, 392 (1950).

⁷ Lavollay, J., "L'autooxydation des Diphénols" (Paris: Hermann, 1943).

⁸ Javillier, M., and Lavollay, J., *Helv. Chim. Acta*, 29, 1283 (1946).

⁹ Clark, W.-G., and Geissman, T.-A., *J. Pharm. Exp. Ther.*, 95, 363 (1949).

'Y-Protein': a New Component of the Muscle Machine

We have recently shown¹ that if rabbit muscle is extracted with a solution of high ionic strength (potassium chloride > 0.5), the electrophoretic test (Tiselius-Longsworth, see ref. 2) shows, besides the proteins already described³⁻⁷, a new component that we have called 'Y-protein'. At ionic strength 0.40 and pH 7.4, it moves more slowly than β -myosin.

This new protein exists only in extracts of normal resting muscle. If the muscle has been previously subjected to contracture by pharmacological means (moniodoacetate) or *rigor mortis*, neither Y-protein nor β -myosin can be found in the extracts if potassium chloride is used for the extraction⁸. This is due to the presence of strong linkages between those proteins and muscle stroma, in the contracted state. These linkages cannot be broken by potassium chloride, but are broken by potassium iodide or pyrophosphate solutions¹.

We have succeeded in isolating Y-protein. Normal resting muscle is progressively frozen in a $-20^\circ C$. chamber, cut in 50- μ slides by use of a freezing-microtome⁹, extracted for ten minutes with 5 vol. Weber-Edsall solution (0.6 M potassium chloride, 0.04 M sodium bicarbonate, 0.01 M sodium car-

bonate; the addition of 0.3 per cent adenosine-triphosphate does not affect the amount of Y-protein extracted). Extraction and the following operations are carried out in a cold room at $\pm 1^\circ C$. After centrifugation, the liquid is progressively mixed with saturated ammonium sulphate solution at pH 6.00 (53 c.c. sulphate solution for 100 c.c. extract, which corresponds to 35 per cent saturation).

The electrophoretic analysis of the precipitate shows that it contains α -myosin (actomyosin), Y-protein and some myogenes. β -myosin remains in the solution (it is precipitated only at 37-40 per cent saturation in ammonium sulphate at pH 6.00⁹). The precipitate containing Y-protein is washed with ammonium sulphate solution at 35 per cent saturation and dissolved in 0.1 M potassium chloride at pH 7.6. The solution, which is rather turbid owing to the presence of actomyosin, is dialysed against a solution of μ 0.05 (0.016 disodium hydrogen phosphate, 0.002 sodium dihydrogen phosphate). The actomyosin is precipitated and Y-protein remains in the solution.

Electrophoretic controls show that only one component is present. Its velocity is:

at μ 0.40, pH 7.40: -2.7×10^{-5} cm./volt/sec.

at μ 0.10, pH 7.60: -3.4×10^{-5} cm./volt/sec.

In an extract of whole muscle, in which Y-protein is present as well as all the other muscle proteins, the velocity of the Y-protein is much reduced:

at μ 0.40 and pH 7.40: $2.65 \pm 0.1 \times 10^{-5}$ cm./volt/sec.

Solutions of Y-protein are rather clear, slightly viscous and do not show double refraction of flow. Optical properties do not depend on ionic strength between 0.005 and 0.5.

Y-protein is insoluble in distilled water, fairly soluble in solutions at μ 0.005, pH 7.4. The fact that the new protein can be extracted only by using solutions of at least a hundred times ionic strength indicates that Y-protein is not free in the muscle cell, but connected to some part of the insoluble stroma from which it is liberated by solutions of high ionic strength.

This property explains why Y-protein has not been isolated before. It cannot be present in low ionic strength extracts such as those prepared for isolation of the proteins of the myogen group or the X-globulins. When extracts of high ionic strength are made, they are usually prepared to extract the myosins—which are precipitated by dilution—and Y-protein is not precipitated in those conditions, being still soluble at μ 0.005 M.

That the binding forces which maintain the Y-protein in the muscle cell are considerably increased in the contracted state seems to be a proof that the new protein takes part in the constitution of the muscle machine.

M. DUBUISSON

Laboratory of General Biology,
Faculty of Sciences,
University of Liège. July 27.

¹ Dubuisson, M., *Biochem. et Biophys. Acta*, 5, 489 (1950).

² Dubuisson, M., Distèche, A., and Debot, A., *Biochem. et Biophys. Acta* (in the press).

³ Dubuisson, M., "Les protéines musculaires". Exposés Annuels de Biochimie Médicale, 9ème série (Paris, 1948).

⁴ Dubuisson, M., *Bull. Acad. Sci. Belg.*, 33, 769 (1947).

⁵ Dubuisson, M., *Bull. Acad. Sci. Belg.*, 34, 978 (1948).

⁶ Dubuisson, M., *Biochem. et Biophys. Acta*, 4, 25 (1950).

⁷ Dubuisson, M., "Le cycle de la contraction musculaire et la structure physico-chimique de l'édifice contractile", Congrès de l'Assoc. des Physiol. de Langue Française (Montpellier, June 1949).

⁸ Dubuisson, M., *Experientia*, 3, 372 (1947).

⁹ Dubuisson, M., *Experientia*, 2, 412 (1946).