of the oviduct, and at the higher levels it resembled the fully active organ. On the other hand, in all cases the ovaries were quiescent, indicating that endogenous hormone production was negligible.

The results show that the increase in serum riboflavin evoked by the administration of oestrogen consisted entirely of free riboflavin, and that the levels of combined riboflavin were not affected. They also indicate that changes in serum riboflavin can be followed by estimating the amount of vitamin present in trichloroacetic acid filtrates of the serum, thereby greatly simplifying the analytical procedure.

W. BOLTON

Poultry Research Centre,
West Mains Road,
Edinburgh 9.
July 15.

¹ Common, R. H., Rutledge, W. A., and Bolton, W., *J. Endocrinol.* **5**, 121 (1947).

² Burch, H. B., Bessey, O. A., and Lowry, O. H., *J. Biol. Chem.*, **175**, 457 (1948).

Plasmacytosis of the Bovine Udder during Colostrum Secretion and Experimental Cessation of Milking

THE concomitance of bone marrow plasmacyte formation and induced hypersensitivity was noted by Kolouch¹ in 1938. This was succeeded by a large number of papers indicating the part played by the plasma cell in the secretion of specific antibodies and by clinical studies reporting the strict parallelism between hyperglobulinemia and plasmacytosis in various diseases. These have been reviewed by Kolouch, Good and Campbell². The present communication records that the plasmacytosis in the bovine udder is synchronous with heightened periods of immune-body formation at parturition and during experimental cessation of milking.

The concentration of immune bodies in the colostrum of cows has been investigated by Smith and Little³, who showed that transmission of antibody from dam to calf was made through the colostrum milk, rather than by placental transmission as in the human. The chemical nature of these proteins has more recently been described by E. D. Smith. The great concentration of the immune globulin fraction of the colostrum—many times that of the maternal serum globulin—is noteworthy. Notwithstanding this fact, it has been thought that the source of the immune bodies is in the maternal blood-stream⁴. This work has been reviewed by Porter⁵ and extended by the demonstration that experimentally the antibody content of milk may be caused to increase to the level of that in colostrum by cessation of milking for a week or more.

As the concept of the plasmacytic origin of specific antibodies demands that any such spectacular rise in immune globulin be accompanied by a properly timed wave of plasma cell formation, experiments were set up to make a rigorous test of the hypothesis. Three cows were used for the preliminary observations. Aseptic serial biopsies of the udder were performed with local anaesthesia in which several grams of secretory tissue were removed. These were fixed and sectioned according to usual techniques.

Biopsies of No. 278 (the cows used were from the herd of the Department of Dairy Husbandry, University of Minnesota), which was neither pregnant

Biopsy of mammary gland of cow 854 made on day of calving. Note plasma cells in connective tissue septa between alveoli. Cresyl Violet and Erythrosin stain

nor lactating, showed no plasma cells in the udder. Similar biopsies of No. 854 showed the abundance of plasma cells eight days prior to parturition. These were again found three days before calving, on the day the calf was dropped (see photograph) and four days later. A biopsy 18 days after calving revealed no plasma cells. In like manner, No. E607, late in a normal disease-free lactation, was biopsied and a very small number of plasma cells observed to be present. Milking was discontinued in two quarters of the udder, and four days later a large population of plasma cells was revealed in the biopsy. These were again noted eleven days after milking had ceased. Following this third biopsy, the two quarters were again milked regularly, and fifteen days later only a very small number of plasma cells were to be found.

BERRY CAMPBELL
ROBERT M. PORTER
W. E. PETERSEN

Departments of Anatomy and
Dairy Husbandry,
University of Minnesota.
July 8.

¹ Kolouch, F., *Proc. Soc. Exp. Biol. and Med.*, **39**, 147 (1938).

² Kolouch, F., Good, R. A., and Campbell, B., *J. Lab. and Clin. Med.*, **32**, 749 (1947).

³ Smith, T., and Little, R. B., *J. Exp. Med.*, **36**, 453 (1922).

⁴ Smith, E. L., *J. Dairy Sci.*, **31**, 127 (1948).

⁵ Porter, R. M., thesis, Univ. of Minn., 1948.

Complexes of Calcium with Citric Acid and Tricarballic Acids measured by Ion Exchange

WE are engaged in a study of the composition and stability of the complex ions which result from the interaction between metal ions and anions, particularly those of biological importance. The determinations are made by measuring the degree to which tracer levels ($\sim 10^{-8}$ M) of a radioactive cation are taken up by a synthetic organic cation exchanger in the presence of known concentrations of a complex ion-forming anion¹⁻³. This ion-exchange procedure has several unique advantages¹. For example, the dissociation constants of several different metal-organic acid complex ions can be measured simultaneously in the same solution, no corrections are