about 28° C., and were observed continuously for the achievement of complete wilting (Table 2 and accompanying illustration).

The results can be summarized as follows. Cotton wilting—in culture—is not only stimulated by the metabolites of *Fusarium*, but it may be similarly induced by those of a non-specific saprophytic fungus (that is, *Trichoderma*). In the case of cotton cut shoots, the wilting is even more rapid in *Trichoderma* than in *Fusarium* filtrate. The intervention of roots—in cotton seedlings—may be responsible for the prolongation of the time required to achieve complete wilting. The metabolites, of both *Fusarium* and *Trichoderma*, are found to be thermostable.

This work emphasizes the importance, in such studies of biological control, of considering the relationship between the host plant and the fungal associate responsible for such a biological control. The study of the different aspects of such a relationship elucidates to a great extent the possible role of the fungal associate in influencing the resultant balance between the host plant and the specific fungal pathogen.

MECHANISM OF ENZYME ACTION

By DR. S. J. LEACH and DR. H. LINDLEY

Biochemistry Unit, Wool Textile Research Laboratories, Commonwealth Scientific and Industrial Research Organization, Melbourne

SINCE the original suggestion by Szent-Györgyi¹ that proteins may exhibit properties of semi-conductors, much evidence has accumulated to support this hypothesis. Evans and Gergely² have collected most of the salient points of evidence, and have also provided theoretical justification for the suggestion by showing that a system of hydrogen-bonded peptide links should show banded energy-levels associated with the π -electrons and should behave as a conductor on suitable excitation. It is an obvious extension from this to consider an enzymic protein as a large assembly which promotes interaction between two reactants by virtue of conduction pathways. On such a basis an important aspect of enzymic specificity may be the matching of the energy-levels of substrate and enzyme³. Geissman³ has recently considered these ideas in some detail for a number of enzyme systems.

While we consider this general hypothesis is extremely attractive and explains many aspects of enzyme behaviour, we feel that it is deficient in that it does not explain why enzyme activity is not a property of all proteins. Recent suggestions on protein structure by Pauling, Corey and Branson⁴ emphasize the current view that hydrogen-bonding plays an important part in the architecture of proteins generally. Moreover, there is evidence from infra-red studies that the degree of hydrogen-bonding of the —NH residues of pepsin is greater in the denatured (inactive) form than in the native (active) state⁵. It is apparent, therefore, that something more than the existence of a strongly hydrogen-bonded system of peptide links is necessary in order to confer enzymic properties on a protein molecule. It

is also obvious that the most likely source of such properties lies in the order and arrangement of the amino-acid side-chains of the enzyme molecule.

One of the great merits of the hypothesis of enzyme action outlined above is that it provides a general mechanism for all types of enzyme reaction, and hence it is possible that all enzymes may exhibit similar regularities in the arrangement and character of some of their side-chain residues. Such regularities, if they exist, might be associated particularly with the amino-acid side-chains which are capable of forming hydrogen-bonds and thus building up inter-chain networks of electron pathways. Examples of such side-chains are those of the dicarboxylic acids and their amides, the hydroxy acids including tyrosine, cysteine, tryptophane and histidine. We have accordingly examined published data on the amino-acid composition of enzymes and non-enzymic proteins, and the accompanying table lists three regularities we have discovered.

Although no single one of these relationships is entirely characteristic of enzymes, avidin and possibly γ -globulin are the only two examples of non-enzymic proteins (out of a total of twenty-three) which give values for all three relationships similar

RATIOS OF COMBINATIONS OF THOSE AMINO-ACIDS IN ENZYMES AND OTHER PROTEINS WHICH ARE LIKELY TO BE CONCERNED IN THE FORMATION OF HYDROGEN-BONDS

Protein	Ratio of Aspartic acid + asparagine to Glutamic acid + glutamine	Ratio of Dicarboxylic acids + amides to Serine + threonine	Ratio of Dicarboxylic acids + amides + tyrosine to Hydroxy acids + histidine + tryptophane + cysteine
<i>Enzymic proteins</i>			
Pepsin (ref. 6)	1.5	1.0	1.1
Chymotrypsinogen (ref. 6)	1.4	0.71	0.64
Ribonuclease (ref. 6)	1.2	1.0	1.1
Triosephosphate dehydrogenase (ref. 6)	2.9	1.0	0.86
Lysozyme (refs. 7, 8)	3.2 (4.7)	1.0 (1.5)	1.1 (0.83)
Aldolase (ref. 6)	0.94	1.2	1.1
Hæmoglobin (horse) (ref. 6)	1.3	1.5	1.0
Trypsin (ref. 9)	1.7	0.8	—
Mould protease (refs. 9, 10)	1.8	1.3	—
<i>Non-enzymic proteins</i>			
Insulin (ref. 6)	0.40	2.6	2.5
Edestin (ref. 6)	0.64	2.5	2.2
β -Lactoglobulin (ref. 6)	0.58	2.9	2.3
Myoglobin (horse) (ref. 6)	0.53	2.2	1.3
Myosin (ref. 6)	0.45	2.6	2.5
Tropomyosin (ref. 6)	0.30	4.3	4.5
Albumin (human) (ref. 6)	0.66	2.5	2.1
Gelatin (ref. 6)	0.58	2.4	2.3
Zeln (ref. 6)	0.19	2.3	0.5
Gliadin (ref. 6)	0.03	5.0	4.3
Casein (ref. 11)	0.35	2.0	1.9
Serum albumin (bovine) (ref. 12)	0.73	2.2	1.9
Human plasma mucoprotein (ref. 13)	0.77	2.0	1.8
Lactogenic hormone (ref. 14)	0.91	1.8	1.5
Ovalbumin (ref. 6)	0.62	1.6	1.4
Silk (ref. 6)	1.4	0.25	6.2
Wool (ref. 6)	0.52	1.0	1.0
<i>Cl. botulinis</i> toxin type A (ref. 6)	1.4	2.3	2.5
Fibrinogen (ref. 6)	1.0	1.7	1.4
γ -Globulin (ref. 6)	0.82	0.81	0.85
Conalbumin (ref. 8)	4.7	1.5	1.5
Ovomucoid (ref. 8)	2.2	1.65	1.6
Avidin (ref. 8)	1.62	0.9	0.75
Ratios apparently characterizing enzymes	>1 (Range 0.94–4.7)	Approx. 1 (Range 0.7–1.5)	Approx. 1 (Range 0.6–1.1)

Possible mode of hydrogen-bonding between serine and asparagine residues in adjacent polypeptide chains in (a) the keto form and (b) the extreme enol form. P = protein chain

to those which characterize the enzymic proteins. While more analyses, especially of enzyme proteins, are desirable before any final conclusions are drawn, we consider that the range of enzyme types covered by existing analyses is sufficiently broad to warrant an attempt to interpret the results. We regard these ratios as implying the existence in enzymic molecules of frequent interchain hydrogen-bonds, probably largely of the carboxyl-hydroxyl type, which reinforce the intra-chain hydrogen-bonding of peptide links to form a highly resonating network over which electron flow can occur. We think it is significant that, for many of the side-chains we have considered, it is possible to envisage hydrogen-bonding not only between the side-chains but also simultaneously between each side-chain and an adjacent peptide bond. An example of such a system involving residues of asparagine and serine is shown in the accompanying formulae. The side-chain of glutamic acid with its additional CH_2 group does not lend itself so readily to such a configuration, and this may explain the preponderance of aspartic acid in enzyme molecules shown in the first relationship in the table.

We hope to publish a fuller account of these and other implications in the near future. [Aug. 15.]

- ¹ Szent-Györgyi, A., *Science*, **93**, 609 (1941).
- ² Evans, M. G., and Gergely, J., *Biochim. et Biophys. Acta*, **3**, 188 (1949).
- ³ Geissman, T. A., *Quart. Rev. Biol.*, **24**, 309 (1949).
- ⁴ Pauling, L., Corey, R. B., and Branson, H. R., *Proc. U.S. Nat. Acad. Sci.*, **37**, 205 (1951).
- ⁵ Uzman, L. L., and Blout, E. R., *Nature*, **166**, 862 (1950).
- ⁶ Tristram, G. R., "Advances in Protein Chemistry", **5**, 83 (Acad. Press, New York, 1949).
- ⁷ Fromageot, C., Cold Spring Harbor Symp. Quant. Biol., **14**, 49 (1950).
- ⁸ Lewis, J. C., Snell, N. S., Heischmann, D. J., and Fraenkel-Conrat, H., *J. Biol. Chem.*, **186**, 23 (1950).
- ⁹ Authors' preliminary data from paper chromatograms.
- ¹⁰ Crewther, W. G., and Lennox, F. G., *Nature*, **165**, 680 (1950).
- ¹¹ Gordon, W. G., Semmett, W. F., Cable, R. S., and Morris, M., *J. Amer. Chem. Soc.*, **71**, 3293 (1949).
- ¹² Stein, W. H., and Moore, S., *J. Biol. Chem.*, **178**, 78 (1949).
- ¹³ Weimer, H. E., Mehl, H. W., and Wingler, R. J., *J. Biol. Chem.*, **185**, 561 (1950).
- ¹⁴ Li, C. H., *J. Biol. Chem.*, **178**, 459 (1949).

EFFECT OF THE ANNULAR ECLIPSE OF MARCH 7, 1951, ON RADIO-WAVE PROPAGATION*

DURING the annular eclipse on March 7, 1951, observations were made at Quartz Hill, Wellington, N.Z., to ascertain the effect of the eclipse on radio-wave propagation. The eclipse occurred at Quartz Hill at 1907 G.M.T., which was too soon after sunrise to obtain satisfactory discrimination of some of the effects, and this difficulty in discrimination was increased by the occurrence of a magnetic storm of moderate intensity which commenced on March 7, and lasted intermittently until March 14.

As was expected, medium-wave signal strengths increased to near night-time values and short-wave signals faded out as the umbra crossed the transmission paths. Along the line of the eclipse, observations were made of signals from the B.B.C. station GSD, 11.75 Mc/s., beamed on a London azimuth of 260° . On March 7, this station was not heard until 1940 G.M.T., but its signal strength then rose steadily to 1945 G.M.T., when it remained at a steady value some 20 db. below that received on the preceding days. The low signal strength and the fact that the station was not heard prior to 1940 G.M.T. may have been partially due to effects of the magnetic storm in northern latitudes. It was of interest that the fading characteristic of the station during the eclipse was identical with that obtained when the transmission path is in darkness.

Short-wave observations across the path of the eclipse were confined to the signals from station WVVH (10 Mc/s.) at Hawaii. The strength of signal from this station decreased at 1847 G.M.T., faded out at 1902 and returned suddenly at 1936 and then increased steadily to reach its usual strength at 1945 G.M.T.

Noise on the lower short-wave frequencies, 1.4–6 Mc/s., was higher, and noise on 30 Mc/s. was lower during the period of the eclipse as compared with the same period on the control days. The normal increase in noise on 30 Mc/s. following sunrise occurred some 90 min. later on the day of the eclipse, and a sudden drop of 4 db. in this noise occurred at 1920 G.M.T. some 13 min. after the

* Substance of a paper at the New Zealand Geophysical Conference by L. H. Martin, N.Z.B.S. Receiving Station, Quartz Hill, Wellington, N.Z.