CORRIGENDUM

Risk and prognostic factors for Japanese patients with chronic graft-versus-host disease after bone marrow transplantation

Y Atsuta, R Suzuki, K Yamamoto, S Terakura, H Iida, A Kohno, T Naoe, K Yano, A Wakita, H Taji, M Hamaguchi, Y Kodera, H Sao, Y Morishima, N Hamajima and Y Morishita

Bone Marrow Transplantation (2006) **38**, 583. doi:10.1038/sj.bmt.1705497

Correction to: *Bone Marrow Transplant* (2006) **37**: 289–296; advance online publication, 9 January 2006. doi:10.1038/sj.bmt.1705247

below. Eosinophilia (>4%) has been changed to Eosinophils (<4%) and the corresponding *N* has been changed to 57.

The authors have identified an error in Table 1 of the above paper; a corrected version of the table is given

Table 1 Prognostic significance of cGVHD parameters for cGVHD-specific survival

	N	HR (95% CI)	P
<i>Mode of presentation of cGVHD</i>			
De novo	33	1.0	
Quiescent	73	1.1 (0.44–2.6)	0.89
Progressive	4	1.8 (0.83–4.0)	0.14
<i>Grade of cGVHD</i>			
Limited	30	1.0	
Extensive	80	1.6 (0.60–4.3)	0.34
<i>Cutaneous extent</i>			
Code 1 (no skin involvement)	45	1.0	
Code 2 (50% or less of all skin surface area)	54	0.61 (0.26–1.5)	0.27
Code 3 (more than 50% of all skin surface area)	9	1.3 (0.71–2.2)	0.42
<i>Oral involvement</i>			
Code 1 (no clinical evidence of cGVHD)	25	1.0	
Code 2 (clinically evident oral symptoms)	78	0.62 (0.25–1.5)	0.31
Code 3 (severe oral cGVHD causing functional impairment)	4	1.1 (0.37–3.1)	0.91
<i>Thrombocytopenia</i>			
Code 1 ($\geq 100\,000/\text{mm}^3$)	56	1.0	
Code 2 ($< 100\,000/\text{mm}^3$ and $\geq 50\,000/\text{mm}^3$)	40	1.2 (0.48–2.8)	0.73
Code 3 ($< 50\,000/\text{mm}^3$)	14	2.1 (0.74–6.2)	0.16
Eosinophils (<4%)	57	3.5 (1.3–9.5)	0.01
Hyperbilirubinemia (>1.2 mg/dl)	22	1.7 (0.71–4.2)	0.23
Elevated alkaline phosphatase ($> 2.5 \times \text{ULN IU/L}$)	26	1.8 (0.79–4.2)	0.16
<i>Weight loss</i>			
None	61	1.0	
<10% from baseline; body weight before BMT	22	0.65 (0.22–2.0)	0.45
$\geq 10\%$ from baseline; body weight before BMT	9	1.4 (0.75–2.6)	0.29
Chronic diarrhea	7	2.1 (0.62–6.9)	0.24
Presence of infection	20	5.7 (2.5–13.2)	<0.0001
Reduced serum IgG (<500 mg/dl)	33	2.0 (0.84–4.7)	0.12
KPS <80%	30	5.8 (2.5–13.3)	<0.0001
Use of corticosteroids	23	3.9 (1.8–8.7)	0.001

Abbreviations: BMT = bone marrow transplantation; CI = confidence interval; cGVHD = chronic graft-versus-host disease; HR = hazard ratio; IgG = immunoglobulin G; KPS = Karnovsky performance score; ULN = upper limit of normal.