EFFECT OF FORMALIZED CULTURES OF *Cl. welchii* ON FRESH SHEEP RED CELLS

Type of <i>Cl. welchii</i>	Dilutions of cultures					
	1/16	1/32	1/64	1/128	1/256	1/512
A	+	—	—	—	—	—
B	+	+	+	+	+	—
C	+	—	—	—	—	—
D	+	+	+	+	—	—

+, Agglutination of red cells; —, no agglutination.

from agar showed a higher titre of agglutinins than those from fluid medium: supernatant fluids from broth cultures were almost inert.

As was the case with *Cl. septicum*, the reaction was inhibited by the appropriate immune sera. In order to avoid confusing non-specific agglutination of red cells by these sera, a sheep red cell + horse or rabbit immune serum system was used and gave clear-cut results. Serial two-fold dilutions of antisera were incubated with red cells for half an hour and were then completed with two hæmolytic doses of bacterial culture. Both antibacterial and antitoxic sera inhibited agglutination, but the latter were the more effective. It was possible to distinguish type A from types B, C and D in this way, but the overlap between B, C and D prevented any sharp differentiation of these groups.

Treatment of antitoxic and antibacterial sera with heavy suspensions of the corresponding organisms failed to remove group antibodies. The end-points in hæmagglutination were also unaffected when the antisera were added to red cells, previously treated with the homologous strain, in an attempt to remove anti-agglutinins.

These experiments, which confirmed in part those reported by Gürtürk³, show that the agglutinating factor in *Cl. welchii* cultures is associated with the bacterial cell but not with the toxin. As this factor is neutralized not only by the homologous antibody but also, except in the case of type A, by other antisera, the test can be used in identifying *Cl. welchii*, but not for typing the different strains.

E. N. DAFAALLA
M. A. SOLTYS

Department of Veterinary Pathology,
University of Liverpool.
Feb. 16.

¹ Dafaalla, E. N., and Soltys, M. A., *Brit. J. Exp. Path.*, **32**, 510 (1951).

² Wilsdon, A. J., *Second Rep. Inst. Anim. Path. Camb.*, 53 (1931).

³ Gürtürk, S., *Z. Hyg.*, **133**, 573 (1952).

Control of a Seed-borne Disease by *Chaetomium cochlioides* Pall., under Natural Conditions

MANY workers have reported striking antagonism between plant pathogens and saprophytes. Most of this work has been done in pure culture or in soil partially or completely sterilized before introduction of the test organisms. There are comparatively few records of disease control under natural conditions by the introduction of an antagonistic organism.

The work briefly reported below, based on earlier investigations in the Division of Plant Pathology, University of Minnesota, is an example of such control. It was then shown that when certain Brazilian isolates of *Chaetomium cochlioides* Pall. were introduced into unsterile soil, the severe blight caused by *Helminthosporium victoriae* M. and M. on oat seed-

lings was prevented. Similar results have now been obtained with a more active strain of *C. cochlioides* using oat seed heavily infested naturally with *Fusarium nivale* (Fr.) Ces.

Of several media used to increase inocula of the antagonist, oat-straw saturated with a 2 per cent glucose solution was most satisfactory. This was loosely packed into litre jars, sterilized and incubated for three to four weeks at 20° C. after inoculation. The straw became permeated with the fungus and covered with perithecia and was easily incorporated into soil in this condition.

A medium loam taken from plots under continuous cultivation was used with no treatment other than those given below. Oat seed heavily infested by *F. nivale* was obtained from the Norwegian Government Seed Testing Laboratory, Vollebakk, Norway. 4-inch pots were two-thirds filled with soil, and 20 gm. of the oat straw inoculum was spread over the surface on which were placed thirty seeds. These were finally covered with about 3 cm. of soil. The pots were kept in a cool frame at a temperature of 13–16° C.

In the following experiment, typical of a number, the treatments were: (1), (2) and (3) oat-straw cultures of three strains of *C. cochlioides*; (4) seed treated with an organo-mercurial fungicide; (5) uninoculated oat-straw. Six replicates were used, and the results obtained after five weeks are shown in the accompanying table.

CONTROL OF SEEDLING BLIGHT OF OATS

Treatment	Strain	Mean percentage emergence	Condition of the plants at 2/3 leaf stage		
			% Dead	% Blighted	% Healthy
1 Oat-straw culture	A	87.2	0	0	87.2
2 " " "	B	81.7	8.9	20.6	52.2
3 " " "	C	31.2	20.6	10.6	0
4 Organo-mercurial fungicide	—	88.3	0	3.9	84.4
5 Uninoculated oat-straw	—	26.1	17.8	8.3	0

The results show quite clearly the excellent control of disease obtained by strain A of the antagonist. This was as good as, if not slightly better than, that obtained by the proprietary seed-dressing. The importance of using suitable strains is demonstrated by the failure of strain C to produce any control and the intermediate results obtained with strain B. Results comparable to the above have been obtained with strain A using a variety of inoculation methods.

The mechanism of the control which is obtained has not yet been determined, although a certain amount of evidence has been accumulated which suggests that it is by the production and subsequent diffusion into the outer layers of the infested seed of one or more antibiotic substances. It is interesting to note, however, that the fungus is not strikingly antagonistic to *F. nivale* or other pathogens in pure culture. This aspect of the problem is being investigated, as is also the use of strain A in the control of other seed-borne and soil-borne pathogens, and together with the above will be reported in greater detail at a later date.

I wish to thank Dr. R. K. S. Wood for valuable help and criticism throughout these investigations.

M. TVEIT

Botany Department,
Imperial College of Science and Technology,
London, S.W.7.
Feb. 2.