(3) So far as one really knows, neither can be regenerated.

(4) The two types of genes are distinguishable because in sexual reproduction nuclear genes follow precise segregational patterns based on chromosomal mechanics while cytoplasmic genes do not.

(5) The criteria and the experimental means for the detection of plasmagenes require much more careful consideration than has usually been given them in the past. (This point has often been ignored by geneticists but is critically treated by Ephrussi on pp. 116-118 of his book.)

Second is Waddington's objection that bringing plasmagenes into play to explain differentiation leads to confusion. His argument is that "Either the plasmagenes arise *de novo* during development, in which case they are doing exactly that thing the impossibility of which is used as a means of detecting them; or they are all present, possibly in an inactive form, in the unfertilized egg, in which case they are nothing but our old and discredited friends the 'organ-forming substances' appearing under a new name and with their hair curled in the latest fashion". In all fairness, it should have been pointed out that any difficulties in the interpretation of differentiation that arise out of the second alternative apply to nuclear genes as well as plasmagenes.

Third, in the final paragraph of the review, Waddington states: "True, he [Ephrussi] starts off on the wrong foot, arguing rightly enough (on p. 4) that development involves a 'differential [which] must have its seat in the cytoplasm', but drawing the complete *non sequitur* that 'if the cytoplasm causes differentiation it must be endowed with the power of perpetuation of cell-type'". This statement of *non sequitur* is based on the quotation of two isolated fragments taken out of context, and appears to me to be a serious misrepresentation of Ephrussi's position. Actually, in the text, both before and between the two fragments quoted in the review, there is important material that establishes the general line of argument. Essentially this material is in reference to highly significant experiments in tissue culture, which show that differentiated cell types may breed true indefinitely. *Non sequitur* implies that a conclusion has been drawn fallaciously from irrelevant premises. Reference to the text clearly shows that Ephrussi has not been guilty of such an error, since his statement in question is not derived from the purported premise but is reasonable interpretation of a set of investigational observations that are under discussion.

ADRIAN SRB

Department of Plant Breeding,
Cornell University,
Ithaca, New York.

¹ *Nature*, 172, 784 (1953).

I WELCOME the opportunity to try to clarify a little further the difference of opinion between Prof. Srb and myself, which concerns rather the question of how the 'plasmagene' theory should be incorporated into general biological thought than the particular instance of Ephrussi's book, as to the high value of which we both agree, though it seems from different points of view.

The point I wished to make in my review was that many approaches to the 'plasmagene problem' seem

to me to take it back-to-front. Thus Srb himself suggests that we should start by defining plasmagenes; he gives four points which say in effect that they are just like nuclear genes except that they happen to lie in the cytoplasm; and it is only in his fifth and last that he raises the "criteria and the experimental means for the detection of plasmagenes", which to my mind is a demure way of hinting that quite possibly no such entities as have been defined exist at all. I believe that a more useful method of discussion would be to start from the empirical facts and proceed cautiously towards the definition.

Prof. Srb's second paragraph does not persuade me to change my opinion that it is more difficult to account for a process of change (of cell-type during differentiation) in the face of two sets of persistent entities (genes and plasmagenes) than of only one set. As to the two quotations referred to in his third paragraph, they are both taken from p. 4 of Ephrussi's book. The two sentences which separate them rather tend to reinforce than to contradict the possibility that the persistence of differentiated cell-types is due more to irreversible nuclear changes than to cytoplasmic determinants; and they certainly do not refer to the tissue culture evidence, which in any case does not prove the point one way or the other.

I am glad to reiterate the point with which I ended my review. The great merit of Ephrussi's book is that, starting by an exposition of the arguments which have led many people to suggest that the idea of plasmagenes gives us a simple explanation of development, he comes at the end to advocate the more cautious and empirical approach. His discussion, in the Addendum to the main body of the book, of the criteria for the recognition of plasmagenes (to which Srb refers) is particularly valuable; but the whole progression of his thought is, as I hope I sufficiently emphasized, of the greatest interest, and makes his book one of outstanding importance in the history of these questions, which are among the most fundamental with which biology is concerned at the present time.

C. H. WADDINGTON

Institute of Animal Genetics,
West Mains Road,
Edinburgh 9.
Feb. 15.

Are Inbred Strains Suitable for Bio-assay?

GRÜNEBERG¹ has directed attention to two properties of inbred strains which should counsel caution in their use for some experimental purposes: (1) They cannot be relied upon to remain constant in their heritable properties with the passage of time, and may diverge quite rapidly when split up into separate sub-strains and colonies. (2) They do not constitute phenotypically uniform material, but may, on the contrary, be strikingly more variable than the F_1 hybrids between strains.

This second proposition, for which Livesay² obtained experimental evidence in 1930, calls into question the established practice of using inbred material for the bio-assay of food factors, hormones, drugs, poisons, pathogens, etc. In such experiments precision in the estimates, and hence uniformity in the test organisms, is of the highest importance.

The characters measured in bio-assay are of the type usually termed 'physiological', whereas the