This investigation is in progress as an item in the study of fish parasites, and a detailed account may be awaited. We are grateful to Mr. M. Devidas Menon, assistant director of fisheries, Madras, for providing facilities, to Dr. G. Dubois, Neuchâtel, for valuable information, and to the Director of Industries and Commerce, and the Director of Animal Husbandry, Madras, for permission to publish these observations.

FLORENCE G. ABRAHAM
M. ANANTARAMAN

Freshwater Fisheries Research Station,
Madras 10,
and

Veterinary College, Madras 7. Sept. 2.

¹ Dawes, Ben, "Trematoda" (Camb. Univ. Press, 1946). Haderlie, E. C., *Univ. Calif. Pub. Zool.*, 57 (5), 303 (1953). Hughes, R. C., *Trans. Amer. Micro. Soc.*, 46, 248 (1927). Hunter, G. W., and Hunter, W. S., *J. Parasit.*, 20, 328 (1934).

² Hyman, L. H., "The Invertebrates", 2 (1951). Kaw, B. L., *Ind. J. Helm.*, 2 (2), 67 (1950).

³ Lal, M. B., *Nature*, 171, 130 (1953).

⁴ Chandler, A. C., *Amer. Midl. Nat.*, 45 (3), 711 (1951).

Molecular Orientation in the Wings of a Termite

THE orientation of complex molecules in certain tissues is well known¹. I have recently observed that, although the main structure of the wing of a termite (*Macrotermes* sp.) is isotropic, the two principal anterior veins (costal and sub-costal), running approximately parallel to the leading edge and to each other, show very marked anisotropy with the extinction direction parallel to the vein axis. These brown veins appear to contain both chitin and sclerotin.

X-ray photographs taken about the vein axis using copper K_{α} radiation lead to the idealized fibre diffraction patterns shown in Fig. 1 (a and b). In both cases the rings are barely visible except very near the sickles. The equatorial spacing of the

Fig. 1. (a) Original fibres; (b) treated fibre. Left: principal veins on wing, indicated by arrows; orientation corresponds to that of (a) and (b)

untreated vein, Fig. 1a, is 4.63 Å., the meridional one is 3.48 Å., with the sickles corresponding to not more than 7° disorientation. These spacings could correspond either to fully extended β -keratin-type chains running in the direction of the vein axis ($\pm 7^\circ$) with randomly orientated cross-linkages, or to orientated chitin molecules. It seems likely that both these sources are contributing to the total diffraction pattern.

The veins may be cleaned and bleached by boiling them in concentrated potassium hydroxide for about twenty minutes. If the bleached vein is washed with ethanol and water and air-dried, the diffraction pattern of Fig. 1b can be obtained. This shows that the meridional spacing is no longer present; but that another equatorial spacing has appeared at 4.13 Å., in addition to the former spacing at 4.63 Å. Hence the principal chain must be largely disintegrated by boiling with potassium hydroxide.

RALPH HULME

University College, Achimota,
Gold Coast. Sept. 14.

¹ Richards, A. G., "The Integument of Arthropods", 14ff. and 59ff. (Minnesota, 1951).

Photoelectric Cell Observations of Insects in Flight

RECENT work in this Laboratory involved the exposure of a red-sensitive caesium photoelectric cell to direct sunlight, the cell output being fed to a high-gain audio amplifier and a loud-speaker instead of the more usual d.c. amplifier-galvanometer chain. It was observed that occasional brief (< 0.25 sec.) random 'bursts' of sound occurred, of apparent frequency between 100 c./s. and 500 c./s. The number of 'bursts' in a given time varied considerably during the day, but appeared to rise to a peak in the late afternoon. No moving object which might immediately explain the phenomenon was visible to the naked eye, and the sound produced was quite different from the continuous 'rumble' produced when tenuous cloud drifted across the sun's disk.

The recording system was amended to use two photoelectric cells close together and feeding separate amplifiers. Observation of a number of 'bursts' showed that in many cases a burst recorded in one channel was followed after a discrete time-delay by a similar burst in the other channel. This suggested moving objects as the cause, and from the geometry of the system and the duration of a burst, established their speed as of the order of 5–10 m.p.h. and their range as relatively close.

Several of the bursts were recorded on magnetic tape and later played back, displayed on an oscilloscope, and their wave-forms photographed (p. 129). Fig. 1a shows a typical burst of duration about 0.15–0.2 sec., and Fig. 1b shows the same occurrence displayed on a faster sweep (150 m.sec.).

It is almost certain that these bursts of sound are caused by insects flying across the optical path between the sun's disk and the photoelectric cell, the generated frequency being due probably to the oscillation of the insect's wings, and the insects themselves being beyond the range of the unaided human eye or ear.

Several different types of captive insect, for example, bee, common housefly, have been exposed in a transparent container to the photoelectric cell and their wave-forms obtained. Fig. 1c shows the wave-form produced by a bee fluttering its wings, photographed at approximately the same speed of sweep as was Fig. 1b. It will be seen that there is a close