another study², where analogous observations were made in normal animals treated with hydrocortisone, biochemical investigation of the heart tissue confirmed the histological observation that deposition of calcium is associated with these lesions.

These incidental findings obtained with a potent hormonal agent emphasize that, when one has recourse to a hormone for one specific therapeutic effect, it is essential to bear in mind the tremendously far-reaching effects that may be exercised by that hormone on all tissues of the body.

This investigation was supported in part by grants from the Cancer Board of the University of California, and the Albert and Mary Lasker Foundation.

LOWELL L. SPARKS
WERNER ROSENAU
REX N. MACALPIN
THOMAS A. DAANE
CHOW HAO LI

Hormone Research Laboratory and
the Department of Anatomy,
University of California,
Berkeley, California.

¹ Daane, T. A., Sparks, L. L., Lyons, W. R., Li, C. H., Cole, R. D., and Hayashida, T., *Nature*, **173**, 1085 (1954).

² Lostroh, A., and Li, C. H., following communication.

Deposition induced by Hydrocortisone of Calcium in the Heart Tissue of Female C₃H Mice

It is reported in the previous communication that chronic administration of hydrocortisone acetate to tumour-bearing C₃H mice in high dosages produces degeneration of the cardiac muscle fibres. Histological examination of the heart tissue revealed deposition of a mineral material which stains positively with an alizarin dye, suggesting the presence of calcium. These observations have now been confirmed in acute experiments with normal female mice of the C₃H, Heston A and B-albino-C strains. Progressively higher values for the calcium content of the heart tissue, as determined by the spectrophotometric method, were noted in hydrocortisone-treated animals as the hormonal dosages were increased, paralleling the degree of mineral deposition observed grossly. These findings invite interesting speculation with regard to the similarity between what is noted here and the pathological changes seen in certain clinical conditions^{1,2}, although extreme caution must be employed in translating data from one species to another.

Virgin female mice (fifty days of age) of the C₃H strain were given daily subcutaneous injections of either saline or a saline suspension of hydrocortisone acetate (Merck), in a daily dose of 0.25–2.5 mgm. As a prophylactic measure, all animals were also injected intraperitoneally on the first two days with 1 mgm. of terramycin in 1 ml. of a 5 per cent glucose solution; despite this precaution, several of the treated mice succumbed to infection. Only those surviving the entire period of injection have been included in the final analysis.

On the day following the tenth injection, the animals were anaesthetized with nembutal; the heart, adrenals and thymus were removed and weighed. For the calcium determinations, the heart tissue from four or five animals was pooled in a porcelain crucible and ashed in a muffle furnace at 900° F. The ash was taken up in concentrated hydrochloric acid and

the calcium was precipitated as the oxalate³ at approximately pH 4. Following centrifugation, the precipitate was thoroughly washed with 2 per cent ammonium hydroxide, dissolved in 1 N hydrochloric acid and made up to a 5 ml. volume with distilled water. The calcium content was then estimated in a flame spectrophotometer at 554 mμ with appropriate standard solutions⁴.

Results are summarized in Table 1. As was to be expected, hydrocortisone caused thymic involution and adrenal atrophy; in addition, the hormone produced hypertrophy of the heart tissue. The spectrophotometric analysis of the myocardium afforded both identification of the mineral deposition as a calcium salt, and a means for quantitative estimation of the calcium content, which was shown to be a function of the hormonal dosage. While only small increases were obtained with 0.25 mgm. of the hormone, a ten-fold increase was evident at a dose-level of 1.0 mgm. The microgram quantities listed in the table for calcium should not be thought of in absolute terms, since some variation was found from week to week between appropriate groups given a standard amount of hydrocortisone for the 10-day period. Nevertheless, the comparative increases of calcium observable in the groups receiving progressively higher hormonal dosages were remarkably constant.

Table 1. THE EFFECT OF HYDROCORTISONE ACETATE ON ORGAN WEIGHTS AND CALCIUM DEPOSITION IN C₃H MICE

Daily dosage (mgm.)	No. of mice	Body weight at autopsy (gm.)	Adrenal weight (mgm.)	Thymus weight (mgm.)	Heart	
					Weight (mgm.)	Calcium content (μgm./100 mgm.)
0	10	19.5	4.0 ± 0.1*	41 ± 2.4	92 ± 3.6	2.2
0.25	7	17.0	3.0 ± 0.03	nil	124 ± 9.1	6.4
0.50	10	18.5	2.9 ± 0.06	nil	124 ± 3.9	17.2
1.0	5	18.0	3.1 ± 0.1	nil		25.8
2.5	5	17.0	2.8 ± 0.4	nil		27.9

* Mean ± S.E.

No conclusion can be drawn from the present results with regard to the amount of circulating steroid necessary to induce the observed pathological changes. When the hormone is administered subcutaneously as a saline suspension, considerable quantities of unabsorbed material accumulate beneath the skin. Moreover, doubling the dosage offers no assurance that the total absorption is increased proportionately.

Irrespective of these recognized variables in absorption, these results indicate clearly that the hydrocortisone administered is responsible for the relatively constant occurrence of calcific depositions in the right ventricle of the heart and for their incidental appearance in the liver and other viscera. The mechanism of its action and the influence of other endocrine factors remain to be investigated.

We wish to thank the Albert and Mary Lasker Foundation for research grants which have aided greatly in this investigation.

ARDIS LOSTROH
CHOW HAO LI

Hormone Research Laboratory,
University of California,
Berkeley, California.

¹ Albright, F., "Recent Prog. in Hormone Res.", **1**, 293 (1947).

² Albright, F., Harvey Lectures, 1942–43 (Sci. Press Printing Co., Lancaster, Pa., 1943).

³ Michaels, G. D., Anderson, C. T., Margen, S., and Kinsell, L. W., *J. Biol. Chem.*, **180**, 175 (1949).

⁴ Denson, J. R., *J. Biol. Chem.*, **209**, 233 (1954).