North Flugga, North Shetland, on September 5. This specimen had unfortunately been thrown overboard again; but the report from the fishermen with a detailed description of the turtle excludes any doubt about the species. The weight had been estimated at more than 300 kgm.

The three specimens in question most probably came in from the Atlantic north of Shetland, following the strong inflow of Atlantic water in this area.

So far as I know, there are no previous records of the leathery turtle from the northern North Sea. Single specimens of accidental stragglers have now and then visited French and British waters. A few have also appeared in the southern part of the North Sea, but these have more often been recorded dead or stranded on the coast. The present captures seem to be the more remarkable in that they have taken place at short intervals within a comparatively limited area and very far north. The finds may be taken to indicate that a more extensive movement of leathery turtles has taken place in the North Atlantic lately. This suggestion may perhaps be supported by new and supplementary records from other waters.

A more detailed account will be given in a Norwegian periodical.

Note added in proof. I have now obtained information about a fourth record of the leathery turtle off western Norway. It was observed by a tunny fisher on July 31, 17 miles west of Bulandet, county of Sogn og Fjordane (about 61° 18' N., 4° E.). This turtle was in good condition and was probably also an adult: estimated length, about 2 metres; weight, 4,500 kgm. Repeated attempts to catch it failed.

J. F. WILLGOHS

Zoological Museum,
University of Bergen. Oct. 15.

Serum Amylase Activity: Normal Values

In a paper¹ on the determination of amylase activity in biological fluids, we suggested a temporary working 'normal' range of 9-35 of our units per 100 ml. of serum. This working range was derived from a study of the serum amylase activity of 101 hospital patients in whom there was no reason for suspecting an abnormal serum amylase activity.

Fig. 1. Distribution curve of serum amylase activity in 100 'normal' healthy adult persons

We have now used our method¹ for determining the amylase activity of the serum of 100 'normal' healthy adult persons and the results have been examined statistically. The results range from 8 to 38 of our units, and the arithmetical mean is 19.47 of our units. 9 per cent of the results are less than 11 units, 8 per cent greater than 28 units; 3 per cent are less than 9 units and 2 per cent greater than 33 units.

Fig. 1 shows the distribution curve when the results are plotted in groups of three. The curve is skewed slightly to the right in the diagram.

The standard deviation is ± 6.84 and the coefficient of variation is ± 34.61 . Twice the standard deviation on each side of the mean should, theoretically², leave out about 5 per cent of the number of results. $19.47 \pm (2 \times 6.84)$ gives 5.79 and 33.15, and these limits cut off 2 per cent of the results, both of which are above the upper limit. There are no results below 5.79. Three times the standard deviation on each side of the mean gives the limits of the curve as -1.05 and $+39.99$.

We suggest, therefore, that the 'normal' range of results for our method¹ is 6-33 of our units per 100 ml. of serum.

Work is now being carried out on an ultra-micro modification of our present method¹ and the procedure will shortly be published elsewhere.

HAROLD V. STREET
JOHN R. CLOSE

Biochemistry Laboratory,
Department of Pathology,
Crumpsall Hospital,
Manchester 8.

¹ Street, H. V., and Close, J. R., *Clinica Chimica Acta*, 1, 256 (1956).
² Mainland, D., "The Treatment of Clinical and Laboratory Data" (Oliver and Boyd, London, 1938).

The Agama Lizard as Reservoir of *Salmonellae* in Nigeria

BACTERIOLOGICAL examination of the common agama lizard (*Agama agama*), which is plentiful in Ibadan, revealed that 18 of the 160 animals examined (11.2 per cent) were faecal carriers of *Salmonellae*. A variety of types were isolated, including *Salmonella elizabethville* (2), *S. kingston* (3), *S. durban* (1), *S. takoradi* (1), *S. kaapstad* (1), and a new type which has been assigned the provisional antigenic formula 4, 12 . . . i: 1, 6.

None of 100 house gekkoes (*Hemidactylus brookii*) were found to be carrying *Salmonellae*. To the best of our knowledge, lizards have not previously been recognized as a reservoir of *Salmonellae* in Nigeria, although Mackey¹ in East Africa, Lee and Mackennas² in Brisbane, and Hinshaw and McNeil³ in California have published accounts of lizards as carriers of *Salmonellae*.

We should like to thank Dr. Joan Taylor, director of the *Salmonella* Reference Laboratory, Colindale, London, for her help in identifying these organisms.

A detailed account of our findings will shortly be published elsewhere.

PATRICK COLLARD
DAVID MONTEFIORE

Department of Bacteriology,
University College,
Ibadan. Oct. 30.

¹ Mackey, J. P., *East Afr. Med. J.*, 31, 1 (1955).

² Lee, P. E., and Mackennas, I. M., *Bull. Hyg.*, 30, 1010 (1953).

³ Hinshaw, W. R., and McNeil, E., *J. Bact.*, 52, 349 (1946); 53, 715 (1947); 55, 870 (1948).