irregularities. The alternative view is that whiskers may well be imperfect, but high strength may still be observed because of the pinning of dislocations and inhibition of sources due to the small dimensions of the specimen. Both the latter strengthening mechanisms have been shown to be plausible for thin films from direct observation in the electron microscope, and it would be valuable to have more direct information of a similar nature about whiskers. Equally it would be valuable to have much more information

about the relation between strength and thickness for thin films.

It is important to know more about changes in mechanisms and modes of deformation with thickness and also whether a small thickness alone is adequate to achieve high strength. The structure of the film arising from the particular method of preparation may also turn out to be important.

J. E. GORDON
J. W. MENTER

CHEMISTRY OF PLANT PRODUCTS

THE Royal Australian Chemical Institute's Western Australian Branch, in co-operation with the University of Western Australia, held a symposium on "Plant Products" at the Chemistry Department of the University on March 20 and 21. The first day was devoted to general and technological papers and the second mainly to structural chemical investigations. About 45 members and guests attended the two sessions.

After introductory remarks by Dr. D. E. White, president of the Branch, who welcomed the visitors, Dr. J. R. Price (Division of Industrial Chemistry, Commonwealth Scientific and Industrial Research Organization, Melbourne) outlined the development of phytochemistry in Australia, from the pioneers, like the Surgeon of the First Fleet, to its present flourishing state. Work in this field is carried on in most of the Universities and also by the Commonwealth Scientific and Industrial Research Organization. After discussing briefly the role of these institutions, Dr. Price summarized the scope, organization, objectives and potentialities of current work in the Division of Industrial Chemistry. Mr. R. H. Pearce (Kwinana Chemical Co., Kalamunda) described the application of solvent extraction and chromatography to the separation of fractions from *Xanthorrhoea* resins. Variations with species and age of the resin were discussed and some components were found to be unstable in light and air. The techniques used appear much more promising than those previously used in attempts to isolate compounds from this resin.

An account of the use of growth regulators in viticulture was given by Mr. L. T. Jones (Department of Agriculture, Perth), who described the effect of hormone spraying in place of the obsolete and harmful practice of cincturing vines. Local research work illustrated the superiority of such hormones as PCPA, 2,4-dichloro- and 2,4,5-trichloro-phenoxy-acetic acids and gibberellic acid, resulting in increased production, improved fruiting, prevention of pre-harvest fruit drop and the killing of unwanted vines. Reference was made to the disadvantages of hormone spraying, for example, damage to bunches and vines, but these were due to incorrect application of the spray. Mr. Jones pointed out that gibberellic acid (not strictly a hormone), had been used experimentally only recently, and the results obtained suggest a promising future for the use of this material in viticulture. A striking contrast between the sizes of bunches on treated and untreated vines was shown in colour slides.

Mr. W. E. Hillis (Division of Forest Products, Commonwealth Scientific and Industrial Research

Organization, Melbourne) discussed chemical aspects of the sapwood-heartwood transformation, referring particularly to the increased polyphenolic content of the heartwoods of Australian timber trees. These components, which contribute to durability, may be derived from the cambium, the leaves or the starch in the sapwood, and evidence indicates that an appreciable portion arises from the latter. The significance of the shikimic acid content of *Eucalyptus sieberiana* leaves of different sizes, before and after shading, was discussed.

Dr. H. Jakobs (Leederville, W.A.) described how chemical research had made possible much of the agricultural development in the Dutch East Indies. He exemplified this in detail by reference to the tea industry, which is now of major importance in Indonesia.

Dr. D. E. White discussed a study made with D. K. Dougall on the 'primary' tars, obtained by distillation of jarrah and karri sawdust *in vacuo*. Among the compounds isolated catechol, 2:6-dimethoxyphenol, 1:2:3-trihydroxybenzene, 5-methylfurfural and D-glucosan (1:5) β (1:6) were identified. Decomposition of the latter was responsible for the low decomposition temperature of the tar.

The second day's proceedings opened with a paper on the chemistry of stock poison-plants by Dr. C. C. J. Culvenor (Division of Industrial Chemistry, Commonwealth Scientific and Industrial Research Organization, Melbourne) read by Dr. J. R. Price. The liver-damaging pyrrolizidine alkaloids and their N-oxides isolated from Compositae, Leguminosae and Boraginaceae were described with special reference to the recognition of monocrotaline from *Crotalaria retusa* as the cause of Kimberley horse disease. Recent work on other *Crotalaria* spp. was also discussed.

Dr. P. R. Jefferies (Chemistry Department, University of Western Australia) outlined the principles of gas chromatography and showed how it could be used for the quantitative analysis of simple mixtures. Complex mixtures as found in essential oils can be studied qualitatively, first on the whole oil, and then on distillation fractions. In this way the isolation of pure compounds is greatly facilitated and the presence of high-boiling components can be easily detected.

The constitution of the sesquiterpene alcohol globulol was discussed by Mr. G. J. H. Melrose in a joint paper with P. R. Jefferies and D. E. White. He described new sources of this alcohol in Western Australian Myrtaceae, and the synthesis of an epimer by the reaction of methyl magnesium iodide with apoaromadendrone. This alcohol also affords aromadendrene on pyrolysis of its 3:5-dinitro-

benzoate. Hence the junction of the 5 and 7 membered rings in aromadendrene must be the more stable (*cis*) type. Other features of the stereochemistry of globulol and *epi*-globulol were also discussed and evidence for the attachment of the cyclopropane ring to C₆ and C₇ was outlined.

The chemistry of *Lunasia* alkaloids was discussed by Dr. J. R. Price, who compared the American work on *L. amara* with that done in the Commonwealth Scientific and Industrial Research Organization on *L. quercifolia*. In addition to the tertiary bases, lunacrine and lunacridine, water-soluble bases were present, the major component of which analysed for C₁₇H₂₂O₃N⁺. A possible structure for this compound, which forms lunacridine when treated with alkali, was presented.

Mr. J. O. Knight presented the final paper in collaboration with A. R. H. Cole, D. C. Shaw and D. E. White, on triterpenoid constituents of the Myrtaceae. After discussing the common occurrence of betulinic acid in members of this family he dealt with the structure of melaleucic acid. The structure originally assigned (3 β -hydroxylup-20(29)-en-25:28-dioic acid; *Chem. and Indust.*, 926; 1956) cannot be correct since the lactone which was assumed to involve the 3-hydroxyl and the 25-carboxyl affords a tertiary hydroxyl group on reduction with lithium aluminium hydride. The second carboxyl group must be on C₂₆ or C₂₇ and its lactonization involves re-arrangement. Means for distinguishing between these possibilities were discussed.

D. E. WHITE

THE INSTITUTE OF BIOLOGY ANNUAL CONVERSAZIONE

THE annual conversazione of the London Branch of the Institute of Biology was held at University College on the evening of July 3. Prof. G. P. Wells and Dr. K. Mellanby (chairman of the London Branch) received members and their guests in the North Cloisters of the College. A large number of exhibits was on display in the Zoology Department, by courtesy of Prof. Wells.

The range of activities of this year's guest Society, the Botanical Society of the British Isles, was well shown by its exhibit. Of particular interest was its progress in the scheme to produce a map of the distribution of British plants. The mass of data for this scheme has been reduced to a minimum of space by the use of a punched-card system, and it is hoped that an atlas of British plants will be produced in 1960. A related project has been undertaken by the British Naturalists' Association, whose exhibit illustrated their work on a phenological survey which will map the changes in the flora and fauna which occur during the year and also the fluctuations of longer period.

Hospital and Laboratory Supplies, Ltd., provided this year's trade-stand, which gave a sample of the comprehensive range of apparatus and materials which this firm can supply. Specimen publications of the Universities Federation for Animal Welfare and of the Institute of Biology were on display, and the Department of Extra-Mural Studies of the University of London had produced an exhibit to illustrate the work of the Department in the field of biology.

Aspects of the Falklands Islands Dependencies Survey biological research were shown, including the method of ageing individual seals by the annual increments of dentine or cementum in the teeth, and also illustrating work on the reproductive cycle of the southern elephant seal. The Ministry of Agriculture, Fisheries and Food provided a stand showing advances in food research and development, particularly in food dehydration, and its Infestation Control Division was represented by Mr. G. A. Brett's exhibit of a common tropical cockroach, *Pycnoscelus surinamensis*, which is occasionally imported into Britain.

In addition to the exhibits of organizations, a large number of exhibits had been provided by members and guests. Studies on pest control provided the basis of several of these. An example of a serious infestation of a house by the ant *Lasius brunneus*

was found in 1957 (A. A. Green), and there is some evidence of an increased tendency for this ant to invade houses. Population studies of locusts (R. F. Chapman) have made it necessary to be able to separate the egg-pods of locusts from those of other species; various characteristics of the pod can be used. The way in which infestation in a bulk of grain causes heating (D. H. Burges) introduced a model showing the rate of development of colonies of the flat grain beetle (L. P. Leffkovitch), colonies of up to 550,000 individuals being produced in 4 months.

A more gradual example of population increase was shown by the frog *Rana ridibunda* (J. I. Menzies); 12 specimens were introduced into England in 1935, and their descendants have spread at the expense of the British *Rana temporaria* and now cover 120 square miles of S.E. England. New anatomical work on the classification of the lugworm genus *Arenicola*, which can be divided into two groups, was shown by Prof. G. P. Wells, with a map of the zoogeography of this genus. Two very different types of colour change in the Orthoptera were shown (K. H. L. Key), some acridids showing a physiological change from black to blue as a direct reaction to increasing temperature, while some phasmatids show centromorphic colour changes related to population density.

Some exhibits were concerned with genetical studies. Studies on finger-print patterns (S. B. Holt) have shown that the total ridge-count is determined by additive genes with an independent effect but without dominance. In the fungus *Aspergillus nidulans* (M. Mahoney and D. Wilkie) some spores give rise only to asexual colonies; this appears to be due to the loss of a self-replicating cytoplasmic unit. The rare genetically determined biochemical defect in man known as gargoylism (U. Mittwoch) can be detected by the presence of abnormal inclusions in the lymphocytes.

An exhibit of the different degrees of specificity of both the bacteria and the leguminous plant in their symbiotic relationship (P. S. Nutman) showed also the way in which different mutant strains make it possible to study how the relationship develops. New methods of studying the cellular constituents of yeast (A. A. Eddy) have resulted from the discovery that protoplasts can be prepared by treating yeast cells with certain digestive enzymes of the snail *Helix*.