The reduction rates of different dinitrophenylhydrazones proved to be different. The results obtained with a number of dinitrophenylhydrazones after development of a paper chromatogram are summarized in Table 1.

Table 1. RESULTS FOUND BY TITRATION OF DINITROPHENYLHYDRAZONES ON WHATMAN-1-PAPER WITH TITANOUS CHLORIDE: recoveries from 20.0 μ gm. carbonyl compound

2,4-Dinitrophenylhydrazone of	Time and temperature of reduction:		
	5 min. 20°C.	15 min. 20°C.	5 min. 50°C.
Formaldehyde	~16	17.0	20.6
Acetaldehyde		17.0	19.2
Propionaldehyde	~15	19.3	19.4
Acetone	~19	19.9	20.0
Butanone	~16	20.0	
Pentanone-2	16.4		21.2
Cyclohexanone	16.4	~18	19.3
Heptanone-2	20.6	~22.8	24.0*

* Reproducibility excellent

From these results it is clear that a reaction period of 5 minutes at 50°C. will, in general, suffice for the quantitative reduction of the nitro groups. We found that under these conditions the reproducibility is very satisfactory.

Difficulties due to the stability of the titanous chloride solution² were completely overcome by proper exclusion of air. In contrast with a statement in the literature² we found that light had no influence. A 0.003 *N* solution of titanous chloride in 0.6 *N* hydrochloric acid, stored in a colourless glass bottle did not show a decrease of titre for about one month.

The method described is not restricted to carbonyl compounds. Work is in progress to adapt the method to the analyses of alcohols (dinitro esters), phenols, amines, and amino-acids (dinitro aryl-derivatives). The quantitative reduction of many of these derivatives on paper is possible. The quantitative conversion of the parent compounds to the derivatives is being studied. In principle the method is applicable to all substances containing groups which are reduced by titanous chloride in a reproducible way.

A more detailed description of this work will be published elsewhere.

L. BLOM
J. CARIS

Centraal Laboratorium,
Staatsmijnen in Limburg,
Geleen, The Netherlands.

¹ Matthias, W., *Naturwiss.*, **43**, 351 (1956).

² Ma, T. S., and Early, J. V., *Microchim. Acta*, 130 (1959).

BIOCHEMISTRY

Distribution of 5-Carboxymethylhydantoinase

THE enzyme 5-carboxymethylhydantoinase, which reversibly cyclized carbamylaspartate to 5-carboxymethylhydantoin, was demonstrated in *Zymobacterium oroticum* by Lieberman and Kornberg¹. No enzymatic activity was found in rat liver or in two corynebacterium strains². In recent studies on pyrimidine metabolism in man this enzyme was found to be absent in erythrocytes and normal or leukæmic leukocytes^{3,4}. The absence of this enzyme in mammalian cells and its presence in bacteria suggested the possibility that chemotherapeutic agents might be developed as competitive inhibitors of 5-carboxymethylhydantoinase. Inhibition of the growth of *Lactobacillus casei* by thiohydantoin-5-acetic acid has been reported⁵. A survey of the distribution of the enzyme in micro-organisms was therefore undertaken.

dl-5-Carboxymethylhydantoin-¹⁴C was synthesized from orotate-¹⁴C as previously described³ with specific activity of 1.9×10^8 c.p.m./ μ mole. Enzyme assay was based on the hydrolysis of 5-carboxymethylhydantoin-¹⁴C (3×10^{-4} moles) to carbamylaspartate-¹⁴C during 1 hour incubation at 37°C., pH 8.2 (*tris*). Following the addition of 10 μ moles carbamylaspartate carrier and protein precipitation, carbamylaspartate was isolated by elution from a 'Dowex'-1-formate column (1 $\times$ 10 cm.) with sodium formate buffer pH 3.2 (50 column volumes wash using 0.02 *M*; 20 column volumes using 0.05 *M*). Specific activity was determined in duplicate tubes³ and the rate of synthesis calculated by the standard carrier formula. *Zymobacterium oroticum* was grown using a modification of the method of Friedmann and Vennesland⁶. Other bacteria were grown in standard laboratory media, harvested by centrifugation, and disrupted by sonication for 10 min. using a Raytheon 10 kc./s. Oscillator. By analogous reactions as for 5-carboxymethylhydantoin synthesis, 1-5-carboxyethylhydantoin was synthesized from 1-carbamyl aspartic acid: m.p. 164–165°C. (dec.) Calculated (per cent): C 41.86, H 4.68, N 16.28; found C 42.07, H 4.91, N 16.39; and 1-5-sulphonylmethylhydantoin from 1-carbamylcysteic acid: potassium salt m.p. 270–274°C. (dec.) Calculated (per cent): C 20.70, H 2.59, N 12.08, S 13.80; found C 20.75, H 2.35, N 12.27, S 13.52.

The results can be summarized briefly. The presence of 5-carboxymethylhydantoinase in *Zymobacterium oroticum* was readily confirmed with rates of synthesis of carbamylaspartate of approximately 13 μ moles/mgm. protein/hr. Trace enzymatic activities, approximately 1–2 per cent of that in *Zymobacterium oroticum*, were found in *Pseudomonas fluorescens*, *Proteus vulgaris*, and *Staphylococcus aureus*. No significant enzymatic activity could be detected in *Bacillus subtilis*, *Alcaligenes faecalis*, *Lactobacillus leichmannii*, *Escherichia coli* B, beta hæmolytic streptococcus, *Salmonella st. paul*, Brewer's yeast, or the Ehrlich ascites cell tumour. Neither of the structural analogues were an effective inhibitor of 5-carboxymethylhydantoinase from *Zymobacterium oroticum*. These results and those previously cited demonstrate that 5-carboxymethylhydantoinase has a very limited biological distribution. The metabolic significance of this spur reaction off the general pathway of pyrimidine synthesis has remained obscure, and its limited activity in pathogenic bacteria does not recommend it as a focus for chemotherapeutic attack.

This study has been supported by Grant No. A-1606 (C2) from the Institute of Arthritis and Metabolic Diseases, National Institutes of Health, Public Health Service. We are grateful to Dr. H. C. Friedmann for help in obtaining and culturing *Zymobacterium oroticum*.

LLOYD H. SMITH, jun.
FAITH A. BAKER
MARGARET SULLIVAN
LAWRENCE J. KUNZ

Departments of Medicine and Bacteriology,
Massachusetts General Hospital,
Boston, Mass.
July 28.

¹ Lieberman, I., and Kornberg, A., *J. Biol. Chem.*, **207**, 911 (1954).

² Reynolds, E. S., Lieberman, I., and Kornberg, A., *J. Bacteriol.*, **69**, 250 (1955).

³ Smith, L. H. jun., and Baker, F. A., *J. Clin. Invest.*, **38**, 798 (1959).

⁴ Smith, L. H. jun., and Baker, F. A., *Blood* (in the press).

⁵ Sós, J., Csálay, L., Kemény, T., Harnos, G., Perényi, L., Schnell, M., and Jóna, M., *Acta Physiol. Acad. Sci. Hung.*, **10**, 397 (1956).

⁶ Friedmann, H. C., and Vennesland, B., *J. Biol. Chem.*, **233**, 1398 (1958).