The results of observation on the incidence of the rhinoceros beetle and the extent of damage done by it under field conditions were presented by Mr. C. P. Ramachandran and Dr. C. Kurian (Kayangulam). Detailed studies of *Perisierola nephantidis*, an ectophagous larval parasite of *Nephantis serinopa*, were reported by Mr. J. Antony and Dr. C. Kurian. Mr. K. V. Joseph (Vellayani), based on the work done by him, emphasized the need to exercise caution in the matter of using DDT for large-scale

spraying in coconut areas because of the risk involves in adversely affecting the population of parasited and predators that normally help to keep pests under control in Nature.

Other papers were presented and discussed, and some were submitted only. All the papers submitted for the Conference are being printed as a special volume of the *Indian Coconut Journal* (published by the Indian Central Coconut Committee, Ernakulam, South India).
K. M. PANDALAI

MEDICAL RESEARCH IN THE BRITISH CARIBBEAN

THE fifth annual scientific conference organized by the Standing Advisory Committee for Medical Research in the British Caribbean was held at the University College of the West Indies, Jamaica, during April 2-4. The topic for the meeting was the application of the laboratory to problems of clinical medicine. Doctors from the Caribbean islands, British Guiana, British Honduras, the Bahamas, Great Britain and the United States attended. Thirty-five papers were read and discussed. Prof. A. C. Frazer (University of Birmingham) took the chair on the first day.

The first session was devoted to cardiovascular disorders. A number of hospital studies and field surveys were reported from Jamaica, Trinidad, British Guiana and St. Kitts, by Drs. J. A. Tulloch, G. Wattle, H. Hamilton and K. Stuart. Hypertensive heart disease was the most common cardiac cause for hospital admission followed by luetic and then rheumatic heart disease. Acute rheumatic fever was common, and in Trinidad affected the East Indian to a greater extent than the Negro. Cor pulmonale was uncommon in both Trinidad and Jamaica, but was seen with greater frequency in British Guiana. In British Guiana and Trinidad it occurred almost exclusively in the East Indian. No satisfactory explanation of this racial difference was forthcoming. Myocardial infarction was uncommon in all the territories. This was of interest as aortic atheroma is as common in Jamaica as in the United States of America, although coronary artery atheroma is less severe in Jamaica than in the United States. The association of aortic and coronary atheroma with essential hypertension, diabetes, age and sex was discussed by Dr. W. B. Robertson. The importance of heart disease in the general population was illustrated by the report from Barbados of a chest X-ray survey of 3,000 subjects. This showed a 10.7 per cent incidence of cardiovascular abnormality, and a considerable proportion of these had a positive serological test for syphilis. Yaws has not been recognized in Barbados for many years. This stressed the fact that symptomless cardiovascular syphilis may be more common than has been thought and therefore poses an important public health problem. In the discussion of this paper the urgent need for a test, serological or otherwise, that will differentiate syphilis from yaws was stressed.

Hypertension is common and field surveys carried out in Jamaica and St. Kitts were reported by Dr. K. L. Stuart. The levels of blood pressure in St. Kitts were slightly higher than those reported by the Medical Research Council team in South Wales, whereas in Jamaica the figures were comparable to

those from South Wales. An investigation into the occurrence of significant bacilluria in the general population was also reported by Dr. Stuart. There was a considerably higher incidence of bacilluria in the hypertensive female, and this increased with parity. The relationship of bacilluria to hypertension was discussed, and the possibility that pyelonephritis was responsible for the greater incidence of hypertension in the elderly female was suggested. A post-mortem study of hypertensive cases reported by Drs. Ling and Stuart did show that renal hypertension was responsible for 50 per cent of the cases, and that chronic pyelonephritis accounted for one-third of the renal cases.

Diabetes mellitus was discussed at the second session. It was pointed out by Dr. J. A. Tulloch that the prevalence of diabetes in Jamaica was a little more than 1 per cent of the adult population, and that only about 40 per cent of the cases were already known to have diabetes. The aetiology was reviewed and was considered to be very similar to that reported from temperate countries. A consideration of the J type cases was reported by Drs. D. MacIntosh and J. Tulloch, which indicated that this was not a distinct type, but was a phase in the natural history of the badly treated type II patient. The results of treatment with chlorpropamide were reported from Trinidad by Dr. T. Poon-King. A considerable proportion of his cases responded satisfactorily. The distribution of blood groups in diabetic subjects was also reported from Trinidad. In both male and female cases there was a low incidence in group O. Males showed an excessive group B and females an excess of group A over that occurring in the local population. A study of liver enzymes in human diabetes was reported by Dr. S. J. Patrick. This largely confirmed animal experiments.

On the second day Prof. E. T. C. Spooner (London School of Hygiene and Tropical Medicine) took the chair. Dr. R. D. Montgomery of the Medical Research Council Tropical Metabolism Unit reported that the oxygen uptake per unit of body weight was not reduced even in severe malnutrition. During recovery the increase became much greater and evidence was given that this was related to thyroid activity. Drs. J. C. Waterlow and D. I. M. Picou reported preliminary observations from isotope studies that, in protein malnutrition, the economy of protein is achieved not by a decreased rate of catabolism but by an increased degree of re-utilization. Although the extent of protein synthesis may be reduced through shortage of supplies, the capacity for synthesis seemed in most cases to be unimpaired. The occurrence of magnesium deficiency in malnutrition was reported

by Dr. R. D. Montgomery. This paralleled potassium deficiency and supported the concept of a primary loss of intra-cellular constituents. Further investigations into the megaloblastic anaemia noted in malnourished infants was reported by Drs. MacIver and Back. They showed that the cause was a dietary deficiency of folic acid, and vitamin B₁₂ was not implicated. Dr. MacIver also reported on the nature of the severe anaemic episodes in sickle-cell disease and showed that these were of two types. In one folic acid deficiency was the aetiological factor and in the other there was an acute aplastic state. A possible relationship of this latter state to preceding infection was suggested. Non-Addisonian vitamin B₁₂ deficiency in the adult was reported from Trinidad by Dr. G. Habib. All the cases were Hindus and with one exception they were strict vegetarians eating no eggs, meat, fowl or fish. It was considered that poverty and a vegetarian diet combined to produce this deficiency state.

A field survey on the size of liver in apparently healthy children was reported by Dr. C. G. Miller. Hepatomegaly was found in about 95 per cent of all children under the age of 12 years. Its presence was not related to malnutrition, but the possible role of infection was considered. The autopsy incidence of cirrhosis of the liver, as noted at the University College Hospital of the West Indies, was much the same as in America. One-third of all cirrhosis cases resulted from veno-occlusive disease of the liver. Liver cancer occurred with moderate frequency. This was a more common finding than in Denmark, but was much less common than in Africa. Hepatocellular carcinoma was more common than cholangio carcinoma and occurred particularly in the young adult male.

Virological investigation of human yellow fever cases in Trinidad was reported by Dr. W. G. Downs of the Trinidad Regional Virus Laboratory. The

cases noted in 1959 were diagnosed by isolation of virus from the blood and the difficulty of serological diagnosis was stressed. The clinical occurrence of encephalitis in Jamaica was analysed by Dr. R. A. Irvine and it was noted that sequelae were a common occurrence. The intra-dermal inoculation against Asian influenza was discussed by Drs. M. M. Sigel (Miami University) and L. S. Grant. In their opinion the intra-dermal injection of antigen would engender an antibody response and such a method could well be adopted in an emergency during epidemics before large supplies of suitable antigens became available.

Prof. G. Bras took the chair on the third day. The possibility that the unusual spastic neuropathies seen and reported from Jamaica were related to treponemal infection was considered by Dr. R. D. Montgomery (Medical Research Council). It was argued that the neuropathy could be a late manifestation of treponemal infection, whether venereal or otherwise, and that it could be due to a strain acting as a poor antigenic stimulus, but producing a typical meningo-vascular lesion in the nervous system. A second neurological disorder noted in Jamaica, that of spinal arachnoiditis, was reported by Prof. E. K. Cruickshank and Dr. P. E. Rodgers. The aetiology of the condition was largely unknown, but it had been found that cases seen early in the disease responded well to cortisone administration. An investigation into tropical plants used for medicinal purposes was reported by Prof. Haynes. It had been found that the breadfruit contained γ -aminobutyric acid, and this factor gave rise to a slight reduction of blood pressure in the anaesthetized dog. Other plants had also been shown to contain this substance.

The meeting was brought to a close by Prof. Bras who thanked all contributors and all visitors from the many islands, Great Britain and the United States for attending the conference.

J. A. TULLOCH

ABUNDANCES OF THE RARE-EARTH ELEMENTS, LANTHANUM TO LUTETIUM, IN CHONDRITIC METEORITES

By DR. R. A. SCHMITT, A. W. MOSEN, C. S. SUFFREDINI, J. E. LASCH,
DR. R. A. SHARP* and D. A. OLEHY

John Jay Hopkins Laboratory for Pure and Applied Science, General Atomic
Division of General Dynamics Corporation, San Diego, California

THE abundance distribution of nuclear species is considered an important requisite for a satisfactory theory of nucleogenesis¹⁻³. Compilations of cosmic abundances of the elements have been published by Goldschmidt, Brown, Urey, Aller, Suess and Urey¹, and Cameron², and others. It has been found that the relative cosmic abundances vary by a factor of $\sim 10^{13}$, and decrease logarithmically and rather smoothly from the lightest to the heaviest elements.

Rules of relative atomic abundances, for example, regular and smooth decrease in the abundances of those odd-mass-number nuclides which have masses greater than ~ 50 , were assumed to be valid for all elements, because it had been observed that the rare-earth elements (lanthanum to lutetium) showed

a regular and smooth decrease in their abundances¹. Since it is well known that the fourteen rare earths possess a high degree of chemical similarity, because of the filling of the 4-f electron shell far inside the valence electrons, the rare earths are assumed to have migrated geochemically through time and space as an 'entity'⁴. Consequently, accurate determinations of the abundances of rare earths in primordial matter are required in order to establish empirical relationships among the abundances of elements formed in nucleogenesis.

Noddack, using X-ray spectroscopic techniques⁵, has determined the abundances of the rare earths in composite mixtures of chondritic and achondritic meteorites (~ 9 parts of chondrite to 1 part of achondrite in the mixtures); and Minami⁶ has used

* Present address: Sharp Laboratories, Inc., La Jolla, California.