of concrete. J. De Yong, in a paper with H. G. Higgins, described draining studies on a pad of cellulose fibres, representing an early stage of paper formation. Compressibility of the porous medium complicates both the Kozeny and the drag equations. L. Dintenfuss presented a rheological classification of suspensions, separating the properties of the solid and the liquid phases. He correlated colloidal structural properties such as sedimentation volume with the flow-curves of paints and other systems.

In the following section, "Hygroplasticity", several papers were devoted to wood and one to wool. Dr. G. N. Christensen described the effect of rheological properties on sorption. He had observed that the rate of water sorption by thin layers of wood is inconsistent with diffusion control, and suggested it might be dependent on the relaxation of swelling stresses. Two important practical effects arising on drying wood, namely, 'collapse' and 'set', formed the subject of a paper by Dr. W. G. Kauman, who correlated macroscopic observations with the molecular processes in the cell wall. M. Feughelman discussed the effect of humidity on stress relaxation in wool, and suggested a model in which hydrogen bonds break in the presence of 'mobile' water. L. D. Armstrong presented experimental evidence on the great increase in creep of wood when changes in moisture content take place while the material is loaded. The influence of changes in moisture content was also observed by J. W. Gottstein, who used strips of wood veneer held at constant deformation and subjected to humidity cycles.

The fourth section was concerned with "Specific Materials". Dr. F. A. Blakey reviewed factors influencing the creep of gypsum plaster, and again emphasized the effect of free moisture content on the stress-strain-time relations. Various practical requirements of glazing compounds were discussed by E. R. Ballantyne, who compared the properties of oil-based and rubber-based compounds. C. Massey described tests on several thermoplastics with reference to their suitability for model structural analysis.

In a section on "Instruments", Dr. B. I. Aldrich, in a joint paper with E. L. Newman, described a Couette viscometer designed for testing syrup during fractionation; it accommodates a wide range of shear stresses and has close temperature control. K. G. Martin described a micro-viscometer and discussed factors in its operation.

"Continuum Theory" was the subject of the last section. Dr. P. U. A. Grossman discussed Prof. K. Weissenberg's ideas leading from an assumed restriction on stress directions for a given strain history to observable phenomena, such as cross-stresses.

A policy session was held, at which the possibility of future co-operation in the field of rheology was discussed. In New South Wales the branch of the Society of Rheology has already held several meetings and has arranged a full programme of activities for the current year, including lecture courses on rheology. Plans were discussed for broadening the interest in this field, and for holding future meetings both at the local and national level. P. U. A. GROSSMAN

GAS CHROMATOGRAPHY

THE Gas Chromatography Discussion Group of the Institute of Petroleum Hydrocarbon Research Group held its annual general meeting in conjunction with an informal symposium at the Brunel College of Technology, Acton, London, W.3, on April 22. The secretary, Mr. D. H. Desty, reported that the Group had had another successful year in which it consolidated its position following the rapid expansion of the past few years. Interest in the Group's activities has been well maintained and its status in the field is now well recognized on an international basis.

The abstracting service organized by the Group has made rapid progress, and more than 100 British and foreign journals are abstracted regularly by a team of 25 abstractors. The inaugural volume of "Gas Chromatography Abstracts", 1958, was published by Butterworths, London, on May 5, and has largely been the work of Mr. C. E. H. Knapman, the editor of the Group's editorial committee. This volume provides abstracts and, more particularly, a detailed subject index for 1,468 papers in this field published up to January 1, 1959. Succeeding volumes will appear annually.

Dr. A. T. James and Dr. D. Ambrose have taken part in the preparation by an international committee of preliminary recommendations on the nomenclature and presentation of gas chromatographic data, which were presented to the Analytical Section of the International Union of Pure and Applied Chemistry in September, 1959. These have now been approved in the International Union for publication. The Group's financial position is very sound, and its

financial year has been adjusted to coincide with that of the Institute of Petroleum Hydrocarbon Research Group (that is, to the end of September). No change in the annual fee (1 guinea), which is now primarily intended to cover the cost of publications issued by the Group, is therefore necessary.

The informal symposium which followed the annual general meeting was organized and introduced by Dr. J. A. Skollon of Brunel College of Technology, who commented on the active research interests of the College in the field of gas chromatography. The first paper, and perhaps the highlight of the meeting, was that by Dr. J. H. Purnell on the modification of solid supports. Dr. Purnell described measurements, made in his laboratory, of the adsorptive properties of 'Celite' and brick dust which show that whereas 'Celite' has the lower adsorption, brick dust has the more reproducible properties. These experiments were made with the view of studying treatments of the support which might reduce adsorption and minimize tailing of peaks. One promising method of treatment is the use of hexamethyldisilazane, which reacts with hydroxyl groups. Its practical application is simple and the treated support is unwettable and has reduced adsorption. Although chromatograms obtained on the treated support are improved, a further improvement is obtained if 0.1 per cent polyethylene glycol is added to the column with the stationary phase. One disturbing fact is that treatment of the support in this way has a slight effect on values of the specific retention volume even when the columns contain as much as 20 per cent of stationary phase.

The other contributions to the meeting concerned the study of pyrolysis products. Dr. J. Girling described studies on the identification of the primary products of the pyrolysis of coal, in which field the introduction of gas chromatography has enabled the scale of experiments to be reduced from 20 kgm. to 20 mgm. A 20-mgm. sample is heated to a series of steady temperatures ranging from 50° up to 700° C. and the products of pyrolysis at each temperature are swept into a refrigerated trap. The analytical sample is injected on to the column, containing polyethylene-glycol or squalane, by means of an automatic valve of 0.1 ml. capacity; there was considerable interest in the design of this valve, the fixed sample volume of which is formed by a length of 1-mm. bore nylon tubing between two sections of the tubing which are compressed by cam-operated plungers. Dr. J. Blenkin described the analysis of hydrocarbon elastomers by combined pyrolysis and gas chromatography. It was hoped that elastomers could be identified if the monomers from which they are made could be obtained from them and analysed by gas chromatography: to a large extent this hope has been realized. Samples are pyrolysed by dropping them into a heated silica tube attached to the apparatus, so that the products of pyrolysis are swept into the column and chromatographed; for example, polybutadiene gives

a chromatogram of butadiene, natural rubber one of a mixture of isoprene and dipentene. Dr. Lehrle described similar experiments in which the degradation was carried out in stages. A sample is placed in a coil of resistance wire in the form of a basket which is located at the entrance to the column. The temperature of the wire is varied by altering the current passing through it, and within half an hour a series of chromatograms can be obtained corresponding to the degradation of the one sample at each temperature. The series of chromatograms is characteristic and suitable for qualitative and quantitative analysis.

The symposium concluded with a general discussion on sample injection, which was introduced by Dr. Purnell. Most speakers were agreed on the difficulty, if not present impossibility, of injecting representative samples for quantitative analysis which are small enough to be satisfactory with ionization detectors. The lower limit of sample introduction possible with pipettes and syringes is probably about 25 micrograms. Various suggestions—for designs of valves or introduction of the sample in solution—were made, but the simplest proposal for quantitative analysis was made by Dr. A. B. Littlewood, namely, to use a large sample and correspondingly insensitive detector.

D. AMBROSE

MEDICAL RESEARCH IN MALAYA

THE valuable research work being done in the vast field of tropical medicine is well brought out in the annual report of the Institute for Medical Research, Federation of Malaya, for 1958.

The Institute is divided into a number of divisions, each of which is carrying out investigations into diseases which are baffling and wide-spread. In the Pathology Division, for example, a curious type of allergic granuloma, apparently hitherto undescribed and believed to be the result of an auto-immunity reaction, has been investigated for some years. The variations in the behaviour of melanocytes when surrounded by atypical or malignant epidermal cells have also been studied.

A number of important investigations are being made in the Division of Nutrition and Biochemistry. Severe anaemia is one of the main complications of pregnancy in Asian women, and one in eight of the beds available for ante-natal patients in the General Hospital, Kuala Lumpur, is occupied by a patient requiring treatment for anaemia. Premature births, deaths of infants and of mothers were all increased in this group of women.

The relationship of hookworm infection to anaemia was also studied: the greater the degree of anaemia, the higher was the proportion of patients with hookworm infection. The use of radioactive chromium enabled calculations of the daily loss of blood apparently due to hookworms; with heavy infections this could amount to as much as 80 ml. (almost 3 fl. oz.) of blood a day. Other studies were made on the changes in blood volume in chronic anaemia, on the factors influencing response to treatment, and on the reasons for the frequent low levels of vitamin B₁₂ to be found in patients with megaloblastic anaemia.

Dietary inquiries made in the homes of a number of the patients indicated that various causes, besides

poverty, were responsible for the poor diets which finally resulted in their admission to hospital with nutritional anaemia.

In almost every area where sufficient numbers of 'toddlers', children aged 1-4 years, have been examined, examples of protein malnutrition can be found; this is true even in the heart of Kuala Lumpur. Only a few come to hospital; they must be sought out by home visits and detailed inquiries. As might be expected, the condition is most common in isolated areas where communications are poor and unemployment is prevalent. A detailed investigation was made of kampongs on the Perak River where protein malnutrition is common; the diets there were found to be most unsatisfactory, and well below the recommended allowances.

In the Bacteriology and Serology Division, research activities were directed to the production of vaccines to combat epidemics of cholera and smallpox.

The search for Malayan streptomycetes which may produce antibiotics continued, but without success; resistance of bacteria to the commonly used antibiotics is becoming more noticeable.

Resistance to the tubercle bacillus is now reaching the point where it presents a serious problem. Judging by sensitivity tests, about 10 per cent of local strains are completely resistant to streptomycin, about 10 per cent are resistant to *p*-aminosalicylate, and 2 per cent to isoniazid. Partial resistance to one or other of these drugs is found in 30-50 per cent of strains tested.

The Virus Research and Medical Zoology Division continued its work on arthropod-borne and other viruses. Further work was done on the survey of animal reservoirs of leptospirosis, and infected rats were obtained from Maxwell Hill in Perak, from Langkawi Island, and from parts of Trengganu. By