Fig. 1. Glycogen synthetase activity. ●, Muscle; □, total embryo.

suggestions and the supply of eggs. This work was done during the visit of one of us (T. A. I. G.) to California.

T. ADESANYA IGE GRILLO
KENJI OZONE

Department of Anatomy,
University College,
Ibadan, Nigeria
and

Department of Poultry Husbandry,
University of California,
Berkeley, California.

¹ Grillo, T. Adesanya I., *J. Histochem. and Cytochem.*, **9**, 386 (1961).

² Bot, G. Y., Andrassy, K. O., and Kovacs, E. G., *Kiserl. orvostud* **12/4**, 380 (1960).

³ Leloir, L. F., and Goldemberg, S. H., *J. Biol. Chem.*, **235**, 919 (1960).

⁴ Dalton, A. J., *Anat. Rec.*, **68**, 393 (1937).

⁵ Lee, W. H., *Anat. Rec.*, **110**, 465 (1951).

⁶ Grillo, T. Adesanya I., *J. Endocrinol.*, **22**, 285 (1961).

Zinc in Bone

THE role of zinc in biological systems has been the subject of many investigations during recent years. Addink¹ suggested a possible correlation between the zinc content of blood and liver and the presence of cancer. No clear confirmation of this correlation has been obtained. Zinc has been found to be an integral portion of several enzymes by Vallee². The high levels of zinc present in the prostate, in leucocytes and in sperm have been observed by a number of investigators³. In general, the understanding of the role of zinc in biological systems is still largely reduced to a description of concentration differences rather than of metabolic interactions.

In a recent article Taylor⁴ infers that the zinc content of rat bone increases with age, for example, 77 p.p.m. to 136 p.p.m. to 200 p.p.m. at 56, 313 and 679 days respectively. If this is so, this finding would be quite useful in ageing studies and in the general understanding of bone structure. This suggestion prompted a re-examination of bones from several previous strontium studies for their zinc content.

The stable zinc-levels were determined by means of an X-ray fluorescence method and confirmed by emission spectrographic analysis. The satisfactory recovery of zinc under the conditions of ashing (12 h at 600°C) was verified using metabolically incorporated zinc-65.

In Table 1 are shown the results from a group of Wistar-strain rats reared on a diet which had a constant zinc to calcium ratio of 2.75 mg zinc/g

calcium. This experiment was terminated at 414 days, as compared with the 679 days of Taylor's experiment. No elevation of zinc in the bone was observed. In fact, since the ends contain approximately 30 per cent more zinc than the shaft, one might expect that the femur zinc-level would go down with age as the ratio of compact to cancellous bone increases, if it should be expected to change at all.

Table 1. ZINC CONTENT OF RAT FEMUR

Age (days)	Zinc (p.p.m. of ash)		Femur
	Shaft	End	
New-born	338	430	399
57	362	431	402
67	339	412	375
145	323	419	371
169	327	426	380
262	302	369	336
314	278	398	353
365	—	—	384
374	319	451	393
409	340	455	412
414	306	407	370
Mean $\pm$ S.D.			380 $\pm$ 24

In Table 2 are shown data from human ribs selected at random from two sets of samples previously analysed for strontium⁵. The results are tabulated according to increasing zinc-level in order to observe any possible correlation with age or locality. No such correlation is apparent, however, the six-fold increase in variance for the human samples, as compared with the rats on a controlled diet, suggests a definite correlation with the zinc and calcium ratio of the diet. It seems likely that the level of zinc in bone is directly related to the zinc to calcium ratio of the diet, as was found to be the case for strontium. The inferred retention factor for zinc in rat bone would be 0.40 as compared with 0.28 for strontium.

Table 2. ZINC CONTENT OF HUMAN RIB

Age (yr.)	Location*	p.p.m. Zn (ash)	Age (yr.)	Location*	p.p.m. Zn (ash)
40	F	106	14	F	155
16	B	122	31	A	155
15	C	123	35	C	170
22	A	131	17	B	171
16	G	134	13	E	180
9	B	140	75	B	188
17	A	141	47 $\pm$	B	202
5	B	143	38	G	209
34	B	144	51	B	300
47 $\pm$	B	147	0.7	B	306
47 $\pm$	B	149	63	B	330
30	D	150	38	G	372
11	D	153	Mean $\pm$ S.D. = 181 $\pm$ 70		

* A, Fresno and Oakland, California; B, Los Angeles, California; C, Phoenix, Arizona; D, Tulsa, Oklahoma; E, Madison, Wisconsin; F, Minneapolis, Minnesota; G, Philadelphia, Pennsylvania.

This work was supported by contract AT(04-1) GEN-12 between the U.S. Atomic Energy Commission and the University of California at Los Angeles.

GEO. V. ALEXANDER
R. E. NUSBAUM

Laboratory of Nuclear Medicine
Radiation Biology,
Department of Biophysics and
and Nuclear Medicine,
School of Medicine,
University of California,
Los Angeles.

¹ Addink, N. W. H., *Rec. Trav. Chim. Pays-Bas*, **70**, 168 (1951).

² Vallee, B. L., *Adv. Protein Chem.*, **10**, 317 (1955).

³ Koch, H. J., Smith, E. R., Shimp, N. F., and Connor, J., *Cancer*, **9**, 499 (1956). Mawson, C. A., and Fischer, M. I., *Nature*, **167**, 859 (1951). Wetterdal, B., *Acta Radiol.*, Supp. 156 (1958). Vallee, B. L., and Gibson, J. G., *J. Biol. Chem.*, **176**, 445 (1958).

⁴ Taylor, D. M., *Nature*, **189**, 932 (1961).

⁵ Hodges, R. M., et al., *J. Biol. Chem.*, **185**, 519 (1950). Alexander, G. V., and Nusbaum, R. E., *J. Biol. Chem.*, **234**, 418 (1959).