These participants, confirming the world-wide extension of the *Gravesia* zones, believe themselves faithful to the spirit of d'Orbigny when he created the stage in considering these zones as the base of the Portlandian. Four solutions have been proposed by R. Enay in his report on "The Tithonian Stage". In the course of the discussion a fifth, compromise proposition was formulated. The participants were inclined to agree either with Enay's fourth solution, or with the fifth.

Fourth Solution (R. Enay, 1962, "L'Etage Tithonique". Pre-print for the Colloquium on the Jurassic, Luxembourg, p. 29). "A last solution is to re-define the Portlandian in such a way that it begins with the *Gravesia* zones, a usage that has persisted for a very long time and still remains in use in numerous countries. Thus the Tithonian becomes the exact equivalent of the Portlandian, and at the same time its lower boundary coincides with that of the Russian Volgian. This is the solution that had already been adopted by E. Haug in 1898."

Fifth Solution: provisional solution agreed between and proposed by: K. W. Barthel, J. H. Callomon, R. Enay, A. J. Lloyd, A. Zeiss and B. Ziegler, August 3, 1962:

North-west Europe			Tethys		Russia
Stage	Cretaceous				
Portlandian			<i>Berriasella chaperi</i>	Tithonian	Volgian
	Sub-stages	<i>albani</i>			
Kimmeridgian	Upper (Dorsetian)	<i>pallasioides</i> <i>Gravesia</i>	<i>Gravesia (hybonotum)</i>		<i>Gravesia</i>
	Lower (Crussolian)	<i>autissiodorensis</i> <i>baylei</i>	<i>beckeri</i> <i>platynota</i>	Crussolian sub-stage	Crussolian sub-stage

In view of the impossibility of proposing a unanimous resolution, it was decided to refer the question back, for consultation among interested specialists.

(Luxemburg, August 4, 1962.)

In 1961, at the request of the International Mediterranean Mesozoic Committee, a British Mesozoic Committee was set up for the purposes of liaison with the international organizations, and to attempt to obtain a consensus of opinion among British Mesozoic specialists on matters of international significance. This Committee was recognized and approved by the British National Committee on Geology.

The British Mesozoic Committee met in February 1963 to consider the foregoing recommendations, which it approved but for the following points:

(1) *The Lower/Middle Jurassic junction.* The British Committee maintains that on both logical and historical grounds the Lower/Middle Jurassic junction should coincide with the Lias/Dogger junction, and should not come above the Aalenian as recommended at Luxembourg.

(2) *The Portlandian.* The British Committee approved the first of the two alternatives recommended by the Luxembourg Colloquium, namely, that the Portlandian should begin at the bottom of the *Gravesia* zones, following the first strict application of the term (by Salfeld) and making it the equivalent of the Volgian and the Tithonian. The Purbeckian is accepted as a facies term which should be dropped and the Portlandian should be regarded as ending at the top of the Jurassic.

(3) *Sub-stages.* The British Committee did not recommend the definition of named sub-stages.

(4) *Definition of stages.* The British Committee recommended, as a general principle, that the base of each stage should be regarded as fixed for all time (preferably by reference to a specified point in a type section). This would still allow for future adjustments, if needed, for example, to accommodate newly recognized strata, at the top of each stage. This principle would aid greatly the stability of stratigraphical nomenclature.

The Committee considered that other Mesozoic workers should be given this opportunity of considering these recommendations, and of communicating their views to the Secretary, before the Committee's views are passed on to a meeting of the International Committee in September 1963. It is hoped to have the Jurassic stages finally agreed before the next International Geological Congress in 1964.

DATA OF GEOCHEMISTRY

SIXTY-FIVE years ago F. W. Clarke, chief geochemist of the United States Geological Survey, prepared a critical summary of what was then known on the chemistry of the Earth's crust, the oceans, and the atmosphere. This monograph on *The Data of Geochemistry* rapidly became a widely used reference book and teaching text, and between 1908 and 1924 it was revised and enlarged four times. Since then, to meet continuing demand, the 1924 edition has been reprinted again and again, both in English and Russian translation; and as recently as 1960 it was referred to, in a widely used text-book of geochemistry, as "still the finest and most readily available collection of analytical data on geological material". Clarke retired in 1925 after 41 years' service, and died in 1931. His name has been commemorated by the Russians in the term used to denote the average content of an element in the Earth's crust.

Recognizing the need for an up-to-date revision of "Clarke", the United States Geological Survey some years ago invited geochemists in the United States and abroad to produce a collaborative sixth edition, each author preparing a chapter in his own field. The technical editor of this commendable enterprise is Dr. Michael Fleischer. It was recently announced that the new *Data of Geochemistry, Sixth Edition*, would comprise 36 chapters, each to be issued separately as a section of the United States Geological Survey *Professional Paper*

No. 440 (published by U.S. Government Printing Office, Washington, D.C.). Six of these have just been published and it looks as if the full work, to appear over several years, will amount to about 1,500 quarto pages.

The six chapters published comprise contributions on: the chemical composition of sub-surface waters, by D. E. White, J. D. Hem and G. A. Waring (Chapter F); chemical composition of rivers and lakes, by D. A. Livingstone (Chapter G); volcanic emanations, by D. E. White and G. A. Waring (Chapter K); chemical composition of sandstones, by F. J. Pettijohn (Chapter S); non-detrital siliceous sediments, by E. R. Cressman (Chapter T); and marine evaporites, by F. H. Stewart (Chapter Y). Each consists of text, lengthy tables of analyses, extensive references, and index; and to encourage further research the authors in each case take pains to emphasize the principal gaps in our knowledge. Each is, in effect, a text-book in itself, drawing on the literature in all languages. Prof. F. H. Stewart of the University of Edinburgh, who has produced an outstanding summary of what is known about marine evaporites, is as yet the only non-American contributor.

The review copies do not state the price. With all publications of the United States Geological Survey this is nominal, being based on the cost of printing an overrun for public sale after producing the edition required for approved repository libraries. C. F. DAVIDSON