occurring in the caecum of the rat and rabbit and in the rumen of the sheep. Experiments are now in progress with the view of identifying the organisms responsible and so far we have found that they are not inhibited by sulphanilamide but are inhibited by 'Dimol'.

A. N. BOOTH*
R. T. WILLIAMS

Department of Biochemistry,
St. Mary's Hospital Medical School,
London, W.2.

* On leave of absence from the Western Regional Research Laboratory (U.S. Department of Agriculture), Albany, California.

¹ Booth, A. N., Murray, C. W., Jones, F. T., and DeEds, F., *J. Biol. Chem.*, **223**, 251 (1956).

² Scheline, R. R., Williams, R. T., and Wit, J. G., *Nature*, **188**, 849 (1960).

³ Smith, A. A., *Biochim. Biophys. Acta*, **69**, 203 (1963).

⁴ Booth, A. N., Emerson, O. H., Jones, F. T., and DeEds, F., *J. Biol. Chem.*, **229**, 51 (1957).

⁵ Shaw, K. N. F., Gutenstein, M., and Jepson, J. B., *Proc. Fifth Intern. Cong. Biochem.*, Moscow, 369 (1961).

⁶ Benjafield, J. G., *Lancet*, ii, 782 (1962).

⁷ Scott, T. W., Ward, P. F. V., and Dawson, R. M. C., *Biochem. J.*, **87**, 3P (1963).

Mobilization of Fat Body Glycogen by an Extract of Corpus Cardiacum

HUMORAL involvement in the carbohydrate metabolism of *Periplaneta americana* (L.) has been reported by Steele¹, who demonstrated a 150 per cent increase in blood trehalose and a drop in blood glucose following injection of an aqueous extract of the corpus cardiacum. The work recorded here, conducted as part of an overall investigation of the regulation of sugar utilization in the South American cockroach, *Blaberus discoidalis* (Serville), confirms the rise in trehalose, but reveals also a significant increase in blood glucose and a concomitant decrease in fat body glycogen.

Gland extracts were prepared after Steele's method and injected intra-abdominally in 50 μ l. of distilled water. Blood sugar determinations (trehalose and glucose) were made on individual cockroaches by the method of Friedman². For this purpose 1 μ l. of blood was removed from an antenna, thoroughly chilled to prevent coagulation, diluted, deproteinized, centrifuged, and the supernatant made up to volume for the subsequent analyses. Glycogen obtained from the fat body was determined by the method of Good, Kramer and Somogyi³.

Results in Table 1 establish the significant elevation of both trehalose and glucose produced by the injection of the equivalent of 0.1 pair of corpora cardiaca (CC). An extract of corpus allatum (CA) prepared as the equivalent of one pair per injection is ineffective at the 5 per cent level. The lack of activity of the corpus allatum is a happy circumstance, since analysis of its function in any process in this animal will be complicated by anatomical arrangements which make it impossible to obtain the gland completely free of corpus cardiacum.

It is clear from the results in columns 1 and 2 of Table 1 that an increase in free trehalose does not come about at the expense of free blood glucose, since this is also raised. There is, however, some indication of the possible origin of

both these compounds. Column 3 of Table 1 contains the results of analyses of fat body glycogen made 5 h after injecting the named substances, and it may be seen that there is a substantial decrease in glycogen within the period in which the blood sugars are increasing. This information, coupled with the knowledge that in the locust⁴, at least, trehalose is synthesized from glucose in this organ, leads to the reasonable hypothesis that corpus cardiacum extract exerts its influence on glycogen synthesis and/or degradation in fat body. The regulatory capacity of the extract is, therefore, at present under investigation. It may be noted in passing that the effect is not mediated through other cephalic glandular material, since the extract is active on injection into the body cavity when the head of the animal is tied off.

This work was partially supported by David Ross pre-doctoral grant No. 2258 from Purdue University to one of us (W. S. B.).

W. S. BOWERS*
S. FRIEDMAN

Department of Entomology,
Purdue University,
West Lafayette, Indiana.

* Present address: Entomology Research Division, Insect Physiology Laboratory, U.S. Department of Agriculture, Beltsville, Maryland.

¹ Steele, J. E., *Nature*, **192**, 680 (1961).

² Friedman, S., *Arch. Biochem. Biophys.*, **87**, 252 (1960).

³ Good, C. A., Kramer, H., and Somogyi, M., *J. Biol. Chem.*, **100**, 485 (1933).

⁴ Candy, D. J., and Kilby, B. A., *Biochem. J.*, **94**, 19, P (1960).

L-Phenylalanine: an Organ Specific, Stereo-specific Inhibitor of Human Intestinal Alkaline Phosphatase

HUMAN alkaline phosphatase of intestine but not of liver, bile, bone, lung, kidney and spleen is almost completely (78 per cent) inhibited by L-phenylalanine and is not so affected by D-phenylalanine.

Specimens of fresh human tissue taken at autopsy were each processed for alkaline phosphatase as described earlier¹. These partially purified enzyme preparations were assayed in the presence of 0.005 M D-phenylalanine and of 0.005 M L-phenylalanine (Calbiochem, Los Angeles).

Table 1. SPECIFICITY OF INHIBITION BY PHENYLALANINE (μ g of phosphorus liberated from β -glycerophosphate* in 2 h)

	No phenyl- alanine	0.005 M D-phenylalanine	0.005 M L-phenylalanine	% Inhibition D-L $\times$ 100
Intestine	2.9 18.0 5.35 5.95	3.05 20.25 5.15 6.15	0.8 4.6 1.1 1.25	74 77 79 79
Gastro- intestinal washings	11.7 12.5	14.6 14.6	4.1 4.0	72 67
Liver	4.1 2.85	3.7 3.0	3.35 2.55	9 15
Bile	4.45 7.5 5.95	4.2 10.15 7.6	3.75 9.4 7.25	10 8 5
Rib	1.3	1.5	1.3	13
Vertebra	1.0	1.2	1.3	0
Vertebra	2.3	2.25	2.25	0
Kidney	1.5	1.25	1.1	12
Lung	1.45	1.45	1.2	17
Lung	2.0	2.2	1.9	13
Spleen	7.8 7.25 4.35 4.74	7.45 7.5 4.2 4.6	6.85 7.15 4.2 4.15	8 5 0 10

* Assays were done according to Fishman, Green and Inglis procedure (ref. 1), modified from Shinowara, Jones and Reinhart (ref. 2).

Table 1 gives the results obtained with a number of human tissues, from which results it is clear that only alkaline phosphatase of the intestine (as well as washings of the stomach and intestine) is strongly inhibited by L-phenylalanine as compared with D-phenylalanine, which is not an inhibitor of any preparation tested. There is, therefore, a double specificity, one dependent on the organ source of alkaline phosphatase and the other exclusively the property of one stereoisomeric form of phenylalanine.

Table 1. EFFECT OF INJECTION OF EXTRACTS ON BLOOD SUGAR AND FAT BODY GLYCOGEN

Injection	Mean changes in blood sugar concentration over 5 h post-injection period (mg/ml.)	Mean value of fat body glycogen 5 h after injection (μ g/mg dry-weight)	
	Trehalose	Glucose	
Water (4)*	-0.7	2.0	118.4
None (4)	-1.0 ($P = > 0.10$)†	1.4 ($P = > 0.10$)	98.9 ($P = > 0.10$)‡
CA extract (8)	2.3 ($P = > 0.05$)	1.5 ($P = > 0.10$)	80.4 ($P = > 0.05$)
CC extract (8)	23.7 ($P = < 0.002$)	7.6 ($P = < 0.02$)	45.9 ($P = < 0.001$)

* No. of individuals.

† Significance of change compared with change incurred on injection of water ('Student's' t -test).

‡ Significance of difference of mean from that of water-injected animals ('Student's' t -test).