Further information on future developments is given in No. 17 of *The B.B.C. Record* (February 1963), which emphasizes the value of the two television services. Moreover, it outlines the first year's proposals for adult education in television, and in the Sixth Bishop Bell Memorial Lecture, "Future Prospects in Broadcasting", delivered by H. Carleton Greene, the director-general of the Corporation, on April 30, and now published as a pamphlet; Mr. Greene stresses the complementary functions of the Corporation and the importance of concerning itself with the whole of the national life, with mass inter-

ests and with minority interests. He suggests that if broadcasting planners do their jobs properly they can create and stimulate cultural and educational demands as well as meet them. He directs attention to the use of television programmes in teaching and to the introduction of experimental programmes for use in technical colleges and colleges of further education. Furthermore, he urges that it would not be right to make an educational channel a charge on licence revenue: it should receive a separate grant-in-aid, as do the External Services.

SOILS AND LAND USE IN NORTH WALES

A SURVEY of the soils of North Wales was started by the late Prof. G. W. Robinson and his colleagues, who published several reports and maps in the *Welsh Journal of Agriculture*. The work has been completed recently with a memoir, and accompanying maps, which covers an area of 240 sq. miles, stretching from Penrhaeth, in Eastern Anglesey, and Llandudno in the north to Waenfafrw and Bettws-y-Coed in the south*. The Conway valley and most of the high land of Snowdonia are included.

The climate is maritime and its most important influence in soil development is the rainfall, which increases steeply from some 30 in. on the coast to 120 in. on the mountains in the south. The parent materials of the soils include acid and basic igneous rocks, schist and shale and their derived drifts, non-calcareous clay and limestone, peat and various mixed drifts from igneous and sedimentary rocks. The soil mapping and classification follow the custom of the Soil Survey and profile description includes information on site, drainage, parent material, horizons, colour, texture and structure. Soils of five major soil groups have been recognized, namely, brown earths, calcareous, gley, podzolized and organic soils, but there is also a considerable area of immature soils occurring on youthful sands and alluvium. Each is described and sub-divided into sub-groups and soil series, and again

* Agricultural Research Council. Memoirs of the Soil Survey of Great Britain—England and Wales. *The Soils and Land Use of the District Around Bangor and Beaumaris*. (Sheets 94 and 106.) By D. F. Ball. Pp. ix + 182 + 13 plates. (London: H.M.S.O., 1963.) 35s. net.

grouped into lowland, marginal and mountain soil zones.

The larger part of the memoir is naturally taken up with detailed descriptions of the soil series and with appendixes on particular profiles and analytical data, but there is a chapter on land use, compiled with the assistance of G. Edwards, H. E. Roberts and W. A. Lindsay-Smith. This opens with a historical review of the area from the time of the disappearance of the valley glaciers, with inferences to be drawn from studies on pollen grains and dating by radioactive carbon, on changes in climate, flora and human settlement. Recent agricultural statistics show the variation in size of holdings and the changes in crops and livestock between 1935 and 1955. An interesting aspect is the continuous rise in sheep production, but there has also been a pattern of technical improvement and change in farm economy to meet the seasonal fluctuations in demand. This is discussed in relation to the five regions of the area. There are about 12,000 acres of woodland, about half planted by the Forestry Commission—largely conifer high forest. Many factors, of course, determine tree growth, but it seems clear that goshawk oak and elm are to be found on freely drained slopes of soils derived from mixed drift although falling short of conifers in yield production.

Numerous figures and tables are of great assistance in assimilating the information presented, and several excellent photographs illustrate the outstanding features of the area in question.

A. M. SMITH

COMPARATIVE BIOCHEMISTRY OF THE LEGUMINOSAE

A SYMPOSIUM on "The Comparative Biochemistry of the Leguminosae", organized by the Plant Phenolics Group, was held at the John Innes Institute, Bayfordbury, Hertford, during September 10–11. The aim of the meeting was to bring together workers in diverse scientific fields but sharing a common interest in this group of plants. The Leguminosae, which contains some 550 genera and 12,000 species, is, in economic importance, second only to the Gramineae, and members of the family contain a remarkably large number of interesting and unusual chemical constituents. In his opening paper, N. W. Simmonds (John Innes Institute) reviewed the systematics of the family and summarized the many uses legumes are put to as foods, forages, timbers, gums, resins and tannins. Dr. R. Cooper (Rothamsted Experimental Station), in dealing with root nodulation, pointed out that 90 per cent of the fixed nitrogen in the world had come from leguminous root nodules. His review of the conditions required for successful nodulation by *Rhizobium* showed that many factors, including the occurrence of the correct host-symbiont specificities and the presence of the right kind of polysaccharide in the host roots, were involved.

Chemistry may well be, in the future, of considerable value to the systematists dealing with legumes. Its present contribution is slight, mainly because very few taxa have been thoroughly surveyed chemically. Dr. R. E. Alston and his colleagues in the University of Texas are therefore concentrating their efforts on a single genus, *Baptisia*. Dr. Alston illustrated his talk on this genus with pictures of the phenolic spots obtained on two dimensional chromatograms of leaf and flower extracts. The patterns so obtained can be used not only to distinguish most of the eighteen or so species but also many of the interspecific hybrids. Compounds which, at the moment, show more promise as taxonomic markers than the phenols are the free amino-acids. Besides having the common protein amino-acids, legume seeds contain a range of unusual ninhydrin positive substances, such as canavanine, homoarginine, γ -hydroxyhomoarginine and lathyrine. Dr. A. E. Bell (King's College, London), in the next paper, showed that most species of *Lathyrus* could be divided into four groups, depending on their characteristic amino-acid patterns; species of *Vicia* could be divided in a similar manner.