strated within the lysosomes in cryostat sections of normal tissues which have been suitably treated.

In a work of this kind, which is presumably intended to encourage workers in histochemistry, it is disappointing to find that the authors regard the subject in such a narrow fashion. It would appear that their initial approach is purely morphological, and that is certainly no disadvantage. When it is stated that "deriving its theoretical foundations from chemistry, histochemistry remains a morphological tool" (p. 342) one feels that the extensive possibilities of the subject remain unexplored by the authors. This is decidedly surprising for much attention has been directed to the demonstration of enzymes which are essentially dynamic in nature. Again on p. 342 they say that "finite measurement is not the immediate goal of microscopic histochemistry". The large number of biochemical comparisons made in recent years between normal and abnormal tissues have served to emphasize that while qualitative differences do exist they are greatly outnumbered by quantitative variations. It is hard to see how these differences can be identified if quantitative histochemistry is not developed. Biochemistry has done much to advance our knowledge of the cell, but a precise localization of chemical components can be achieved only by histochemical methods. Whether the method of study be the demonstration of carbohydrates, lipids, proteins, or enzymes, or the rate of uptake of labelled metabolites, the aim is the same, namely, to piece together our information on the metabolism of the cell. At the present time it is too much to hope that one discipline will provide the answer to any one problem, but it is quite certain that histochemistry in its future and fuller development will play a vital part in the understanding of cellular physiology and pathology. It would indeed be sad if it merely demonstrated a number of new structures, the function of which could not be determined.

GEORGE J. CUNNINGHAM

VIRUSES

Advances in Virus Research

Vol. 9. Edited by Kenneth M. Smith and Max A. Lauffer. Pp. viii+312. (New York and London: Academic Press Inc., Publishers, 1962.) 92s.

ONE-THIRD of this number of *Advances in Virus Research* is occupied by an article by St. Groth called "The Neutralization of Viruses". The article is in fact more limited than this for it deals almost exclusively with the influenza viruses. This makes it rather specialized, but it is written in a lively and colourful manner so that the reader is carried along in the stream in spite of possible disagreement with many of the *obiter dicta*. The influenza viruses are an admirable vehicle for carrying the general argument for they have an unusual number of measurable activities—infectivity in several hosts, toxicity, an enzymic activity, and haemagglutination—all of which are inhibited by suitable antibodies. St. Groth discusses the peculiarities and limitations of each type of measurement, the effects on them of virus and antibody heterogeneity, and, when possible, the mechanism of the action. Wisely, each section is followed by a summary so that the reader unfamiliar with work of this type can keep track of what he has been told.

The scope of the article by Porterfield on the arthropod-borne viruses, that is the viruses that can multiply in both mammalian and arthropod hosts as distinct from those that are transmitted mechanically by arthropod hosts, is somewhat similar. The article is factual rather than interpretative; it shows the complexity of the antigenic structure of the viruses and the variation in the hosts' responses to them. Non-specialists would have understood the article more fully if the editors had explained the terminology, especially the system of units used. It

ends with the comforting conclusion that there is no good evidence for the appearance of new viruses in this group though the antigenic components may regroup in suitable conditions.

The causative agents in a group of clinically rather ill-defined infections of the respiratory tract, and of some organs in the abdomen and elsewhere, in man and animals are grouped together as the adenoviruses. Electronmicrograms of several members of the group show icosahedrons with rounded corners. Brandon and McClean discuss both the structure of the virus particles and the reactions of the infected host to them. The story is fascinatingly complex with a cytopathic protein made by the infected host, at least three antigens that seem to be fragments of the intact virus and, in long-established infections, intranuclear crystals that seem to be virus aggregates. As the authors point out, this multiplicity of effects makes the group peculiarly suitable for an investigation of the general nature of virus infection.

The article by Smith is more general in scope. He gives a great deal of information on what might be called the natural history of the arthropod viruses and shows that many of the old ideas about them are ill-founded. They are by no means as host-specific as used to be thought, several can multiply in both insects and plants and cause pathological changes in both hosts, and latency is so widespread as to render all claims for the chemical induction of new virus infections suspect. The last phenomenon is particularly interesting because an uncongenial environment, or even a change of diet, can make the latent virus patent. There seem to be no general differences in morphology or chemical composition between arthropod viruses and those of plants or mammals, but the arthropod host seems able to tolerate an unusually large amount of virus before it finally succumbs—the only analogies with this extensive replacement of host tissue by virus is found in some of the infective mammalian warts, and in nitrogen-deficient plants.

Markham contributes a cheerful article on the information that can be gained about viruses by ultracentrifugation. He stresses that the instrument is easy to use, the calculations easy to make, and that a great deal can be found out with even one run on unpurified material. There is much shrewd and entertaining comment here. Those who were engaged in the early work on tobacco mosaic virus assumed that linear aggregation would have little effect on sedimentation constant; they will therefore be surprised to learn that the fact that this is so is unexpected.

Ten years ago, few virus workers could tell a regular from a rhombic dodecahedron; but we are all Platonists nowadays and the icosahedron is our juju. The significance of the regular solids (or the helix) is, however, widely misunderstood. These structures are likely to be common because they offer unusual opportunities for the formation of a stable shell. Speaking anthropomorphically, for the sake of brevity, the two main problems confronting a virus are: to find a system in a host that can be deluded into synthesizing virus, and to be stable enough to avoid the host's scavenging systems. We may therefore expect certain structures to appear often, as caffeine appears in at least 14 families of plants and the benzene ring (probably) in all organisms, though it serves many purposes and is made in many different ways. That is a reaction to the final article, by Andrewes, on the classification of viruses. We classify from a point of view and with an end in view. At one point Andrewes agrees and says "taxonomy is an artificial business"; he seems nevertheless to be searching for a classification that will command the general agreement that Linnean classification gets. But satisfactory Linnean classification summarizes the course of evolution, and there is no reason to assume that viruses, except within closely related groups, have undergone a similar process.

N. W. PIRIE