LETTERS TO THE EDITOR

PHYSICS

Ultra-violet Gas Laser at Room Temperature

ULTRA-VIOLET coherent light has been generated directly at room temperature in a pulsed nitrogen-gas laser. A group of 20 strong simultaneous lines have been observed in the ultra-violet. Thirty lines have been identified. These range in the ultra-violet spectrum from 3000 Å to 4000 Å. The strongest line has been observed at 3371 Å; the strongest visible line is at 3400 Å. The output radiation is believed to be produced as the result of inversion in the triplet state of nitrogen, resulting in subsequent emission in the second positive group, $C^3\pi_u \rightarrow B^3\pi_g$. The beam has a full angular width of 1 milliradian, as measured 100 ft. from the confocal laser cavity.

The plasma is excited with a 100–150-kV pulse of sub-μsec duration. The coherent-light pulse width is as short as 20 nsec. Despite the short pulse duration the output spot of the laser is exceedingly brilliant. The output spot for the confocal Fabry–Perot cavity does not contain the usual mode structure associated with single line operation, a characteristic that is noted when several sets of transitions compete simultaneously.

Output powers of the order of 10-W peak have been obtained summing over the lines. The plasma tube has a 48-in. discharge-length. Cold cathode excitation is used. The silver-coated flint glass end mirrors are spaced at 150 cm. The radius of curvature of the mirrors is approximately 400 cm. One mirror is heavily silvered, whereas the other transmits 2–5 per cent.

The infra-red nitrogen band spectra¹, which can be produced with 40-kV, 30-amp pulses of 1 μsec duration, are not observed when high-voltage pulses are applied.

H. G. HEARD

Energy Systems Inc.,
Palo Alto, California.

Mathias, L., and Parker, J. T., *J. App. Phys. Letters*, 3, 16 (1963).

Exchange Currents at Liquid Electrodes

RATES of non-equilibrium processes cannot be calculated *a priori* without quantitative hypotheses as to energies (heats) of activation. I emphasize this truism for electrochemical interface phenomena because ions are field particles in the sense that the potential barrier, over which they pass between the bound state and the solvated state, is itself changed by their passage. Only in thermodynamic equilibrium do ions become mere test particles, and it is perhaps for this reason that reversible thermodynamics does not always appear fruitful in this field.

In a recent communication¹ a model of the potential barrier, separating ions of a solid metal from solution, has been put forward and compared with known observations. I now reconstruct that model for interfaces between liquid metals and polar (aqueous) electrolytes. The basis of the model is retained: the height of the potential barrier (activation energy for dissolution) is the corresponding height, V_0 , with respect to *vacuo* (the real potential) divided by the dielectric constant, $\epsilon(\omega)$, of the electrolyte taken at the vibration frequency, ω , of the bound ions in accordance with Debye's theory² of the dispersion of the dielectric constant. Thus the electrolyte is viewed as a continuous dielectric medium with regard to the weakening of electrostatic interactions, but as a set of molecular dipoles with regard to its temperature-dependent response to the local electrical perturbations caused by the oscillations of ions bound at the interface.

The barrier has, therefore, a different height for each vibration frequency, the probability of dissolution per ion and unit time being again¹ $(\omega/2\pi)\exp[-V_0/\epsilon(\omega)kT]$. We thus need to know the distribution of bound surface ions over frequencies, and for this purpose the specific heat model of Debye³ has been adapted¹ for solid electrodes. Its counterpart for the vibrations of liquid metal surfaces has been given by Frenkel⁴, who considered surface capillary waves rather than elastic volume waves and obtained a cut-off frequency by regarding twice the mean particle separation as the shortest admissible wavelength, corresponding to a maximum wave number $K = K_{\max}$. The dispersion of capillary waves is given by:

$$\omega_s^2 = 2\pi K^{3/2}(\sigma/\rho)^{1/2} \quad (1)$$

σ being the surface tension and ρ the density of the metal. The number of vibration modes per unit surface area (which may be equated to three times the surface concentration of ions, in the manner of Debye³) between K and $K + dK$ is $2\pi K dK$, so that the absolute dissolution rate is:

$$R = \frac{2\pi(\sigma/\rho)^{1/2}}{3} \int_0^{K_{\max}} \exp[-V_0/\epsilon(K)kT] K^{5/2} dK \quad (2)$$

Wave numbers in the region of $K_{\max}$ correspond by equation (1) to frequencies larger than the inverse Debye relaxation time $1/\tau$ of water molecules. At such frequencies the orientational part of the polarizability vanishes and the activation energy becomes prohibitive at room temperatures (about $V_0/3$). It is for this reason that a good first approximation for R is obtained by putting $K_{\max} \sim \infty$ together with $V_0/\epsilon \approx (V_0/\epsilon_s)(1 + \omega^2\tau^2)$, where $\epsilon_s \approx 80$ is the static dielectric constant of a dilute aqueous solution:

$$R \approx \frac{\Gamma(7/6)}{9(2\pi)^{7/3}} \left(\frac{\rho^2}{\tau^2 \sigma^2} \right)^{1/3} \frac{\exp(-V_0/\epsilon_s kT)}{(V_0/\epsilon_s kT)^{7/6}} \quad (3)$$

$\Gamma(7/6) \approx 0.93$ being a Γ -function. The correspondence between equations (2) or (3) and the corresponding expressions for solid metals¹ may be brought out by comparing Frenkel's cut-off frequency with Debye's.

When the liquid electrode comes to equilibrium with a solution of its ions, the surface tension σ is connected with the electrode potential according to Lippmann's equation⁵ (note that R is the smallest at the potential of the electrocapillary maximum). We may then identify⁶ zeR (ze is the ionic charge) with the exchange current density obtained by extrapolating Tafel's overvoltage relations to zero net current⁵. It can be shown that Hg^{++} ions are dissolved in monovalent form ($V_0 \approx 6.5$ eV from a well-known cycle) and are ionized further by a fast redox step, and if we disregard the slight electrocapillary effect of the Hg^{++} concentration in solution on σ , equation (3) yields some 1.4 amp cm^{-2} at room temperature (not counting the electron current; if we count only one degree of freedom per surface ion, this result is to be multiplied by 3). This agrees with the observation⁵ that the exchange current density should exceed 0.4 amp cm^{-2} .

On the other hand, from the expression for R of solid metals¹ we obtain for Cu^+ at room temperature the exchange current density of 3.8×10^{-5} amp cm^{-2} as compared with the observed⁵ value 2×10^{-5} amp cm^{-2} . Considering the complexity of the physical situation, the calculated values for the liquid as well as the solid electrodes agree well with measurements. The two appropriate frequency distributions being very different from