It is noteworthy that normal nutritional or metabolic reactions produce a series of virogenic and mutagenic substances.

KAZUO YAMAFUJI

Agricultural Chemistry Institute,
Kyushu University,
Fukuoka, Japan.

KENJI WATANABE

Pharmaceutical Institute,
Fukuoka University,
Fukuoka, Japan.

- ¹ Yamafuji, K., and Shirozu, Y., *Biochem. Z.*, **317**, 94 (1944).
- ² Yamafuji, K., and Yoshihara, F., *Enzymologia*, **14**, 124 (1950).
- ³ Yamafuji, K., and Hirayama, K., *Enzymologia*, **17**, 229 (1955).
- ⁴ Yamafuji, K., and Yoshihara, F., *Enzymologia*, **22**, 89, (1960).
- ⁵ Yamafuji, K., and Omura, H., *Enzymologia*, **14**, 120 (1950).
- ⁶ Freese, E., Bautz-Freese, E., and Bautz, E., *J. Mol. Biol.*, **3**, 133 (1961).
- ⁷ Gierer, A., and Mundry, K. W., *Nature*, **182**, 1457 (1958).
- ⁸ Wyss, O., Stone, W. S., and Clark, J. B., *J. Bact.*, **54**, 767 (1947).
- ⁹ Jones, L. W., and Scott, A. W., *J. Amer. Chem. Soc.*, **46**, 2172 (1924).
- ¹⁰ Yamafuji, K., *Nature*, **167**, 770 (1951).
- ¹¹ Yamafuji, K., *Nature*, **171**, 745 (1953).
- ¹² Yamafuji, K., and Osajima, Y., *Nature*, **190**, 534 (1961).
- ¹³ Yamafuji, K., Osajima, Y., and Omura, H., *Nature*, **185**, 162 (1960).
- ¹⁴ Yamafuji, K., and Cho, T., *J. Agric. Chem. Soc. Japan*, **20**, 441 (1944).
- ¹⁵ Yamafuji, K., *Agric. Biochem. Kyushu Univ.*, **19**, 5 (1959).
- ¹⁶ Yamafuji, K., and Akita, T., *Enzymologia*, **15**, 14 (1951).
- ¹⁷ Yamafuji, K., Tonokawa, M., and Yoshihara, F., *Enzymologia*, **23**, 111 (1961).
- ¹⁸ Yamafuji, K., *Agric. Biochem. Kyushu Univ.*, **23**, 13 (1963).

Biosynthesis of L-Serine Ethanolamine Phosphodiester

FROM recent work¹ it would seem that L-serine ethanolamine phosphodiester (L-SEP) is found only in the tissues of fish, amphibians, reptiles and birds. D-SEP, on the other hand, appears to be confined to earthworms², where it is the biological precursor of lombricine³. The status of the work on both isomers has been summarized by Ennor and Rosenberg⁴. These authors had previously shown¹ that dinitrophenol (3×10^{-4} M) completely inhibited the biosynthesis of L-SEP and that cytidine triphosphate (CTP) appeared to be required. Thus it was shown that chicken kidney homogenates, supplemented by CTP, incorporated more inorganic phosphorus-32 into L-SEP than when this nucleotide was omitted.

This communication is concerned with an extension of this observation and with the development of a theory to account for the biosynthesis of L-SEP.

Crude chicken kidney homogenates have been used throughout and were prepared by homogenizing 10 g of fresh kidney in 20 ml. of ice-cold 0.25 M sucrose containing EDTA (10^{-3} M) in a Potter-Elvehjem type homogenizer for 30 sec. The homogenate was centrifuged at 500g for 10 min at 0° and the supernatant was used without further treatment. Incubation mixtures containing 1.0 ml. of the supernatant were incubated for 1 h in phosphate buffer pH 7.2 at 37° in the presence of various compounds labelled with carbon-14 and in a total volume of 2.0 ml. (Table 1). At the end of the incubation period trichloroacetic acid was added; a crude phospholipid fraction was prepared from the precipitated material and L-SEP was isolated from the supernatant by previously described chromato-

graphic techniques⁵. The radioactivities of the phospholipid fraction and of L-SEP were determined in a Packard 'Tricarb' scintillation counter.

The results (Table 1) indicate that appreciable incorporation of ¹⁴C-ethanolamine into both phospholipid and L-SEP was achieved only when CDP-¹⁴C-ethanolamine was the isotopic source and that this incorporation was enzyme dependent. It should also be noted that neither ¹⁴C-ethanolamine nor ¹⁴C-ethanolamine phosphate was incorporated into the phospholipids or L-SEP and that no isotopic dilution occurred when unlabelled ethanolamine phosphate was added together with CDP-¹⁴C-ethanolamine.

These results suggest that the final step in the biosynthesis of L-SEP involves the reaction:

which is presumably catalysed by a specific enzyme. Work now in progress is designed to determine the existence of this reaction and the results, together with full details of the present work, will be submitted for publication elsewhere.

We thank Dr. K. P. Strickland, of the Department of Biochemistry, University of Western Ontario, for a gift of CDP-¹⁴C-ethanolamine.

H. ROSENBERG
A. H. ENNOR
SADAKO SUGAI *

Department of Biochemistry,
John Curtin School of Medical Research,
Australian National University,
Canberra.

* Present address: Department of Biochemistry, Tohoku University, School of Medicine, Sendai, Japan.

¹ Rosenberg, H., and Ennor, A. H., *J. Biochem. (Tokyo)*, **50**, 81 (1961).

² Rosenberg, H., and Ennor, A. H., *Biochem. J.*, **73**, 521 (1959).

³ Rossiter, R. J., Gaffney, T., Rosenberg, H., and Ennor, A. H., *Nature*, **185**, 383 (1960).

⁴ Ennor, A. H., and Rosenberg, H., in *Amino Acid Pools*, edit. by Holden, J., 187 (Elsevier Publishing Co., Amsterdam and New York, 1963).

⁵ Kennedy, E. P., *J. Biol. Chem.*, **222**, 185 (1956).

⁶ Gaffney, T. J., Rossiter, R. J., Rosenberg, H., and Ennor, A. H., *Biochim. Biophys. Acta*, **42**, 218 (1960).

PHYSIOLOGY

A Biologically Active Metabolite of Vitamin A and Vitamin A Acid

A RECENT report by Zachman and Olson¹ described the formation of acidic water-soluble metabolites from vitamin A (retinol) and vitamin A acid (retinoic acid) by perfused rat liver. We have isolated an acidic metabolite, with potent vitamin A-like activity, from both retinol and retinoic acid. No compound metabolically related to retinol or retinoic acid with growth-promoting activity has hitherto been described.

Circumstantial evidence has for long suggested² that retinol is changed into an 'active form' in the course of its metabolic function. When the vitamin A-like activity of retinoic acid was discovered by Dowling and Wald³, this acid was at first considered to be the 'active form'. However, it could not be detected in the rat, even after administration of it to deficient animals which were being cured⁴. Apparently it was extremely rapidly metabolized. Therefore, the thought arose that retinoic acid, too, is changed into the actual metabolically active compound. The metabolite reported here may be closer than retinoic acid to the active compound, or may be the active compound itself. Whether it arises naturally from retinol via retinoic acid, or independently from either, has not been determined.

Retinol-deficient rats were given retinol (1 mg in 0.2 ml. cottonseed (Wesson) oil per day) orally by intubation or retinoic acid (1 mg in 0.2 ml. aqueous saline, pH 8, per

Table 1. INCORPORATION OF ¹⁴C-ETHANOLAMINE INTO L-SEP AND PHOSPHOLIPID IN CHICKEN KIDNEY HOMOGENATES

Additions	% of added c.p.m. recovered in L-SEP	% of added c.p.m. recovered in phospholipid
(1) CDP- ¹⁴ C-ethanolamine (a)	8.1	22
(b)	8.2	7.4
(c)	9.5	30
(2) As (1a), enzyme boiled	0.4	2.0
(3) As (1a) + unlabelled ethanolamine phosphate	7.7	38
(4) ¹⁴ C-ethanolamine phosphate	0	0
(5) ¹⁴ C-ethanolamine	0.1	0.5
	0.4	2.0

The labelled compound added in each experiment had a radioactivity of approx. 5,000 c.p.m.