Table 3. EFFECTS OF HOMOLOGOUS CHALLENGE OF ANIMALS NO LONGER INFECTED WITH "R" STRAIN *B. bigemina* (AFTER 1 TO 6 PRIOR EXPOSURES)

No. of previous homologous exposures	1	2	3	4	5	6	Total _s
Pathological effects from challenge		2			1		3
Parasitaemia with clinical effects							
Parasitaemia without clinical effects		1	1	4			6
No parasitaemia					1	1	2
Carrier status after challenge					1		1
Carrier two weeks after challenge			1			1	2
No test for carrier two weeks after challenge		3		4	1		8

Table 4. EFFECTS OF HETEROLOGOUS CHALLENGE OF ANIMALS NO LONGER INFECTED WITH EITHER "Y" OR "R" STRAIN *B. bigemina*

No. of previous homologous exposures	1	2	3	4	5	6	7	Totals
No. challenged	3	1	6	2	1	4	1	18
Parasitaemia with clinical effects	3	1	6	2	1	3	1	17
Parasitaemia without clinical effects						1		1

did not diminish thereafter. Resistance to homologous challenge with the "Y" strain was sufficient to prevent detectable parasitaemia in a total of 21 instances. Further investigations (by sub-inoculation and heterologous challenge) of 15 of these showed that in 7 the animal was not carrying *B. bigemina* two weeks after its homologous challenge. In other words, the homologous challenge had failed to set up a new infection. Results were similar but not so spectacular for the "R" strain.

From Table 4 it can be seen that there was a completely different response to heterologous challenge. These animals reacted severely, and several died despite treatment. The reactions resembled those of the splenectomized controls.

Hence, a sterile immunity has been demonstrated in an infection which has always been presented as a classic example of "prémunition"⁴ (co-infectious immunity). It must be pointed out, however, that it is a limited sterile immunity in that it is only effective when the animal has been exposed previously to the challenging strain, and even then not 100 per cent of the animals possess it. This sterile immunity does not exist when a heterologous challenge is performed.

The results also indicate an immunological difference between the two strains of *Babesia* used. A more detailed account of these investigations, the extension of the work to *Babesia argentina* and the epidemiological significance of the findings will be presented elsewhere⁵.

This work was begun in the Department of Parasitology, University of Queensland, and continued at the Animal Research Institute, Brisbane, where I am now employed.

L. L. CALLOW*

Department of Parasitology,
University of Queensland,
St. Lucia, Brisbane.

* Present address: Animal Research Institute, Yeerongpilly, Queensland

¹ Tallaferrro, W. H., *Fourth Int. Congr. trop. Med., Washington, D.C., 1, 776* (1948).

² Callow, L. L., and Hoyte, H. M. D., *Aust. vet. J.*, **37**, 66 (1961).

³ Legg, J., *Pamphl. Coun. sci. industr. Res. Aust.*, No. 56, 48 (1935).

⁴ Sergeant, E., Donatien, A., Parrot, L., and Lestoquart, F., *Studies on Bovine Piroplasmiasis*, 816 (Institut Pasteur d'Algérie, 1945).

ANATOMY

Possible Explosions in Absorptive Cells of the Jejunum

DURING the course of a long-term investigation of isolated, fresh, absorptive cells of the rat's jejunum suspended in physiological saline, prepared by the method described by Mohiuddin and Ramsbotham¹, we have occasionally come across cells showing a particular abnormality. This consists of a dome-like elevation of the plasmalemma over an accumulation of clear fluid covering one surface of the cell or, at the most, two adjacent surfaces. When this change affects the apex the striated border becomes unrecognizable, probably because the microvilli are taken in by the ballooning plasmalemma.

In this respect the condition differs markedly from the transient apical blistering due to rapid pinocytosis reported by Mohiuddin².

In cells isolated from animals engaged in digesting fat, the abnormal cells contained fat globules in the cytoplasm, indicating the recent presence of functional capacity. We are satisfied that extraneous factors such as change in tonicity of the saline or pressure of the coverslip were not responsible for the abnormality, because such cells were usually dispersed among normal cells. We suspect that cells showing such an abnormality might have been observed more frequently if the technique of isolation, which is based on differential centrifugation, was not biased towards elimination of cells of a density differing materially from that of normal absorptive cells.

The appearance of the abnormal cells suggests the imminence of rupture of the ballooned part, although we have not actually witnessed such an end. We are struck by the similarity of appearance between this ballooning and the changes immediately preceding the 'explosions' which seem to occur with such dramatic speed in time-lapse cinemicrographs of cells in cultures.

It is possible that cells as short-lived as the intestinal epithelium (Leblond³) are lost not only through the process of shedding, but also in other ways, including 'explosion'.

A. MOHIUDDIN

University of Lagos Medical School,
Nigeria.

H. L. CHAN
K. F. YAP

Faculty of Medicine,
Singapore.

¹ Mohiuddin, A., and Ramsbotham, M., *Anat. Rec.*, **142**, 95 (1962).

² Mohiuddin, A., *Acta Anat. Basle*, **52**, 139 (1963).

³ Leblond, C. P., in *Dynamics of Proliferating Tissues*, edit. by Price, D. (the University of Chicago Press, 1956).

Location of Renshaw Cells

EVER since their discovery¹, the mammalian spinal cord interneurons known as 'Renshaw cells'² have been the object of extensive physiological and pharmacological research³. The recurrent inhibitory pathway from motor axon collaterals via Renshaw cells back to motoneurons serves as one of the best-documented examples of a negative feed-back mechanism in the central nervous system. Furthermore, the cholinergic synapses made by the motor axon collaterals on Renshaw cells are the most thoroughly analysed central synapses from the point of view of their pharmacology.

The precise location of Renshaw cells is of considerable interest for the understanding of the functional anatomy of the spinal cord. The electrophysiological investigations of Eccles's group have shown that most Renshaw cells are to be found medial to the motor nuclei². Several investigators have described interneurons just medial to the motor nuclei which might qualify as Renshaw cells⁴. However, recently, interneurons in a still more medial position in the ventral horn have been thought to correspond to the Renshaw cells⁵.

In the experiments recorded here, Renshaw cells were located by recording their electrical activity and then marking the position of the tips of the recording electrodes. This provided a physiological control for the histological observations. The preparations used were cats anaesthetized by sodium pentobarbital. The spinal cord was exposed and several ventral roots, usually L6 and L7, were cut peripherally and the central ends mounted on platinum electrodes for stimulation. The recording electrodes were glass micropipettes filled with a solution containing a copper complex, prepared according to the method of Galifret and Szabo⁶. The resistances of the microelectrodes were initially 10–20 megohms, and they were found to be suitable for either extra- or intra-cellular recording.