Fig. 1. Guinier photographs of 9-cyanoanthracene powders. (a) Blue-fluorescent product produced by ultra-violet radiation of original green-fluorescent crystals (b); (c) photodimer

anti-parallel configuration of the photo-associated state as previously suggested for 9-methylanthracene⁶.

B. STEVENS
T. DICKINSON
R. R. SHARPE*

Department of Chemistry,
University of Sheffield.

* On secondment from I.C.I. Ltd. (Dyestuffs Division).

¹ Robertson, J. M., *Proc. Roy. Soc.*, A, **207**, 101 (1951).

² Stevens, B., *Spectrochim. Acta*, **18**, 439 (1962).

³ Stevens, B., and Dickinson, T., *Spectrochim. Acta*, **19**, 1865 (1963).

⁴ Rabaud, H., and Clastre, J., *Acta Cryst.*, **12**, 911 (1959).

⁵ Bowen, E. J., and Tanner, D. W., *Trans. Faraday Soc.*, **51**, 475 (1955).

⁶ Birks, J. B., and Aladekomo, J. B., *Photochem. and Photobiol.*, **2**, 415 (1963).

Nonaboronnonachloride

THE recent preparation of the borane B_9Cl_8H (ref. 1) prompts us to report some of our mass spectral work on the products arising from the thermal decomposition of diboron tetrachloride. When diboron tetrachloride decomposes spontaneously at temperatures more than 0° the main product is boron trichloride with lesser amounts of yellow B_4Cl_4 (ref. 2), red B_8Cl_8 (ref. 3) and dark red $B_{12}Cl_{11}$ (ref. 4); a dark purple solid has also been detected but in amounts too small to allow accurate classical analysis⁴.

The diboron tetrachloride used in this work was prepared by reducing boron trichloride in a mercury discharge⁵ and had been allowed to decompose slowly during

a period of 20 months at room temperature. A series of fractional condensations and sublimations performed in a thoroughly baked-out, all-glass vacuum apparatus allowed separation of the several species; if the baking-out process had not been carried out sufficiently well traces of a permanganate-coloured hydrolysis deposit appeared on the vessel walls. A deep blue crystalline solid, volatile at about 40° (10⁻² mm) was isolated in a virtually pure state and evaporated into a mass spectrometer at 49°; an intense group of mass peaks (407–428) was observed corresponding to the ion $B_9Cl_8^+$; the cracking pattern contained all the peaks for the ions $B_nCl_x^+$ where $n \geq x$ (Table 1). The amounts of this blue solid were too small to allow other analytical techniques to be used, but due to the similarity of the volatility and cracking pattern to B_9Cl_8H we suggest the solid is a new boron sub-chloride, nonaboronnonachloride, B_9Cl_8 .

Forstner, Haas and Muettterties¹ describe a group of peaks in the mass spectrum of B_9Cl_8H due to the ion $B_{10}Cl_{10}^+$ which they believe arose from recombination reactions in the mass spectrometer. The mass spectrum of one of our nonaboronnonachloride samples also showed peaks due to $B_{10}Cl_{10}^+$, $B_{10}Cl_9^+$ and $B_{10}Cl_8^+$. This spectrum was obtained at a lower pressure than those not showing $B_{10}Cl_{10}^+$ peaks, which would appear to rule out the possibility of recombination reactions since they would be expected to occur most readily at the highest pressures. There is therefore a distinct possibility that decaboron-decachloride, $B_{10}Cl_{10}$, exists as a stable entity and was present as an impurity in one sample of nonaboronnonachloride. Products containing boron and chlorine in a ratio 1:1 are to be expected since diboron tetrachloride produces BCl radicals during its thermal decomposition⁶.

We thank Dr. W. L. Mead and Dr. R. A. Saunders for recording the mass spectra on A.E.I. instruments (MS9 and MS8, respectively).

A. G. MASSEY
D. S. URCH

Department of Chemistry,
Queen Mary College,
Mile End Road, London, E.1.

¹ Forstner, J. A., Haas, T. E., and Muettterties, E. L., *Inorg. Chem.*, **3**, 155 (1964).

² Urry, G., Wartik, T., and Schlesinger, H. I., *J. Amer. Chem. Soc.*, **74**, 5809 (1952).

³ Jacobson, R. A., and Lipscomb, W. N., *J. Amer. Chem. Soc.*, **80**, 5571 (1958).

⁴ Schram, E. P., and Urry, G., *Inorg. Chem.*, **2**, 405 (1963).

⁵ Holliday, A. K., Massey, A. G., and Taylor, F. B. (submitted to *Inorg. Syntheses*).

⁶ Massey, A. G., and Zwolenik, J. J., *J. Chem. Soc.*, 5354 (1963).

Table 1. RELATIVE INTENSITIES OF SEVERAL PEAKS IN THE CRACKING PATTERN OF B_9Cl_8

Mass No. for $^{11}B_n^{35}Cl_x^+$	Relative intensities, $B_9Cl_8^+ = 100$
B_9Cl_8 , 414	6.1
B_9Cl_8 , 379	0.5
B_8Cl_7 , 344	7.9
B_8Cl_7 , 368	16.6
B_8Cl_7 , 333	0.25
B_8Cl_6 , 298	43.7
B_8Cl_6 , 263	2.8
B_7Cl_5 , 322	0.45
B_7Cl_5 , 287	10.7
B_7Cl_4 , 252	100.0
B_7Cl_4 , 217	3.9
B_7Cl_3 , 182	13.1
B_7 , 77	0.3
B_6Cl_5 , 276	0.4
B_6Cl_5 , 241	7.2
B_6Cl_4 , 206	16.7
B_6Cl_3 , 171	50.0
B_6Cl_2 , 230	ca. 1.5
B_6Cl_1 , 195	ca. 1.5
B_5Cl_3 , 180	25.6
B , 55	0.9