the Duffy group system. These are reported in the previous communication.

A. G. BAIKIE
O. MARGARET GARSON

University of Melbourne Department
of Medicine,
St. Vincent's Hospital.

R. G. BIRRELL

St. Vincent's Hospital,
Melbourne, Australia.

¹ Uchida, I. A., Lewis, A. J., Bowman, J. M., and Wang, H. C., *J. Pediatr.*, **60**, 498 (1962).

² Day, R. W., Wright, S. W., Koons, A., and Quigley, M., *Lancet*, ii, 154 (1963).

³ Becker, K. L., Burke, E. C., and Albert, A., *Proc. Mayo Clin.*, **38**, 242 (1963).

⁴ Gagnon, J., Katyk-Longtin, N., de Groot, J. A., and Barbeau, A., *Un. méd. Can.*, **90**, 1220 (1961).

⁵ Ford, C. E., Jones, K. W., Miller, O. J., Mittwoch, U., Penrose, L. S., Ridler, M., and Shapiro, A., *Lancet*, i, 709 (1959).

⁶ Haylock, J., Fitzgerald, M., and Danks, D. M. (personal communication).

PSYCHOLOGY

Learning in the Polychaete *Nereis*

THE rapid withdrawal reflex of the polychaetes *Nereis diversicolor*, *N. pelagica* and *Platynereis dumerilii* habituates rapidly to sudden stimuli, such as changes in illumination or mechanical stimuli, but if a foreign stimulus is interpolated in a series of stimulus presentations it is usually followed by greater reactivity to the original stimulus. Responses can sometimes be induced to sudden decreases in light intensity in completely habituated worms by the interpolation of a single tactile stimulus. This process of reflex sensitization has already been described in the polychaete *Mercierella enigmatica*¹.

Most species of *Nereis* are tubiculous or burrowing animals which come to the ends of their tubes or burrows in order to feed. When feeding they are highly susceptible to attack from a wide variety of predators including fishes, seabirds and crustaceans. Habituation of the rapid withdrawal reflex is usually interpreted as allowing for a compromise between the conflicting needs of the animal: to respond to stimuli caused by predators but not to respond to repeated innocuous stimulation such as shadows caused by passing clouds and seaweed floating in the water². Sensitization may be of biological importance because it enables habituated *Nereis* to react to a previously innocuous stimulus if there is a sudden change in the background stimulation.

Recently it has been shown that either sensitization or similar processes due to repeated stimulation by an unconditioned stimulus (US) can be important in other learning situations. In procedures designed to demonstrate classical conditioning and avoidance learning in *Nereis* the effects of the US have been responsible for modifications in behaviour which might otherwise have been interpreted as associative learning.

Attempts to condition the rapid withdrawal reflex in *N. diversicolor* have been made by pairing a sudden change in illumination (CS), to which *Nereis* habituates, with an electric shock (US), which invariably elicits withdrawal. Controls subjected to the CS alone habituate to it whereas worms subjected to a series of US-CS pairings become more reactive to the CS. However a similar effect can be produced by presenting the US between trials so that it is not paired with the CS (Fig. 1). Furthermore, controls habituating to CS presentations become more reactive if one or more groups of ten consecutive electric shocks are given between trials. Sensitization of the withdrawal reflex by repeated effects of the electric shocks is clearly a more tenable interpretation of these data than one of classical conditioning.

The repeated effects of US stimulation have also been shown to have an important effect in a learning situation

Fig. 1. Percentage of worms *N. diversicolor* reacting to sudden decreases in light intensity (CS) during the following training procedures in which an electric shock has been used as a US: CS-US pairing (continuous line, open circles); US between CS presentations (continuous line, filled circles); CS presented alone (broken line). Standard errors are indicated by the vertical lines.

previously described by the author³ in which individual *N. diversicolor*, *N. virens* and *P. cultrifera* could learn to avoid an electric shock at the end of a 'Perspex' channel. Worms introduced into the channel crawl rapidly to the end and continue to do so indefinitely as long as they are not punished in any way. However, the behaviour of worms given slight electric shocks, whenever they reach the end of the channel, becomes modified with training so that they either reverse in the channel before reaching the electrodes or refuse to crawl along it. It has now been shown that electric shocks given outside the apparatus are as effective in inducing this modified behaviour as shocks received at the exit of the channel, and that shocks given to worms with no experience in the channel have the same effect on subsequent performance as shocks given in the apparatus. Strong tactile stimulation as a punishment has been shown to have a similar effect. Although an explanation of fatigue can be excluded because trained worms crawl actively outside the apparatus and refusal itself often involves an active reversal by the worm, there is no doubt that the acquired behaviour cannot satisfactorily be regarded as associative learning. It is more likely to be the result of a process similar to sensitization of the reversal response, which is part of the worm's normal tubiculous behaviour⁴, and a mechanism which inhibits forward movement in a confined space, such as a channel or burrow, when the worm is harmfully stimulated.

The importance of sensitization and processes producing similar effects has yet to be evaluated. The effects of these processes can be regarded as learning in the sense that they are modifications of the animal's behaviour as a result of experience but clearly cannot be regarded as manifestations of associative learning. Although similar processes have not been demonstrated in Nature, the possibility of widespread sensitization and non-associative learning mechanisms in the lower invertebrates cannot be excluded. Indeed, it has already been suggested⁵ that the conditioning claimed in flatworms⁶ may be due to the sensitization effects of electric shocks. A study of these processes may be valuable because they may have been important in the evolution of the more refined learning mechanisms of vertebrates and higher invertebrates.

S. M. EVANS

Department of Zoology,
University of Bristol.

¹ Rullier, F., *Bull. Lab. Mari. Dinard*, **30**, 21 (1948).

² Nicol, J. A. C., *J. Mar. Biol. Ass. U.K.*, **29**, 303 (1950).

³ Evans, S. M., *Anim. Behav.*, **11**, 172 (1963).

⁴ James, R. L., and Halas, E. S., *Psychol. Rev.*, **14**, 1 (1964).

⁵ Thompson, R., and McConnell, J. V., *J. Comp. Physiol. Psychol.*, **48**, 65 (1955).