carbon ages, he states: "It is interesting to find in all three profiles formal ages of the *B* horizon humus *ca.* 400 yr. This can be taken as evidence of a dynamic equilibrium between the supply and breakdown of organic matter in the *B* horizon of the podzol profiles of the type studied."^{2,3} He also collected the humus of *B* horizons from podzols in Lapland and in sixteen samples from different profiles the formal radiocarbon ages ranged from 460 ± 65 to $1,260 \pm 60$ B.P.² Oden collected samples of a soil formed on loess at Hildesheim in Germany, and these were dated as follows:

St-681	8-11 cm	Apparent age: $1,360 \pm 60$ yr
St-1,020	30-50 cm	" $2,090 \pm 70$ yr
St-682	68-71 cm	" $3,170 \pm 65$ yr

His comments imply that the ages are relatively low compared with that of the loess⁴.

So far, no Ando soils appear to have been investigated in this way, but the principles are likely to be the same; however, because of their peculiar organo-mineral complexes they may contain a higher proportion of stable organic compounds and give somewhat greater formal radiocarbon ages than those already cited.

It would thus seem that the organic carbon of a buried soil layer is of more value in dating the deposition of overlying material than in dating the deposition of its own parent material. Thus the age ($8,650 \pm 200$ yr) of the organic carbon of the fifth layer at Choyo (Table 1) gives a maximum age for the deposition of the fourth layer, but gives only a minimum age for the fifth layer which could have been deposited tens of thousands of years earlier.

To determine the approximate age of the fifth layer it will be necessary to date the organic carbon of a buried soil lower in the column, or else to find charcoal (of vegetation burnt at the time of its deposition and incorporated in the base of the deposit) with which to date it, as has been done in New Zealand for the Taupo Pumice and older airfall ash-beds⁵. If it transpires that the age is beyond radiocarbon measurement it will leave in doubt whether the halloysite in this profile is in fact any younger than that at Kanoya. In which case the organic carbon of the buried surface layers 4 and 2 at Kanoya could be dated to help in narrowing down the problem.

J. E. Cox

Soil Bureau,
Box 1005,
Whangarei, New Zealand.

¹ Aomine, S., and Miyauchi, N., *Nature*, **199**, 1311 (1963).

² Tamm, C. O., in Östlund, H. G., and Engstrand, L. G., *Radiocarbon*, **5**, 220 (1963).

³ Tamm, C. O., and Östlund, H. G., *Nature*, **185**, 706 (1960).

⁴ Oden, S., in Östlund, H. G., and Engstrand, L. G., *Radiocarbon*, **5**, 220 (1963).

⁵ Healy, J., Vucetich, C. G., and Pullar, W. A., *N.Z. Geol. Surv. Bull.*, **73** (1964).

Cox objects to the method which Miyauchi and I have used to estimate the age of hydrated halloysite at Choyo¹. It is possible, however, that because of the different nature and properties of volcanic ash soils in New Zealand and Kyushu, some of his objections have arisen from a misunderstanding of the soils in Kyushu.

First, he points out a sharp contrast in the clay mineralogy of the fourth and fifth layers at Choyo, the ages of the ashes of which should not differ. In the profile of volcanic ash soils in Kyushu, an old ash is not necessarily always at a more advanced stage of weathering than a young one, and in certain cases a sharp contrast in mineralogy can be observed even within a given soil layer². The degree of weathering of ash depends not only on the time since deposition but also on many other factors.

Second, Cox has questioned the point that no halloysite was detected in any layer above the sixth layer (older than 30,000 yr) at Kanoya. An answer to his first question would, in fact, also answer this. He also suggests that the dating of the second and fourth layers of this profile would

be more serviceable in narrowing down the problem. Recently, the upper part of the fourth layer (130-150 cm) was estimated to be $5,450 \pm 160$ yr old by the radiocarbon method³, and two layers of another profile, corresponding to the second and fourth layers are $4,640 \pm 80$ and $6,360 \pm 90$ yr old, respectively, according to Matsui (personal communication). Matsui considers that the age of the fourth layer estimated by the radiocarbon method is roughly correct, because earthenware with a strawrope pattern belonging to the early Jomon period (about 5,000-7,000 yr B.P.) has been found below the third layer in Miyazaki, which corresponds to the fourth layer of the Kanoya profile. Discordance might have been between the fifth and sixth layers.

Third, Cox points out that the age of organic matter of a surface soil layer can be much younger than the time taken for the soil to form. This is the usual case, but accumulation of organic matter in Andosols seems to be carried on in a somewhat different way from that in normal soils. With fine ash, organic matter can start to accumulate in the surface layer soon after deposition of the ash, probably within a few years, at the same time as amorphous inorganic colloids are formed. Accumulated organic carbon can reach as high as 20 per cent or more in well drained Andosols under the warm and humid climate of Kyushu, and the amount in surface layers is proportional to the clay content of the layers⁴. It is well known that the humus of Andosols is very stable and exhibits unusual properties, and this is especially true for the humus in the buried layers. This stability is corroborated by the finding that humus 9,000 years old or more is present in abundance in these soils.

Provided that organic matter has not been added to a layer since it became covered with another deposit ash, the accumulated organic matter, as a whole, should be younger than the ash of the layer itself and older than the ash of the overlying layer. If a period of accumulation of organic matter of the layer is negligible in comparison with the age of the ash of the layer, then the age of the layer would be approximately equal to the age of the former. If this period of accumulation could be estimated in some way, the value could be used to correct the age of the overlying ash. The period of accumulation of organic matter of the fifth layer of Choyo was assumed to be 330 years on the basis of 1 ton/hectare/yr of organic carbon accumulated¹. This seems, however, to have been something of an underestimate in that the rate of accumulation was overestimated; recently, this was estimated to be only 0.14 ton/hectare/yr (ref. 3). Because the mechanism of accumulation of organic matter in Andosols is obscure, the radiocarbon dating of organic matter of a buried soil layer cannot be applied to an exact determination of the age of an ash but only to its approximate age.

SHIGENORI AOMINE

Faculty of Agriculture,
Kyushu University, Fukuoka, Japan.

¹ Aomine, S., and Miyauchi, N., *Nature*, **119**, 1311 (1963).

² Aomine, S., and Wada, K., *Amer. Min.*, **47**, 1024 (1962).

³ Aomine, S., and Miyauchi, N., *Soil Sci. Plant Nutrit.*, **11**, 213 (1965).

⁴ Aomine, S., and Wada, K., *Soil Sci. Plant Nutrit.* (in the press).

PSYCHOLOGY

Mechanism underlying the Sleep-Dream Cycle

THE cyclic nature of sleep in mammals, including the regular recurrence of a state characterized by a low voltage desynchronized electroencephalogram, rapid eye movements, and heightened autonomic activity, is now firmly established^{1,2}. This state, associated in humans with dreaming, has been called "dreaming state", "rapid eye-movement sleep" or "paradoxical sleep", and we refer to it as the "*D*-state"³.