Where sky conditions were favourable, observers in the south-western United States saw too many meteors to make a direct count as the shower reached its climax, at 11.55 universal time. The burst of activity started at about 11.30 and lasted for an hour. A group on Kitt Peak, Arizona, estimated that 40 meteors could be seen in a one-second sweep of the observer's head at the peak of the shower, with more than 1,000 meteors a minute from 11.30 to 12.30 (4.30 to 5.30 a.m. mountain standard time). Similar reports, estimating tens of meteors per second at peak activity, came from California, Colorado, New Mexico and Texas. Accounts from the Atlantic states also indicate heavy activity, but are not quite so impressive since dawn began to interfere while the rate was still increasing.

Meteor showers are believed to be the debris of comets, and the 33 year periodicity of the Leonids is caused by an intense concentration of meteoric particles near the point on the orbit where the parent comet is located. The non-appearance of the display over Britain can be attributed to the comparatively small size of this swarm, but it is evident that the Leonids are still capable of giving a brilliant display.

Mechanism of Polarity

from our Cell Biology Correspondent

As Imamoto and Yanofsky (*J. Mol. Biol.*, **28**, 1 and 24; 1967) have now shown, polar mutations cause failure in transcription; genetic polarity is not caused exclusively by defects in translation.

Transcription of polycistronic operons, like the tryptophan and lactose operons of *E. coli*, produces a single mRNA molecule containing information for several enzymes. Amber, ochre and frame-shift but not mis-sense mutations in such operons may show polarity; a polar mutation prevents the expression of the cistron in which it occurs and also leads to reduced levels of the enzymes specified by the subsequent, operator-distal, cistrons of the operon. Not all polar mutations in any one cistron show the same degree of polarity; in the *lac* operon, for example, there is a continuous gradient of polarity across the β galactosidase cistron from the very strongly polar mutations at the 5', N terminal end of the cistron to the very weakly polar mutations at the 3', C terminal end.

How do amber and ochre mutations cause polarity? There are three possibilities: (1) by reducing the frequency of the translation of regions of mRNA beyond the nonsense codon; (2) by preventing transcription of mRNA from DNA after the nonsense codon; or (3) by a combination of both these processes. Because suppressor gene transfer RNA prevents chain termination by amber and ochre codons and also relieves polarity, it seemed that polarity might result from defects in translation, and several groups have devised purely translational models for polarity.

Attardi *et al.* (1963), however, found that extremely polar mutants of the *E. coli* operon fail to make *lac*-mRNA; perhaps polar mutants prevent transcription. Using a hybridization technique to detect *tryp*-mRNA, Imamoto and Yanofsky have now shown that strains of *E. coli* with strongly polar mutations in the tryptophan operon do indeed make less *tryp*-mRNA than identically treated wild type cells

or mis-sense mutants. The *tryp*-mRNA molecules of the polar mutants are short; they lack the regions that are normally transcribed from the DNA beyond (operator distal) the site of the polar mutation. Furthermore, the size of the predominant *tryp*-mRNA species in different mutants increases in relation to the genetic distance between the beginning of the first cistron of the operon and the site of the mutation.

Estimates of the relative proportions of complete and short mRNA molecules made by the different mutants correlate fairly well with their previously observed translational polarity; for example, a mutant with 14 per cent translational polarity—it makes only 14 per cent wild type levels of enzymes specified by the cistrons following the mutated cistron—made only 10 per cent complete *tryp*-mRNA molecules and 90 per cent short molecules. Pulse labelling of the *tryp*-mRNA as it is synthesized revealed that the mutants make normal numbers of mRNA molecules and that the short molecules are not produced by degradation of complete molecules. Polarity, therefore, certainly involves and may even be caused entirely by failure in transcription.

How can this finding be reconciled with evidence that polarity is a translational phenomenon? A nonsense codon itself is not a signal to RNA polymerase to stop transcription, and genetic translation and transcription are closely coupled; ribosomes somehow regulate mRNA synthesis (see *Nature*, **214**, 228; 1967). Hence Imamoto and Yanofsky suggest the continued transcription of DNA by RNA polymerase depends on the normal movement of ribosomes translating that part of the mRNA molecule already synthesized. If this ribosome movement is impeded by a nonsense mutation and a subsequent untranslatable region of mRNA, then the greater the distance between the nonsense codon and the initiation site of the next cistron in the DNA being transcribed the lower the probability that RNA polymerase will continue transcription and so the mRNA molecule will be terminated.

Protean Proteins

from our Molecular Biology Correspondent

It is interesting to speculate whether the native state of a protein represents a single uniquely stable conformation, or whether it is better regarded as one of a number of states occupying potential energy minima, between which transitions can occur in response to environmental adjustments. The proteins trypsin and trypsinogen have recently been subjected to careful scrutiny by several groups working in France, and the results which have been reported have a general bearing on this question.

Lazdunski and Delaage (*Biochim. Biophys. Acta*, **140**, 417; 1967) have studied pH-dependent conformational transitions in both the enzyme and zymogen, and have observed the existence of three reversible equilibria, linking four definable conformational states. The changes are followed on the one hand by difference spectra, which reflect the state of exposure of tyrosine and tryptophan residues, and on the other by optical rotation. Three at least of the observed isomeric states appear not to be denatured forms, in the sense that the protein retains in each case characteristics