conformational isomers, and that the equilibrium is displaced by the added dye in favour of the form to which it most avidly binds. In the present case, this form is evidently one in which the susceptible peptide bonds are relatively unavailable, or in which the structure is comparatively rigid and inflexible. Markus and his co-workers have previously recorded a case in which the binding of a ligand renders a protein more prone to proteolysis, and the converse presumably holds in their latest work.

Several examples of the binding of dyes at or near the active sites of enzymes have recently been reported. The latest of a number of studies by Glazer (*J. Biol. Chem.*, **242**, 4528; 1967) concerns the binding of the anionic dye, Biebrich scarlet, to chymotrypsin. It has proved generally true, as revealed notably by X-ray studies on enzymes, that active sites are situated in crevices, giving on to the predominantly non-polar interior of the molecule. Equally, most organic ligands which are capable of binding strongly to a protein evidently seek an essentially non-polar environment (as shown, for example, by dyes which exhibit fluorescence only in substantially non-aqueous solvent systems, and which become fluorescent when they bind to proteins). Thus it is not surprising that a pattern is beginning to emerge of binding of many such small molecules specifically at active centres. Several cationic dyes have been found to bind at the active centre of chymotrypsin, and it is interesting that Biebrich scarlet, being anionic, binds with comparable affinity. The binding can be followed by changes in the absorption spectrum of the ligand. The dye competes with, and is displaced by, substrates and competitive inhibitors. It does not bind to chymotrypsinogen, which suggests that here the active site is occluded.

Another application of dye-binding is due to Koshland, who uses a covalently attached chromophore as a so-called "reporter" for events involved in the binding and conversion of substrates. Hille and Koshland (*J. Amer. Chem. Soc.*, **89**, 5945; 1967) have bound a compound containing the highly medium-sensitive nitrophenyl chromophore to methionine-192 of chymotrypsin, which is three residues along the chain from the serine of the active site. The activity of the enzyme is preserved, and one can then observe the perturbation of the nitrophenyl spectrum, when for example the pH is changed. The magnitude of the perturbation, in fact, follows a titration curve, with a pK of 7. Addition of substrate eliminates this effect, and it is suggested that the bound chromophoric group is pushed out of the way. The speculation is that the ionizing group is the active-site histidine, and if this is true its pK is thereby established. It must be hoped that a comparison of data of this kind with the environment of the residues in question in the crystallographic structure will ultimately lead to general correlations about their properties, and that this will help in interpretation of the mechanism of catalysis.

Chromosome Breakage

from our Cytogenetics Correspondent

RADIATION and many chemicals used in cancer chemotherapy and as antibiotics cause chromosome breakage. Breakage is a result of the disruption of DNA synthesis

or some other disturbance of chromosome organization and so interest in induced chromosome breakage lies in what this can say about the structure and chemistry of chromosomes. One important question is whether chromosomes in different phases of the cell cycle are equally sensitive to damage. In both animal and plant cells there is evidence that chromosomes are more sensitive to certain agents during the G_2 phase of interphase (that is, between the DNA synthetic stage, S , and mitosis) than during the S or pre- S (G_1) phases of interphase. Methyl coumarin has been shown by Ronchi and her colleagues (*Mutation Res.*, **4**, 615 and 791; 1967) to cause most breaks to bean and onion chromosomes during G_2 as well as to prolong this phase. They also showed that exposing roots previously treated with methyl coumarin to chloramphenicol causes an increase in the number of breaks, and they interpret this as evidence that protein synthesis is required to repair breaks. Breaks are more numerous in the presence of chloramphenicol because many of the breaks induced by methyl coumarin have not been allowed to heal. The requirement of protein synthesis for healing breaks is not a new idea, but it does pose some obvious questions. Is it a repair enzyme or a protein of chromosome structure that is synthesized? And if it is an enzyme, can differences in its activity account for differences in apparent sensitivity of chromosomes during the cell cycle?

Scott and Evans have also shown (*Mutation Res.*, **4**, 579; 1967) that chromosomes of bean at G_2 are most sensitive to X-rays and, further, that chromosomes at different points in the G_2 phase differ in sensitivity. Similarly, Dewey *et al.* (*Int. J. Radiat. Biol.*, **12**, 597; 1967) showed that Chinese hamster chromosomes sustained 2.5 times more chromosome damage in G_2 than in S . In their experiments the chromosome damage was self inflicted as the chromosomes were allowed to incorporate a high level of radioactivity in the form of tritiated thymidine.

Breaks in chromosomes could be caused by interference with nucleic acid metabolism rather than a physical shattering of the chromosome thread. This at least seems to be one of the effects of 9- β -D-arabino-furanosyladenine (ara-A). Its triphosphate has been shown by Furth and Cohen (*Cancer Res.*, **27**, 1528; 1967) to be able to inhibit DNA polymerase of mammalian cells (but not that from cells of the bacterium *Escherichia coli*) and may also inhibit the nucleotide reductase system. But ara-A is able to cause breaks at G_2 ; could it be that there is some DNA synthesis not detectable by conventional methods during G_2 that all these agents inhibit?

What of induced chromosome breakage in meiotic cells? Westerman (*Chromosoma*, **22**, 401; 1967) describes the effects of irradiating meiotic chromosomes during spermiogenesis in male locusts. He found that cells irradiated during the S phase of meiosis showed chromosomes with a lowered chiasma frequency at the subsequent metaphase, while cells irradiated at diplotene or in the S phase of premeiotic mitosis showed an increase in chiasma frequency. The chiasma frequency was most changed in the long chromosomes while that of the short chromosomes was hardly changed at all. It is tempting to speculate on the role of chromosome breakage on the time and mode of chiasma formation, but only further experimental work will allow concrete fact to replace speculation in this perplexing problem.