transcarbamylase and its relation to the function of the enzyme. The central finding (Gerhart and Schachman, *Biochemistry*, **7**, 538; 1968) was that there are six sub-units, two bearing catalytic and four bearing regulatory sites. This unique situation was considered in terms of an allosteric scheme for the enzyme. Hard on the heels of these papers comes a study of the same enzyme by Weber (*J. Biol. Chem.*, **243**, 543; 1968) from which it appears that these results are in error, inasmuch as aspartate transcarbamylase has in fact a rather more commonplace constitution, being made up of four regulatory and four catalytic sub-units.

Whereas Gerhart and Schachman worked with the fragments obtained when the enzyme is dissociated by treatment with an aromatic mercurial, and assumed these products to represent monomeric chains, Weber performed molecular weight determinations in 8 M urea. Sedimentation equilibrium led to values of some 42,000 and 23,000, respectively, for the catalytic and regulatory chains. These results were supported by N-terminal residue determinations and amino-acid analysis: one N-terminal alanine and one threonine were found for an equivalent weight of 73,000, and the amino-acid compositions for the two sub-units together reproduced closely the composition of the intact enzyme. It therefore now seems safe to conclude that the protein is an octamer and that the catalytic and regulatory chains each possess one active site. The number of sites (four of each kind in the native enzyme) was reported by Changeux, Gerhart and Schachman, who have discarded an earlier estimate of eight regulatory sites.

The moral yet again would seem to be that it is hazardous and unnecessary to assume that monomeric chains are present under any but extreme denaturing conditions. Numerous molecular weight determinations of proteins in guanidine hydrochloride solutions—the use of which has been urged notably by Tanford and by Harrington—are now known to have given correct monomer molecular weights, and it is, one must suppose, an indication of the success of this approach that its re-discovery on this side of the Atlantic should already have been described (Ullmann, Goldberg, Perrin and Monod, *Biochemistry*, **7**, 261; 1968).

It may be mentioned that Sia and Horecker (*Arch. Biochem.*, **123**, 186; 1968) are now in agreement with the four-chain structure of the enzyme aldolase, long a subject of contention. They elaborate, moreover, on the remarkable degradation of the polypeptide chain of this enzyme in quite mildly acid solution. The possibility that such phenomena may seriously vitiate molecular weight determinations in special cases must be recognized.

T₄ Regulation

from our Cell Biology Correspondent

DURING the replication of T-even phages, gene functions appear in a time sequence, which is why most viral functions have been classified as either early or late. Refinements in methods of analysis have, however, shown that this is an oversimplification. Levinthal, Hosoda and Shub, for example, have found that early proteins are synthesized asynchronously; they are not a single co-ordinately controlled set. There are also sequential changes in the population of phage mRNA molecules (see *Nature*, **216**, 117; 1967). But

only recently, with the improvements in the sensitivity of DNA/RNA hybridization techniques, has it been possible to measure accurately gene control at the level of mRNA transcription. In *J. Mol. Biol.* (**31**, 325; 1968), Bolle, Epstein, Salser and Geiduschek describe DNA/RNA hybridization experiments which begin to show the complexity of the regulation of T₄ transcription.

Twenty minutes after infection, a third to a half of the phage mRNA molecules are late mRNA species. The remainder of the population of mRNA at 20 minutes consists of early-type molecules which are also present 5 minutes after infection, when the late species are between a hundred and several thousand times less abundant. There is no difference in the average stability of early and late mRNA. It is also clear that the whole T₄ genome is not translated for a brief period immediately after infection; some late species seem not to be made at all in the first 5 minutes after infection. Bolle *et al.* also mention briefly other results to be published in detail later. Apparently there are three classes of early mRNA molecules which are turned on rapidly in succession during the first 5 minutes of infection. Synthesis of the majority of these RNA species is then reduced several-fold after between 5 minutes and 20 minutes, during the period in which the late mRNAs are switched on. Even so, a few of the RNA species present after 5 minutes increase during this period.

It is clear that regulation of T₄ gene expression cannot be solely at the translational level, and the most reasonable explanation of the 200 to 4,000-fold increase in late mRNA between 5 and 20 minutes is that late mRNA transcription is specifically switched on in this period. What the switch mechanism is and whether all late genes are activated at the same time remains to be seen.

In the same issue of *J. Mol. Biol.* (**31**, 463; 1968), Chrispeels, Boyd, Williams and Neidhardt report that during T₄ infection of *E. coli*, a new valyl tRNA-synthetase activity appears with an accompanying decrease in the amount of the original *E. coli* valyl tRNA-synthetase. The T₄ induced enzyme differs in sedimentation properties and thermal stability from the *E. coli* enzyme, but heavy labelling experiments prove that it is formed from polypeptide chain synthesized before infection. Chrispeels *et al.* believe that the phage-induced enzyme is a modified form of the host enzyme, possibly a dimer. This conversion is specific (arginine, aspartic, leucine, phenylalanine, and serine-tRNA synthetases of *E. coli* are not modified by T₄ infection) and it is sensitive to chloramphenicol. For these reasons, Chrispeels *et al.* suggest as a working hypothesis that a new or altered valine tRNA produced by T₄ promotes the conversion of the host enzyme. Whether the new enzyme plays an important part in the physiology of the phage is undecided, but since conversion of all the host enzyme occurs by about 15 to 20 minutes after infection, it is just possible that the conversion is involved in the regulation of early and late protein synthesis.

Globin Metabolism in Thalassaemia

from our Medical Biochemistry Correspondent

THALASSAEMIA is a chronic haemolytic anaemia which is quite common in some parts of the world, particu-