LETTERS TO THE EDITOR

PHYSICAL SCIENCES

Statistical Properties of Pulsar CP 1919

THE pulsar CP 1919 (ref. 1) was observed at Parkes² in March 1968 at a number of radio frequencies. On occasions the frequencies 85, 150, 150.9, 630 and 1,410 MHz were used simultaneously, with the 210 foot telescope tracking the source for some 4 h. The data have now been examined for statistical occurrence properties and a quantitative description of the source can be given. The statistics of pulse occurrence are of general interest because theories of emission (and pulse propagation) require these data. And anyway an observatory embarking on a programme of searching for such objects must know the statistical properties of pulse occurrence to aid in determining an observational programme.

Some of the statistics of pulsars have been described already in general terms. At lower radio frequencies pulses occur in short bursts. At higher frequencies pulses occur in longer trains. Most of the published reports have concentrated on the description of the events when the pulses were strongest. Some preliminary statistics on pulse magnitudes have been given for the frequency of 408 MHz (ref. 3). This communication concentrates on the pulse amplitude occurrence statistics for the pulsar CP 1919 at the frequencies observed at Parkes.

The methods of observation and calibration have already been described². The mean values of pulse amplitude at each frequency were determined from long integrations made in a sequential computer, synchronized to the pulse period. The pulse occurrence data were derived from chart records made with the time constant of 0.1 s. A selection was made of charts where a nearly continuous record was available for at least 5,000 pulses (2 h of tracking).

At a frequency of 85 MHz the noise on the record made the task of identifying pulses less than 20 flux units in amplitude very difficult. The mean value of pulse amplitude was determined to be 1.5 flux units. The inspection of records for a 5,000 pulse period revealed one burst with 7 pulses in the range of 50–100 flux units. A further

Fig. 1. Histograms of pulse occurrence against pulse flux at 150 MHz. Pulses are grouped with sequential number designation.

Fig. 2. Histograms of pulse occurrence against pulse flux at 630 MHz. Pulses are grouped with sequential number designation. The 100 pulse groups show examples of extremes in pulsar activity.

burst was observed in the 5,000 pulse period with a few pulses in the range of 40–60 flux units. At the frequency of 150 MHz the mean value of pulse amplitude was 9 flux units and the detection limit was 4 flux units. Histograms of pulse occurrence were drawn for 5 flux unit steps as shown in Fig. 1. Increases in flux amplitude of pulses in the form of bursts were observed at least once every 200 pulses. The histogram of 5,000 pulses shows several large pulses to be present in the observed sample. At the frequency of 630 MHz the mean pulse amplitude is 1.2 flux units and the detection limit is 0.6 flux units. There is a distinct difference in the histograms shown in Fig. 2 for this frequency as compared with the 150 MHz values. Pulses are weak for some 4,000 pulse periods and then consistently strong for some 500 pulses. The histogram of 5,000 pulses also shows a number of large pulses which, however, occur in one long active period. At 1,410 MHz very few pulses were actually observed. The mean flux was found to be 0.3 flux units. Occasional pulses up to 3.0 flux units, however, were observed in the 5,000 pulse period.

The statistics of pulse occurrence give a complicated picture of the behaviour of pulsars. It has already been noted⁴ that time scale difficulties preclude a simple thin screen scintillation explanation of the pulsar pulse to pulse amplitude variations. The histograms given for the frequency of 150 MHz may allow an explanation by deep scintillations. It would be difficult, however, to fit the 630 MHz results into such an explanation.

For a pulsar search, it seems that a frequency near 150 MHz is optimum with a tracking period of 5 min for each sky point. It would be interesting to seek data to establish whether there is a correlation between pulse occurrence at a particular frequency and pulse period.

I thank Dr B. J. Robinson and Mr B. F. C. Cooper of the Radiophysics Division, CSIRO, and Mr T. L. Landecker of the School of Electrical Engineering, University of Sydney, for cooperation in observations of the pulsar at Parkes.

R. WIELEBINSKI

School of Electrical Engineering,
University of Sydney.

Received July 15, 1968.

¹ Hewish, A., Bell, S. J., Pilkington, J. D. H., Scott, P. F., and Collins, R. A., *Nature*, **217**, 709 (1968).

² Robinson, B. J., Wielebinski, R., Cooper, B. T. C., Gardner, F. F., and Landecker, T. L., *Nature*, **218**, 1143 (1968).

³ Grueff, G., Roffi, G., and Vigotti, M., *Nature*, **218**, 1036 (1968).

⁴ Scheuer, P. A. G., *Nature*, **218**, 920 (1968).