unfair of him to suggest that the theory of Fowler and Hoyle is "motivated" by the apparent similarity of the decay of the light curve of the Crab nebula and of the radioactive decay of ^{254}Cf —that was a kind of bonus in their scheme. Dicke himself starts from the view that even the oldest stars contain proportions of the heavy elements which are comparable with those now found in the Sun, which in his view implies that the processes of nucleogenesis previously thought to be confined to supernovae actually took place in the few hundred million years during which the proto-galaxy collapsed into more or less its present shape. With this starting point, the disagreement with earlier theories follows quite simply, although Dicke also has the advantage of more recent estimates of the ratios of the concentrations of the isotopes of thorium and uranium at present in the Galaxy. The most striking difference is his adoption for the ratio of ^{232}Th to ^{238}U of 3.18 rather than the 3.8 assumed by Fowler and Hoyle.

In fields like these, factors of a mere two may not be much of a surprise. The physical picture which Dicke has provided of the origin of the Galaxy will, however, raise broader issues which are unlikely to be laid to rest by his most recent statement of his case.

PHYSICS

The Intermediate Boson

from our own Correspondent

THE possible existence of the intermediate boson or "W" particle has been the subject of speculation for several years, and Professor G. K. O'Neill of Princeton University devoted the second of his special lectures last week at University College, London, to describing an experiment to locate the W boson using the new intersecting storage rings at CERN.

If the W boson exists it is related to the weak interactions such as those in β -decay in much the same way that the photon is related to electromagnetic processes. The W would thus provide the weak interaction link between nucleons (protons or neutrons) and the leptons (electrons, μ -mesons and neutrinos). Various experiments have been carried out to try to find the W, but all of them have yielded negative results. They imply, however, that the mass of the particle—if it exists—is probably more than three times the mass of the proton. A particle of such high mass is most likely to be observed in reactions where there is a large available energy, and this has prompted various proposals for search experiments using the intersecting storage rings, in which proton collisions of about 1,600 GeV (in the laboratory frame of reference) will take place. Even though this interaction energy is higher than that of the new generation of proton accelerators, which will have energies in the range of 200–300 GeV, the storage ring system produces far fewer interactions per second. This means that an experiment to search for the W would be very time consuming.

It is thought that the principal decay modes of the W would be into a neutrino together with an electron or a μ -meson. Most of the suggestions for experiments to show the existence of the W involve detecting its decay μ -mesons. These particles are likely to have a large momentum because the whole mass of the W is available as energy to be shared between the neutrino

and the μ -meson. Such μ -mesons could be identified by making use of the fact that they can pass through very large amounts of material without undergoing nuclear interactions. This feature distinguishes them from such strongly interacting particles as π and K mesons. But a background contribution of μ -mesons is expected from such processes as the decay of the π and K mesons, which will certainly be produced in large numbers in very high energy interactions.

The special features of Professor O'Neill's proposal are, first, to use a very large spark chamber detection system to cover nearly all the 4π of the solid angle into which interaction products may go and, second, to concentrate chiefly on the electron decay products of the W. The electron detection would be carried out by measuring the momenta of particles before and after they have traversed a thin lead absorber. The very light charged electrons would have their momentum significantly changed by the absorber due to electromagnetic radiation processes. Other heavier particles would be less affected and would thus be excluded. In addition, the system would include the usual μ -meson detection equipment, the whole apparatus being placed inside a large uniform-field magnet.

The background problems should be much less important in the case of electrons than for μ -mesons, according to Professor O'Neill. He suggested that the chief contribution to electron backgrounds will come from neutral π -meson decays into electron-positron pairs, the so-called Dalitz decays, with the second member of the pair not being seen. This background could be small compared with the W decay electrons, provided that the W has a large enough mass to produce very energetic electrons. It is to be hoped that these very high momentum electrons would be found in larger numbers than could be accounted for from any background process.

MATHEMATICS

Random Numbers in the Planes

from our Mathematical Probability Correspondent

DURING the past fifteen years, in step with the rapid development of computers, Monte Carlo methods have been used to solve a wide variety of problems. Estimation in statistics, simulation of random processes, evaluation of integrals and solving of linear operator equations are some of the problems to which Monte Carlo methods have been successfully applied. To find its proper place among applied mathematics, pure analysis and numerical analysis, it has not been enough to find new fields of application for a Monte Carlo approach. The Monte Carlo methods themselves have had to become increasingly sophisticated in order to become powerful tools of analysis with a real advantage over other techniques. It is disquieting, therefore, to discover flaws at the heart and source of Monte Carlo which no amount of subsequent sophistication can overcome.

Monte Carlo involves, essentially, a sequence of operations on a sequence of random numbers $\{u_i\}$, where, ideally, the $\{u_i\}$ are independent and identically distributed. All would be well if the $\{u_i\}$, like the integers, were God given. But in practice the $\{u_i\}$ must be generated artificially in large numbers and at high speed on the computer. The numbers are almost