Interaction between RNA and protein is the basis of ribosome formation. By skilful manipulation of microdissection and microelectrophoresis, J.-E. Edström (Stockholm) has been able to separate events taking place in nucleolus, chromosomes and nuclear sap and to give a picture of the build-up of ribosomal RNA components. He postulates that 28S and 23S components are formed in the nucleolus and then transported to the chromosomes. Here the 23S is further processed to 18S RNA, before both 28S and 18S components are exported.

Interactions of macromolecules in the nucleus do not occur in isolation, but are profoundly influenced by cytoplasmic proteins. This was convincingly shown by the nuclear transplant experiments of J. Gurdon (University of Oxford) in which adult frog brain nuclei inserted into egg cytoplasm swelled markedly and rapidly changed their synthetic activity to coincide with that of the egg cytoplasm. Cytoplasmic proteins up to the size of bovine serum albumin can freely enter through the nuclear membrane, and it is these, rather than proteins transplanted with the nucleus, that control events taking place. In spite of the artificial nature of these experiments, there are enough similarities between nuclear changes here and in normal embryogenesis to hope that these very exciting results will be relevant to the control of nuclear activity in general.

MOLECULAR GENETICS

The Lactose Operon

from our Cell Biology Correspondent

BECKWITH's group at the Harvard Medical School seems to be dominating work on the lactose operon of *E. coli* to almost the extent that Kornberg's group dominates research on DNA polymerase. The fifty-odd pages of the June 10 issue of the *Journal of Biological Chemistry* devoted to the Kornberg group's latest results (see page 347) are eloquent testimony of the latter phenomenon. And in the latest issue of the *Journal of Molecular Biology* (43, 201; 1969) Reznikoff, Miller, Scaife and Beckwith report another of their delightfully elegant manipulations of the *lac* operon, this time designed to prove that the *lac* repressor protein can stop in their tracks RNA polymerase molecules which are actively transcribing DNA. Largely as a result of the work of Beckwith's group, we now know that three genetic elements in the *lac* operon precede, in the following order, the three structural genes of the operon; first there is the *i* gene which specifies the repressor, then the promoter gene *p* at which transcription is initiated and then the operator site *o* which binds the repressor and then the three genes for the enzymes of the operon.

To prove that repressor protein can block an RNA polymerase molecule already engaged on transcription, Reznikoff *et al.* fused, by deletion mutation, the *lac* operon to the tryptophan operon, eliminating the *lac i* and *p* genes in the process. This results in an *E. coli* strain in which the operator and structural genes of the *lac* operon are continuous with the last structural gene of the *trp* operon. The only way in which the *lac* operon can be transcribed is by first initiating and transcribing the *trp* operon. The RNA polymerase reads through from the end of the *trp* operon to the

beginning of the *lac* operon fused to it and lacking its own promoter site.

In this situation, they added *lac* repressor which bound to the still intact *lac* operator (*o*) site and assayed for continued transcription of the *lac* structural genes. The binding of repressor caused up to a ten-fold reduction in synthesis of the *lac* enzymes. Repressor can therefore block the progress of RNA polymerase and this may be the way in which repressor normally effects repression. Moreover, the fusion strains indicate that the crucial factor for read through from *trp* to *lac* operons is the deletion of the *i* gene which specifies *mRNA* for repressor. All the evidence suggests that there is a stop signal at the end of the *i* gene which normally blocks synthesis of *mRNA* beyond the *i* gene and necessitates a reinitiation of transcription at the promoter site.

In a second paper, Reznikoff and Beckwith (*ibid.*, 215) and independently Bhorjee, Fowler and Zabin (*ibid.*, 219) report that deletion mutants which lack the *i*, *p* and most of the *o* genes can produce high concentrations of β -galactosidase. This implies of course that the operator *o* gene is not part of the structural gene for β -galactosidase and although the *o* gene is probably transcribed there is now no reason to believe that the RNA produced is ever translated.

PHARMACOLOGY

Drugs in Discussion

from a Correspondent

THE fourth international congress on pharmacology in Basle (July 13–18) was a sizable affair, with an attendance of about three thousand. There were numerous free communications and symposia, some of which were particularly good—for example those led by A. Katchalsky (Rehovot) on membrane specificities for biological transport, and by B. L. Vallee (Boston) on active sites and metalloenzymes. Two innovations were the holding of “trigger meetings”, and the lumping together of most of the free communications into large discussion groups; of the second, the less said the better. The intention of the trigger meetings was that an invited lecturer should give a short account of some unconventional approach he had brought to some problem, and this approach should then be put into critical perspective by invited discussants. This resulted in several extremely good sessions, for example, that led by J. A. Bevan (Los Angeles) on neuro-muscular transmission in blood vessels, and the experiment may well survive.

Events attended by all participants were limited to the four main lectures. In the first, J.-P. Changeux (Institut Pasteur, Paris) talked about allosteric enzymes under the title “molecular aspects of drug-receptor interaction”. The interaction of allosteric effectors with regulatory sites on proteins causes a conformational change in the protein, and it is possible that receptors for drugs and neuro-transmitters in cell membranes also produce their effect through a conformational change in the membrane, where, as he stressed, the situation is much more complex. He dealt in some detail with work from several laboratories on the question of subunit interactions in soluble proteins, but only touched on the possible relevance